

# Acknowledgements

I need to thank

- Chenyan Wu, Alex Ghitza, Chengjing Zhang
- Bowan Hafey, Oliver and Fei
- Miscellaneous lecturers

for the math that they taught me.

Thanks to my bros for being bros. Thanks to Fei Peng for the thesis template. Arun Ram for helping track down the name for parabolic.

# Contents

# Notation

# Chapter 1

## Classical Groups

We will recall a small amount of the theory of linear algebraic groups to fix conventions, for a more detailed treatment one should consult the litany of sources on this matter: For a full treatment see [?][?][?][?]. Excellent example computations can also be found in [?][?][?][?]. Or for a brief brush up on the main facts consult [?, I.I.1]. The purpose of this section is to treat the classical groups and more specifically  $\mathrm{Sp}_{2n}$  as an example and work out some of the details of the general theory, in order to "get our hands dirty" and have some familiarity with this object, to become fundamental in what follows. Because this theory is made up of simple ideas that can often be obscured by the generality we will make several restrictive assumptions for ease of exposition.

### 1.1 Definition

An algebraic group is for us a group scheme that is reduced, of finite type and defined over a field. A linear algebraic group (LAG) is an affine algebraic group. By [?, 2.3.7(i)] then every LAG is (isomorphic) to a (Zariski) closed subgroup of  $\mathrm{GL}_n$ . A group scheme is called linear if it is (isomorphic to) a closed subgroup for  $\mathrm{GL}_n$ . Hence every linear algebraic group is a *linear* algebraic group. Moreover it is a basic fact that a Zariski closed sub-scheme of an affine scheme is affine [?, II.5.T3].

As Milne points out [?, Abstract] these are supposed to be matrix groups defined by polynomials, which are somehow the natural combinations of symbols that matrix multiplication will lead to. This means that they come with the powerful but cumbersome (for the beginner) technology of algebraic geometry. In particular one must be adept at moving between the following equivalences

**Theorem** ([?, II.6, III.4]). *For  $k$  a field then the following categories are equivalent*

- *Group objects in  $\mathrm{Alg}_k^{\mathrm{opp}}$*
- *Representable (in the category of groups) functors  $\mathrm{Alg}_k \rightarrow \mathrm{Group}$*
- *Group object in the category of affine schemes over  $k$*
- *Commutative Hopf algebras*

The exact groups that an author might mean by classical may vary. Here we will follow [?, §13] or indeed many other places as defining them as the automorphisms of a vector space with a

bilinear form. First let  $V$  be an  $F$  vector space with a bilinear form  $\langle, \rangle$ . An automorphism of this form is a map  $\alpha \in \text{Aut}(V)$  such that

$$\langle \alpha(x), \alpha(y) \rangle = \langle x, y \rangle$$

Therefore we can consider the space of automorphisms of  $\text{Aut}(V, \langle, \rangle)$ . This space, depending on the properties of the bilinear form will define our classical groups.

If the form is degenerate,  $\forall x, y \langle x, y \rangle = 0$  then we define

$$\text{GL}(V) := \text{Aut}(V, \langle, \rangle) = \text{Aut}(V)$$

If the form is non-degenerate and symmetric  $\forall x, y \langle x, y \rangle = \langle y, x \rangle$  then we define

$$\text{O}(V) := \text{Aut}(V, \langle, \rangle)$$

Finally if the form is non-degenerate and skew symmetric  $\forall x, y \langle x, y \rangle = -\langle y, x \rangle$  then

$$\text{Sp}(V) := \text{Aut}(V, \langle, \rangle)$$

There are the further classical groups given by the determinant one subgroups,  $\text{SL}(V), \text{SO}(V)$  respectively ( $\text{Sp}(V)$  one can show already implies that the determinant is one). Moreover every  $F$  algebra is an  $F$  vector space and so we have defined a functor from  $F$ -algebras to groups. This is what we will refer to as "classical groups" although we should note the omission of what the unitary groups that would usually be included.

Another way of motivating the importance of these groups is to consider the classification theory of the split reductive groups. Reductive is a representation theoretic condition that means, worse than semi-simple but still manageable, for more detail see [?, 22.138], this is only for motivation. (Split) Reductive groups over fields have a classification in terms of root datum [?, 22.48] which is a strong parallel of the classification of semi-simple Lie algebras, and in fact is proved largely by bootstrapping from that theory to a more general setting. These correspond to Dynkin diagrams that are listed in [?, Appendix A,B] and of which there are four infinite families, which correspond to the classical groups (plus the unitary groups). This is in the way of motivating the notion of classical groups as "almost all" reductive groups.

## 1.2 Subgroups

From now on we restrict to split reductive LAG because these are the natural adjectives for our classical groups over a number field (in particular characteristic 0)

Subgroups with special properties allow us to reduce and break up problems into smaller ones. Here we will briefly review and compute some examples of special subgroups, with a particular eye on those of  $\text{Sp}$ . The point of these subgroups is two fold. Some of them will help us perform "induction" from smaller simpler groups to larger ones. Others are there essentially as a part of the combinatorial data that classifies the groups we are working with. In particular we will have

the decomposition's (not direct)

$$G(\mathbb{A}) = M(\mathbb{A})U(\mathbb{A})K = T(\mathbb{A})U(\mathbb{A})K$$

### 1.2.1 Parabolics, Levis and Unipotents

Parabolic subgroups have two equivalent formulations, both useful.

Do I need to give a redefinitions?

**Definition 1.1.** A subgroup  $P \subseteq G$  is called parabolic if the following equivalent conditions hold

- $G/P$  is a complete variety
- $P$  contains a Borel (see below)

Completeness is the algebro-geometric analogue of compact, which is always a desirable property. The fact that they contain a Borel gives us an algebraic "parametrisation" of these subgroups, in the case of the classical groups through the use of flags.

**Example** ( $\mathrm{GL}_n$ ). A flag for  $F^n$  is a sequence of subspaces

$$0 \subset W_1 \subset \cdots \subset W_r = F^n$$

note the strict inclusion. When  $n = r$  it is a complete flag.  $\mathrm{GL}_n(F)$  acts on a flag component wise i.e.

$$g.(W_1, \dots, W_r) := (g.W_1, \dots, g.W_r)$$

If  $F^n$  is given the standard basis of  $e_i = (\delta_i^j)_j$  then the standard (complete) flag is

$$0 \subset Fe_1 \subset Fe_1 \oplus Fe_2 \subset \cdots \subset \oplus_i Fe_i = F^n$$

check that this is an isomorphism, it is a bijection but why inclusion still hold?

From now on we consider only subflags of the standard flag.

**Remark.** This is because the stabiliser of a flag is always conjugate to a stabiliser of a sub-flag of the standard flag. i.e. stabilisers of flags are up to conjugacy stabilisers of the standard flag.

It is worth noting that what we do here by fixing a basis of  $F$  is the same as fixing a Borel and then considering only standard parabolics. Thus when we are talking about standard parabolics we are really working up "up to conjugacy".

Stabilisers of such subflags are "staircases":

$$\begin{pmatrix} A_{11} & & * \\ & \ddots & \\ & & A_{rr} \end{pmatrix}$$

**Example.** Consider the flag

$$0 \subset Fe_1 \subset Fe_1 \oplus Fe_2 \oplus Fe_3 \subset F^4$$

lets find the stabiliser under the action of  $\mathrm{GL}_4(F)$ . We need to send  $Fe_1$  to itself

$$\forall x \in F, \begin{pmatrix} a & b & c & d \\ e & f & g & h \\ i & j & k & l \\ m & n & o & p \end{pmatrix} \begin{pmatrix} x \\ \\ \\ \end{pmatrix} = \begin{pmatrix} ax \\ ex \\ ix \\ mx \end{pmatrix} \in F \begin{pmatrix} 1 \\ \\ \\ \end{pmatrix}$$

and so  $e = i = m = 0$ . Next we need to send  $Fe_1 \oplus Fe_2 \oplus Fe_3$  to itself (as a subspace)

$$\forall x, y, z \in F, \begin{pmatrix} a & b & c & d \\ e & f & g & h \\ i & j & k & l \\ m & n & o & p \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ \end{pmatrix} = \begin{pmatrix} ax + by + cz \\ ex + fy + gz \\ iz + jy + kz \\ mx + ny + oz \end{pmatrix} \in \mathrm{Span}_F \left\{ \begin{pmatrix} 1 \\ \\ \\ \end{pmatrix}, \begin{pmatrix} \\ 1 \\ \\ \end{pmatrix}, \begin{pmatrix} \\ \\ 1 \\ \end{pmatrix} \right\} = \begin{pmatrix} * \\ * \\ * \\ \end{pmatrix}$$

So we conclude that  $m = n = o = 0$ . Thus we are left with elements in  $\mathrm{GL}_4(F)$  that look like

$$\begin{pmatrix} * & * & * & * \\ & * & * & * \\ & & * & * \\ & & & * \end{pmatrix}$$

So what is happening here, for each peice of the flag you complete the diagonal that the basis vectors are in into the box that they span and you say that everything directly below that box must be zero.

$$\begin{pmatrix} 1 & & & \\ & \ddots & & \\ & & 1 & \\ & & & \end{pmatrix} \xrightarrow{\text{span}} \begin{pmatrix} \boxed{\begin{matrix} * & & * \\ & \ddots & \\ * & & * \end{matrix}} \\ \end{pmatrix} \xrightarrow{\text{delete}} \begin{pmatrix} \boxed{\begin{matrix} * & & * \\ & \ddots & \\ * & & * \end{matrix}} \\ \boxed{\begin{matrix} & & & \\ & & & \\ & & & \end{matrix}} \end{pmatrix}$$

It is a fact that these are the parabolic subgroups of  $\mathrm{GL}$  up to conjugacy, [?, Exercise 3.2.16, 6.2.11][?]

Parabolics also have the nice property that they split into a semi-direct product where one of the factors is a reductive group  $M$ . For this recall the definition

**Definition 1.2.** A subgroup is unipotent if all its elements are unipotent. The maximal closed, connected, unipotent subgroup  $U \subseteq G$  is the unipotent radical of  $G$ .

Then we have the following fact / definition:

**Lemma** ([?] 11.22). *There is a split exact sequence*

$$0 \rightarrow U \rightarrow P \rightarrow M \rightarrow 0$$

where  $U$  is the unipotent of  $P$ , and  $M$  is a reductive group known as a Levi (unique up to conjugacy).

**Example.** *The staircase*

$$\begin{pmatrix} A_{11} & & * \\ & \ddots & \\ & & A_{rr} \end{pmatrix}$$

has unipotent  $N$  and Levi  $M$  given by

$$U = \begin{pmatrix} I_{11} & & * \\ & \ddots & \\ & & I_{rr} \end{pmatrix}, \quad M = \begin{pmatrix} A_{11} & & 0 \\ & \ddots & \\ & & A_{rr} \end{pmatrix}$$

Then

$$P = M \ltimes U$$

Thus doing things on a parabolic allows us to induce said actions up to the whole group, whilst maintaining the nice property of being reductive.

**Remark** (Bad Etymology). *The origin of the name parabolic is a mystery. Borel in his history [?, VI.§2] attributes it to R. Godement in [?]. Godement conjectures that the quotient  $G(\mathbb{A})/G(\mathbb{Q})$  is compact if and only if every element of  $G(\mathbb{Q})$  is semi-simple, as is the case in classical groups. He says that*

this is probably known

*Lorsque n'est pas compact, il est non moins facile de conjecturer qu'on doit pouvoir définir quelque chose d'analogue aux classiques "pointes paraboliques", lesquelles doivent correspondre à des sous-groupes unipotents non triviaux de  $G_{\mathbb{Q}}$*

which roughly (google) translates to that one can also conjecture that non-trivial unipotent elements should correspond to "parabolic points" in a fundamental domain.

In the case of modular forms the fundamental domain is  $\mathcal{H} = \mathrm{SL}_2(\mathbb{R})/\mathrm{SO}_2(\mathbb{R})$  (using orbit stabiliser theorem). We have the classification of elements of  $\mathrm{SL}_2(\mathbb{R}) - \{\pm 1\}$  as in [?, 3.5] via their trace

$$g \text{ is of type } \begin{cases} \text{Elliptic} & \frac{1}{2}|\mathrm{tr}(g)| < 1 \\ \text{Parabolic} & \frac{1}{2}|\mathrm{tr}(g)| = 1 \\ \text{Hyperbolic} & \frac{1}{2}|\mathrm{tr}(g)| > 1 \end{cases}$$

Being parabolic is equivalent to having eigenvalue 1 hence by the Jordan decomposition we know that parabolics in  $\mathrm{SL}_2$  are conjugate (over  $\mathbb{C}$ ) to

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Clearly the standard parabolic

$$\begin{pmatrix} a & b \\ & a^{-1} \end{pmatrix} \subseteq \mathrm{SL}_2(\mathbb{R})$$

contains these matrices, and moreover all parabolics are conjugate to this parabolic. Hence all parabolic elements are contained in a parabolic subgroup. This classification it seems relies entirely on the aesthetic connection with the classification of the sections of conics via eccentricity.



To connect this to Godement's concept we have two facts from classical geometry. Proper parabolic subgroups of  $\mathrm{SL}_2(\mathbb{R})$  can be realised as the stabilisers of lines in  $\mathbb{R}^2$  under the standard action of  $\mathrm{SL}_2$  on  $\mathbb{R}^2$  [?, 2.6] and moreover some an element of  $\mathrm{SL}_2(\mathbb{R})$  is parabolic if and only if it has one fixed point on  $\partial\bar{\mathcal{H}}$  and none on  $\mathcal{H}$  [?, 3.5].

The take away is that perhaps the folklore of the name being for "para-Borelic", as in kind of a Borel, is probably a better way of thinking of them.

### $\mathrm{Sp}_{2n}$

The case of  $\mathrm{Sp}_{2n}$  is very similar to that of  $\mathrm{GL}_n$ , but we will use this later and so we present the details here. Following [?] and the very explicit calculations in [?, §8]. We let  $(V, \langle, \rangle)$  be a symplectic space, hence  $\mathrm{Sp}(V)$  is the automorphisms preserving the form. A subspace is said to be isotropic if the form is constantly zero on it (in both variables). A flag is isotropic if the proper subspaces in it are isotropic subspaces. A maximal isotropic flag is one with exactly  $n$  components (we elaborate later). The action of  $\mathrm{Sp}$  preserves isotropic flags i.e. it sends an isotropic flag to an isotropic flag. Stabilisers of isotropic flags give parabolics of  $\mathrm{Sp}$  and moreover all parabolics arise in this way (see the above exercises in Springer).

**Example.** First we remind ourselves why the two notions of the symplectic group agree

$$\mathrm{Sp}_{2n} = \{M \in \mathrm{GL}_{2n} : \forall a, b \quad \langle Ma, Mb \rangle = \langle a, b \rangle\}$$

if we make the form into a matrix (by setting the entries to be  $A_{ij} = \langle e_i, e_j \rangle$ ) then we get

$$\forall a, b \quad \langle Ma, Mb \rangle = (Ma)^T A (Mb) = a^T M^T A M b = a^T A b$$

and because this is so for all  $a, b$  we get that

$$M^T A M = A$$

So if we fix a basis of  $V$  such that the form is given by the matrix

$$\begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$$

then we can see what the form does on the standard basis vectors

$$(e_{n+1})^T \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} e_{n+1} = (e_{n+1})^T \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = (e_{n+1})^T \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = 1$$

$$(e_1)^T \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} e_1 = (e_1)^T \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = (e_1)^T \begin{pmatrix} -1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = -1$$

hence

$$\begin{aligned} (e_1 + e_{n+1})^T \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} (e_1 + e_{n+1}) &= (e_1 + e_{n+1})^T \left( \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} e_1 + \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} e_{n+1} \right) \\ &= (e_1 + e_{n+1})^T (-e_1 + e_{n+1}) \\ &= 0 \end{aligned}$$

so the vectors  $e_i^+ := e_i + e_{n+i}$  give a basis for an  $n$  dimensional isotropic subspace. In our reference it is stated more abstractly that an  $n$  dimensional isotropic subspace is maximal and that any flag of isotropic subspaces can be considered as a subflag of a complete flag of a maximal isotropic subspace (it is believable from inspecting the matrix of the form that if your space has a dimension of greater than  $n$  then it cannot be isotropic).

This might all be wrong. The book is giving something different.

Now we have a maximal isotropic flag we can consider subflags and find their stabiliser. Lets look at  $\mathrm{Sp}_4$  and the following flag

$$0 \subset Fe_1^+ \subset Fe_1^+ \oplus Fe_2^+ \subset F^4$$

If we change our basis to these  $e_i^+$  then we can re-use our GL computations to see that the stabiliser is matrices in  $\mathrm{Sp}$  of the form (in this basis)

$$\begin{pmatrix} * & * & * & * \\ & * & * & * \\ & & * & * \\ & & * & * \end{pmatrix}$$

**Remark.** Note that the fact that we change basis here doesn't change that the subspaces are isotropic, this is basis independent. What it does change is the representation of the form, it will no longer be represented by the matrix given above.

Because these are in  $\mathrm{Sp}$  we can find more relations among the entries than in the GL case, we will pursue this further in the maximal case below.

In particular maximal parabolics of  $\mathrm{Sp}$  are stabilizers of *minimal* (non-trivial flags), i.e. stabilisers of non-zero isotropic subspaces.

$$0 \subset V_\ell \subset V$$

where  $V_\ell = \text{span}_F(e_1, \dots, e_\ell)$ . Then the stabilizer is

$$\begin{pmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{pmatrix}$$

with the sizes of the diagonal blocks being (these numbers square)

$$\begin{pmatrix} \ell & * & * & * \\ 0 & n - \ell & * & * \\ 0 & * & \ell & * \\ 0 & * & * & n - \ell \end{pmatrix}$$

these sizes clearly determine the sizes of the rest of the matrix. This has Levi

$$\begin{pmatrix} A & & & \\ & a & & b \\ & & (A^T)^{-1} & \\ & c & & d \end{pmatrix}$$

such that  $A \in \text{GL}_\ell(F)$  and

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Sp}_{2(n-\ell)}(F)$$

and unipotent

$$\begin{pmatrix} 1 & * & * & * \\ & 1 & * & \\ & & 1 & \\ & & * & 1 \end{pmatrix}$$

with relations among the entries.

### 1.2.2 Borel and Torus

One may find it helpful to understand these subgroups to understand the analogous story for Lie groups and their classification [?], however it is not necessary.

**Definition 1.3.** A split torus is an algebraic group that is isomorphic to some products of  $\text{GL}_1^b$ .

**Example** (Bad Etymology).  $\text{GL}_1^2$  is a split torus. Notice that

$$\text{GL}_1^2(\mathbb{C}) = \mathbb{C}^* \times \mathbb{C}^*$$

is isomorphic as abstract groups to  $U(1) \times U(1)$  which when  $U(1)$  is realised as  $\{z \in \mathbb{C} : |z| = 1\}$  is topologically equivalent to  $\mathbb{T}^2 = \mathbb{S}^1 \times \mathbb{S}^1$  which is a torus. Note that it is clear that

$$\text{GL}_1^2(\mathbb{C}) \not\cong T^2$$

as topological groups, as the right hand side is compact whilst the left is not.

Perhaps a more compelling reason to call these Tori is that they play the same role in the classification as the genuine tori in the theory of Lie groups.

example maybe idk

**Definition 1.4.** A Borel is a maximal closed solvable connected subgroup of  $G$ .

**Example.** The standard Borel of  $\mathrm{GL}_n$  is the one given by upper triangular matrices. If  $n$  is even and one intersects this with  $\mathrm{Sp}_{2(\frac{1}{2}n)}$  then we get the standard Borel of  $\mathrm{Sp}_{2(\frac{1}{2}n)}$ .

A Borel can be considered to be a parabolic that is minimal with respect to inclusion. The maximal tori then form the Levis of these parabolics. In particular for a Borel  $B$  we have that

$$B = TU$$

for a maximal torus  $T$  and unipotent  $U$ .

We saw that fixing a minimal parabolic is like fixing a basis in the case of classifying parabolics of classical groups. If a Borel  $B$  is fixed, then a parabolic containing this Borel  $B \subseteq P$  is called standard, the unique Levi of a standard parabolic containing this Borel is called the standard Levi. This gives some intuition as to their importance in the classification of reductive groups via root datum[?, 8], as they correspond to fixing a collection of simple roots. Above also suggests that Borels are "the same data" as maximal tori from which the root datum is constructed.

ref

**Example.**

what is easy to find is definitions, what is hard to find is long drawnout matrix algebra that has been latexed.

maybe put the whole thing in the appendix.

### 1.2.3 Maximal Compact Subgroups

Let our group be defined over the global field  $k$ . We will often need to fix a maximal compact subgroup  $K \subseteq G(\mathbb{A})$ . These are not unique and as such when fixing one it can be arranged to have many nice properties [?, I.1.4]. In particular if we have a group  $G$  and a fixed Borel  $B$ :

fix a global convention

- First require that

$$K = \prod_{\nu} K_{\nu}$$

where the product is over all places of  $k$  and  $K_{\nu} \subseteq G(k_{\nu})$  is maximal compact.

But is the converse true

- For almost all places  $\nu$  of  $k$   $G(\mathcal{O}_{k_{\nu}})$  is defined and is maximal compact in  $G(k_{\nu})$  hence we can require  $K_{\nu} = G(\mathcal{O}_{k_{\nu}})$  at these places.
- We require

$$G(\mathbb{A}) = B(\mathbb{A})K$$

- For every standard parabolic  $P = MU$  we have that

$$P(\mathbb{A}) \cap K = (M(\mathbb{A}) \cap K)(U(\mathbb{A}) \cap K)$$

and  $M(\mathbb{A}) \cap K$  is a maximal compact subgroup of  $M(\mathbb{A})$ .

It is in terms of the third property that we like to think of the maximal compact subgroup, it is the complimentary piece of the Borel. Moreover the fourth property should be thought of as a condition that the maximal compact subgroups are well behaved with the way that we are moving between the bigger and smaller reductive groups.

**Example** ( $\mathrm{GL}_n(\mathbb{A}_{\mathbb{Q}})$ ). *It is a classical result that the maximal compact subgroup of  $\mathrm{GL}_n(\mathbb{R}) = \mathrm{GL}_n(\mathbb{Q}_{\infty})$  is the orthogonal group  $O(\mathbb{R})$ . By [?, II.IV.A1] we have that  $\mathrm{GL}_n(\mathbb{Q}_p)$  for  $p < \infty$  is  $\mathrm{GL}_n(\mathbb{Z}_p)$  and hence the maximal compact subgroup of  $\mathrm{GL}_n(\mathbb{A})$  is the product*

$$K = O(\mathbb{R}) \times \prod_p \mathrm{GL}_n(\mathbb{Z}_p) = O(\mathbb{R}) \times \mathrm{GL}_n(\hat{\mathbb{Z}})$$

### 1.3 Coverings

We are also be interested in certain covering groups of these LAG's. In particular [?, I.1.1] we will be intereseted in  $\mathbf{G}$  some topological group given as a finite central cover of  $G(\mathbb{A})$ . If  $\mathrm{pr} : \mathbf{G} \rightarrow G(\mathbb{A})$  is the projection then to the subgroups listed above we can associate their "lifts" (preimages under  $\mathrm{pr}$ ).

**Example** (Metaplectic Group). *If  $G = \mathrm{Sp}_{2n}$  then there is a unique non-trivial double cover*

$$0 \rightarrow \mu_2 = \{\pm 1\} \rightarrow \mathrm{Mp}_{2n} \rightarrow \mathrm{Sp}_{2n} \rightarrow 0$$

*such group extensions are classified by their second group cohomology, for this extension the relevant cocycle is called the Rao cocycle [?], c. As a set we can think of*

$$\mathrm{Mp}_{2n} = \mathrm{Sp}_{2n} \times \mu_2$$

*with the group operation*

$$(a, b)(x, y) = (ax, byc(a, x))$$

*There is a rich history and representation theory of this group, which we make no pretense of understanding, however some hints can be found in [?] and the references therein.*

There is an analogue of the Langlands program being developed for such groups a nice introduction to which can be found in [?].

Best natural question  
or not these things  
the structure of  
t paper mentions  
lity. Look into it.

## Chapter 2

# Automorphic Representations

We will omit the details of the definition of automorphic form, for the uninitiated the references that will be most helpful are [?, I.II][?] for the general theory, we will follow the notation developed in [?] as it is somewhat standard. For the connection to classical modular forms there is [?][?][?][?]. We will discuss some of the details of their representation theory because it is both subtle and needed later. In particular we want to draw attention to what we think of as the "non-algebraic" nature of the representation theory.

### 2.1 $(\mathfrak{g}, K)$ -Modules

### 2.2 Hecke Algebra

### 2.3 Automorphic Representations

#### 2.3.1 Cuspidal Representations

We will recal the definition of a cusp form in the next chapter.

reference the definition  
next chapter

#### 2.3.2 Tensor Products of Representations

Chengjing example of  
subspaces

### 2.4 Eisenstein Series

[?], [?]

### 2.5 Spectral Decomposition

#### 2.5.1 Definition and Role

This is another one of the tools that can be used to compartmentalise problems in automorphic forms, by dealing with representations that appear in different parts of the spectrum.

**2.5.2 The Decomposition of the Spectrum****2.5.3 Residual Representations of  $\mathrm{GL}_n$** **2.6 L-Functions****2.6.1 In General****2.6.2 Standard L-Functions for Classical Groups****2.6.3 L-Functions of Covering Groups**

## Chapter 3

# Constant Terms

Here we will explain the role of the constant term in our calculation of poles.

### 3.1 Definition and Role

The constant term is an operation defined on a large class of functions and is supposed to generalise the constant term of a fourier expansion, we will see this later, although one may consult [?, 1.6] for some examples as well. In particular [?, I.2.6] give the definition as follows: We consider  $P = MU$  a standard parabolic of  $G$  and  $\varphi : U(k) \backslash \mathbf{G} \rightarrow \mathbb{C}$  a measurable and locally  $L^1$  function then its constant term along  $P$  is

$$\begin{aligned}\varphi_P : U(\mathbb{A}) \backslash \mathbf{G} &\rightarrow \mathbb{C} \\ \varphi(g)_P &:= \int_{U(k) \backslash U(\mathbb{A})} \varphi(ug) du\end{aligned}$$

which inherits many of the properties of  $\varphi$  such as smoothness and moderate growth. For instance if  $\varphi$  is an automorphic form on  $\mathbf{G}$  then its constant term is an automorphic form on  $M$  [?, 6.5].

This allows us to recall the definition of cuspidal automorphic forms or "cusp forms". Let  $\varphi$  be an automorphic form on  $U(\mathbb{A})M(k) \backslash \mathbf{G}$  for  $P = MU$  a standard parabolic. Then  $\varphi$  is cuspidal if for all standard parabolics  $P' \subset P$  we have that

$$\varphi_{P'} = 0$$

Cusp forms have a central role in the theory of automorphic forms, this is for several reasons. They appear historically as interesting examples such as the Ramanujan tau function, by a theorem of Ribet [?, T2.3] the Galois representation associated to a cusp form is irreducible and finally they just make formulas easier to work with.

The constant term itself is of central importance. One reason for this is that it controls the growth of the automorphic forms. More precisely we have the following theorem [?, I.4.10]

**Theorem.** *Let  $P = MU$  be a standard parabolic of  $G$ ,  $V_P \subseteq \mathcal{A}_0(U(\mathbb{A})M(k) \backslash G)$  a finite dimensional subspace,  $\Gamma_P \subseteq X_M$  a compact subset and  $N_P$  an integer. Let  $n \in \mathbb{Z}$ ,  $D \subseteq \mathbb{C}^n$  open and connected,  $f : D \rightarrow \mathbb{C}$  holomorphic and not identically zero. Let  $D' = \{x \in D : f(x) \neq 0\}$ .*



If  $\varphi : D' \rightarrow L_{loc}^2(G)$  is a function such that for all  $z \in D'$ ,  $\varphi(z) \in \mathcal{A}((V_P, \Gamma_P, N_P)_{P_0 \subset P \subset G})$  and for all  $P$

$$\begin{aligned}\psi_P : D' &\rightarrow L_{loc}^2(G) \\ z &\mapsto \psi_P(z) = \varphi_P^{cusp}(z)\end{aligned}$$

is a holomorphic function.

Then  $\varphi$  can be analytically continued to a holomorphic function on  $D$  iff  $\psi_P$  can be continued to a holomorphic function on  $D$ .

Unpacking this a bit we see that an Eisenstein series satisfies the hypothesis on  $\varphi$  by [?, IV.1.9], we think of  $D'$  as the positive open cone on which the Eisenstein series converges, and then  $D = \mathbb{C}^n \setminus S$  as the rest of the cuspidal datum  $\mathfrak{P}$  on which the Eisenstein series is holomorphic ( $S$  is its set of singularities). Therefore the domain on which the Eisenstein series is holomorphic is the same as the domain on which (roughly) its constant term is holomorphic.

## 3.2 Integration Lemmas

**Theorem.** If  $G$  is a locally compact Hausdorff group with a left Haar measure  $\mu$  and if  $\chi : G \rightarrow \mathbb{C}^\times$  is a non-trivial character on  $G$ , then

$$\int_G \chi(g) d\mu(g) = 0.$$

*Proof.* Pick an element  $h$  of  $G$  such that  $\chi(h) \neq 1$ . The equation above then follows from

$$\int_G \chi(g) d\mu(g) = \int_G \chi(hg) d\mu(g) = \int_G \chi(h)\chi(g) d\mu(g) = \chi(h) \int_G \chi(g) d\mu(g). \square$$

Integrating trivial characters gives the volume of the measure space which we typically normalize to be one.

**Theorem** ([?] 5.2, [?] Thm 2.49). Let  $H \leq G$  be a closed subgroup. If  $H \backslash G$  has a right  $G$  invariant measure (iff their modular functions agree on  $H$ ) then the integral is unique up to scalar, namely for a given Haar measures  $dh$  on  $H$  and  $dg$  on  $G$  there is a unique invariant measure  $dq$  on  $H \backslash G$  such that for all  $f \in C_c^0(G)$

$$\int_{H \backslash G} \int_H f(hq) dh dq = \int_G f(g) dg$$

Note that this quotient may not be a group, because  $H$  is not required to be normal.

## 3.3 Constant Terms of Eisenstein Series

This computation forms the heart of a well known theorem, [?, Prop 10.4.2][?, II.1.7][?, 6.2]. I give more detail here than I could find anywhere else.

Notice that the Eisenstein series has a full  $G(k)$  invariance and so we can take its constant terms along any standard parabolic.

### 3.3.1 In General

We will use the following Lemmas to give a simplified expression of the constant term of an Eisenstein series. First fix  $P = MN$  and  $P' = M'N'$  two standard parabolics of suitable group  $G$  over  $F$ , with  $E(x, \varphi, \lambda)$  defined via parabolic induction from  $P$ .

**Lemma.**

$$P(F) \setminus G(F) \cong \coprod_{w \in W_{M'} \setminus W_G / W_M} P'(F) \cap wP(F)w^{-1} \setminus P'(F)$$

**Proof.** Consider the Bruhat decomposition:

$$G(F) = \coprod_{w \in W_{M'} \setminus W_G / W_M} P(F)w^{-1}P'(F)$$

then because the action of  $P(F)$  keeps the disjoint sets disjoint we can move the quotient through and get

$$P(F) \setminus G(F) = \coprod_w P(F) \setminus P(F)w^{-1}P'(F)$$

so we analyse the summands, by the second isomorphism theorem we have a bijection

$$P(F) \setminus P(F)w^{-1}P'(F) \cong P(F) \cap P'(F) \setminus w^{-1}P'(F)$$

now if  $[w^{-1}p] \in P(F) \cap P'(F) \setminus w^{-1}P'(F)$  then its  $pw^{-1}p'$  for some  $p \in P(F) \cap P'(F)$  and hence multiplying by  $w$ , in particular an isomorphism, gives  $wpw^{-1}p' \in wP(F)w^{-1} \times P'(F)$  and so

$$w(P(F) \cap P'(F) \setminus w^{-1}P'(F)) \cong wP(F)w^{-1} \cap P'(F) \setminus P'(F)$$

**Lemma.** Let  $m', n' \in M'(F) \times N'(F)$  then

$$m'n' \in wP(F)w^{-1} \iff m' \in wP(F)w^{-1} \text{ and } n' \in (m')^{-1}wP(F)w^{-1}m'$$

**Proof.** The forward implication is stated in [?], the converse follows from some algebra: First let  $m' = wp_1w^{-1}$  and  $n' = (m')^{-1}wp_2w^{-1}m'$  then

$$\begin{aligned} m'n' &= (wp_1w^{-1})^{-1}wp_2w^{-1}wp_1w^{-1} \\ &= wp_1^{-1}w^{-1}wp_2w^{-1}wp_1w^{-1} \\ &= wp_1^{-1}p_2p_1w^{-1} \in wP(F)w^{-1} \end{aligned}$$

Taking the contrapositive of this lemma will be used below. This is because our sums will be over quotients like  $A \setminus B$  and therefore summing over the "elements" in  $B$  that are not in  $A$ ; by our lemma would be the same as summing over two different such quotients. Now consider the computation:

I have lost the lambda  
below computation

$$\begin{aligned}
E_{P'}(x, \varphi, \lambda) &= \int_{N'(F) \backslash N'(\mathbb{A})} E(nx, \varphi, \lambda) dn \\
([N'] &:= N'(F) \backslash N'(\mathbb{A})) = \int_{[N']} \sum_{\delta \in P(F) \backslash G(F)} \varphi(\delta nx) dn \\
(\text{Lemma 1}) &= \int_{[N']} \sum_{\delta \in \coprod_{w \in W_{M'} \backslash W_G / W_M} P'(F) \cap w P(F) w^{-1} \backslash P'(F)} \varphi(\delta nx) dn \\
&= \sum_{w \in W_{M'} \backslash W_G / W_M} \int_{[N']} \sum_{p' \in P'(F) \cap w P(F) w^{-1} \backslash P'(F)} \varphi(w^{-1} p' nx) dn \\
(\text{Lemma 2}) &= \sum_w \sum_{m' \in M'(F) \cap w P(F) w^{-1} \backslash M'(F)} \int_{[N']} \sum_{n' \in N'(F) \cap (m')^{-1} w P(F) w^{-1} m' \backslash N'(F)} \varphi(w^{-1} m' n' nx) dn \\
(\text{Change Var}) &= \sum_w \sum_{m'} \int_{[N']} \sum_{n' \in N'(F) \cap w P(F) w^{-1} \backslash N'(F)} \varphi(w^{-1} n' n m' x) dn \\
(\text{Unfold}) &= \sum_w \sum_{m'} \int_{N'(F) \cap w P(F) w^{-1} \backslash N'(\mathbb{A})} \varphi(w^{-1} n m' x) dn
\end{aligned}$$

The change of variables is  $(m', n') \mapsto ((m')^{-1} n' m', (m')^{-1} n' m')$ . Again we assume that our  $x$  is sufficiently large so all the integrals converge.

### 3.3.2 Constant Terms of Cuspidal Eisenstein Series

**Lemma (4).** *For  $w \in W_{M'} \backslash W_G / W_M$  we have that  $w^{-1} P' w \cap M$  is a standard parabolic of  $M$  with Levi  $w^{-1} M' w \cap M$  and unipotent  $w^{-1} N' w \cap M$ .*

**Proof.** This is [?, 10.4.1] stated without proof. They give the reference [?, V.4.6] which is in French. We will return to this if we have time / energy.

**Lemma (5).**

$$w^{-1} U' w \cap P = (w^{-1} U' w \cap M) (w^{-1} U' w \cap U)$$

**Proof.** [?, 10.4.1] has some decompositions, as well as the standard decomposition of  $P = MU$  I think I could prove this...

**Lemma (6).**

$$c \backslash (b \backslash a) = (bc) \backslash a$$

Continuing the computation of the constant term above, we will focus purely on the inner integral now

$$\begin{aligned}
\int_{N'(F) \cap w P(F) w^{-1} \backslash N'(\mathbb{A})} \varphi(w^{-1} n m' x) dn &= \int_{w^{-1} N'(F) w \cap P(F) \backslash w^{-1} N'(\mathbb{A}) w} \varphi(n w^{-1} m' x) dn \\
(\text{Lemma 5}) &= \int_{(w^{-1} U' w \cap M) (w^{-1} U' w \cap U) (F) \backslash w^{-1} N'(\mathbb{A}) w} \varphi(n w^{-1} m' x) dn \\
(\text{Unfold} + \text{Lemma 6}) &= \int_{(w^{-1} U'(\mathbb{A}) w \cap M(\mathbb{A})) \backslash A} \int_{w^{-1} U'(F) w \cap M(F) \backslash w^{-1} U'(\mathbb{A}) w \cap M(\mathbb{A})} \varphi(n_1 n_2 w^{-1} m' x) dn_1 dn_2
\end{aligned}$$

the first equality is the change of variables  $w^{-1} n w \mapsto n$  and  $A = (w^{-1} U'(F) w \cap U(F)) \backslash w^{-1} N'(\mathbb{A}) w$ .

Now look at the inner integral here more closely

$$\int_{w^{-1}U'(F)w \cap M(F) \backslash w^{-1}U'(\mathbb{A})w \cap M(\mathbb{A})} \varphi(n_1 n_2 w^{-1} m' x) dn_1 dn_2$$

applying Lemma (6) we see that this is a constant term for a parabolic of  $M$ , of the function  $m \mapsto \varphi(m n_2 w^{-1} m' x)$ .

**Lemma.**  $n_2 w^{-1} m' x \in K$  with variables as above.

This was all in complete generality as well. If we now assume further that the Eisenstein series was induced from a *cuspidal* automorphic representation, then  $m \mapsto \varphi(mk)$  is a cusp form and therefore this last integral will vanish whenever  $w^{-1}U'w \cap M \neq \{1\}$ , because in that case the inner integral doesn't exist (it's over a point).

### 3.3.3 Constant Term Of Eisenstein Series for Conjugate Levis

If we now assume that  $M' = wMw^{-1}$  and recall the definition of our intertwining operator we can use the following

put this in the Eisenstein section

**Lemma** ([?] II.1.7 (6)).

$$U'(k) \cap wP(k)w^{-1} = U'(k) \cap wU(k)w^{-1}$$

to see that

$$\begin{aligned} E_{P'}(x, \varphi, \lambda) &= \sum_w \sum_{m'} \int_{N'(F) \cap wP(F)w^{-1} \backslash N'(\mathbb{A})} \varphi(w^{-1}nm'x) dn \\ &= \sum_w \sum_{m'} \int_{N'(k) \cap wN(k)w^{-1} \backslash N'(\mathbb{A})} \varphi(w^{-1}nm'x) dn \\ &= \sum_w \sum_{m'} M(w, \pi)(\varphi)(x) \end{aligned}$$

In particular we can combine the conjugate and cuspidal cases to get a much simpler expression for some constant terms of some Eisenstein series.

I have mixed up my notation too much...

## 3.4 Siegel Phi Operator

Here we give an example of the constant term which connects it to the classical picture. We thank Chengjing Zhang for showing us this example, and present it here because we cannot find it in the literature. We deal only with the classical Siegel modular forms of full level and moreover are less explicit with the steps as they should be clear after exposure to the previous arguments.

give references for the interpretations of the constant term, there must be some

Because we are trying to connect this to the classical picture it is most convenient to think of things in the archimedean places, recall the way that modular forms are automorphic forms most naturally in the archimedean sense ([?, 6.2]) [?][?][?]. So for this section alone, by automorphic form we will mean automorphic forms on the archimedean places, and the constant term will be

taken only on the archimedean part: i.e. for  $f : G(\mathbb{R}) \rightarrow \mathbb{C}$  and automorphic its constant term along a parabolic of  $G$ , call it  $P = MN$ , is [?, 8.6]

$$f(x)_P = \int_{N(\mathbb{Z}) \backslash N(\mathbb{R})} f(xn) dn$$

We assume here for simplicity (and because it will apply to the examples below) that our groups are unimodular.

### 3.4.1 Siegel Modular Forms

We collect some definitions from [?] to fix notation. Let the Siegel upper half plane be defined as

$$\begin{aligned} \mathcal{H}_g &:= \{ \tau \in M_{g \times g}(\mathbb{C}) : \tau \text{ is symmetric and has positive definite imaginary part} \} \\ &\cong \mathrm{Sp}_{2g}(\mathbb{R}) / U(g) \end{aligned}$$

where the isomorphism is as analytic manifolds and

$$U(g) := \left\{ \begin{pmatrix} A & B \\ -B & D \end{pmatrix} \in \mathrm{Sp}_{2g}(\mathbb{R}) : AA^t + BB^t = 1 \right\}$$

For every  $\gamma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}_{2g}(\mathbb{Z})$  and  $\tau \in \mathcal{H}_g$  we have the action

$$\gamma.\tau = (A\tau + B)(C\tau + D)^{-1}$$

We say that a holomorphic function  $f : \mathcal{H}_g \rightarrow \mathbb{C}$  is a (classical) Siegel modular form of weight  $k$  if

$$f(\gamma.\tau) = \det(C\tau + D)^k f(\tau)$$

with the extra condition that if  $g = 1$  it must be holomorphic at  $\infty$ . Because  $\mathrm{Sp}_2 = \mathrm{SL}_2$  this is a strict generalisation of an (elliptic) modular form.

The space of Siegel modular forms of weight  $k$  and genus  $g$  is denoted  $\mathcal{M}_k(\mathrm{Sp}_{2g}(\mathbb{Z}))$ . There is a useful operator known as the Siegel Phi Operator which allows you to lift known modular forms from lower genus to higher genus [?, 5]

$$\mathcal{M}_k(\mathrm{Sp}_{2g}(\mathbb{Z})) \xrightarrow{\Phi} \mathcal{M}_k(\mathrm{Sp}_{2(g-1)}(\mathbb{Z}))$$

defined by the limit for  $\tau \in \mathcal{H}_{g-1}$

$$\Phi(f)(\tau) := \lim_{t \rightarrow \infty} f \begin{pmatrix} \tau & \\ & it \end{pmatrix}$$

in this context a cusp form is defined to be a Siegel modular form in the kernel of the Siegel  $\Phi$  operator and so it is natural to wonder if there is a constant term that is being taken here.

### 3.4.2 Automorphising

Given a Siegel modular form  $f \in \mathcal{M}_k(\mathrm{Sp}_{2g}(\mathbb{Z}))$  we can associate an automorphic form

$$\tilde{f} : \mathrm{Sp}_{2g}(\mathbb{R}) \rightarrow \mathbb{C}, \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \det(ci + d)^{-k} f\left(\frac{ai + b}{ci + d}\right),$$

where  $a, b, c, d$  are  $g \times g$  matrices such that  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{Sp}_{2g}(\mathbb{R})$ . Fix the Borel of upper triangular matrices. Now for  $1 \leq r \leq g - 1$  an integer we have the standard maximal parabolic of  $\mathrm{Sp}_{2g}$ ,  $P_r = M_r N_r$  such that

$$M_r \cong \mathrm{GL}_r \times \mathrm{Sp}_{2(g-r)}$$

**Theorem** (Zhang). *If  $f$  is a classical Siegel modular form of weight  $k$  and degree  $g$ , then*

$$\tilde{f}_{P_r}(u\gamma) = \det u^k \cdot (\Phi^r f)^\sim(\gamma) \quad (3.4.1)$$

for every element  $\gamma$  of  $\mathrm{Sp}_{2(g-r)}(\mathbb{R})$  and every element  $u$  of  $\mathrm{GL}_r(\mathbb{R})$ .

In particular

$$\tilde{f}_{P_{g-1}} \begin{pmatrix} a & 0 & b & 0 \\ 0 & 1 & 0 & 0 \\ c & 0 & d & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = (\Phi f)^\sim \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

This shows that perhaps the correct generalisation of the Siegel Phi function is just the constant term that we all know and love. It would be interesting to compare this with the work done in [?]. We could also attempt to expand this to Siegel modular forms that are vector valued or not of full level.

### 3.4.3 Base Case

The base case is very instructive, it deals with modular forms. So consider  $f$  a (elliptic) modular form of full level and weight  $k$ , which has a fourier expansion given by

$$f(z) = \sum_{n \geq 0} a_n e^{2\pi i n z}$$

Then one can verify that

$$\tilde{f} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = (ci + d)^{-k} f\left(\frac{ai + b}{ci + d}\right)$$

is an automorphic form on  $\mathrm{Sp}_2$ . The only non-trivial parabolic  $P$  is the one of upper triangular matrices, with Levi and unipotent given respectively

$$M = \begin{pmatrix} m & 0 \\ 0 & m^{-1} \end{pmatrix} \cong \mathrm{GL}_1, \quad N = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \cong \mathbb{G}_a$$

along which we can now compute the constant term

$$\begin{aligned}
 \tilde{f}_P(m) &= \int_{N(\mathbb{Z}) \backslash N(\mathbb{R})} \tilde{f}(mb) db \\
 &= \int_{\mathbb{Z} \backslash \mathbb{R}} \tilde{f} \begin{pmatrix} m & mb \\ 0 & m^{-1} \end{pmatrix} db \\
 &= \int_{\mathbb{Z} \backslash \mathbb{R}} m^k f(m^2 i + m^2 b) db \\
 &= m^k a_0
 \end{aligned}$$

We have chosen normalisation to remove the usual factor of  $1/2\pi$  in the constant term of the Fourier series. Moreover we see that

$$\Phi(f) = \lim_{t \rightarrow \infty} f(it) = \lim_{t \rightarrow \infty} \sum_{n \geq 0} a_n e^{-2\pi n t} = a_0$$

### 3.4.4 Simplifying the Constant Term

maximal parabolic

As we saw in for  $1 \leq r \leq g-1$  an integer we have the standard maximal parabolic of  $\mathrm{Sp}_{2g}$ ,  $P_r = M_r N_r$  such that

$$M_r \cong \mathrm{GL}_r \times \mathrm{Sp}_{2(g-r)}$$

which can be given the explicit matrix representations

$$m(\gamma, A) := \begin{pmatrix} A & & & \\ & a & & b \\ & & (A^t)^{-1} & \\ & c & & d \end{pmatrix}, \quad A \in \mathrm{GL}_r(F), \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{Sp}_{2(g-r)}(F)$$

and unipotent

$$n(s; h, k) := \begin{pmatrix} 1 & 0 & 0 & h \\ -k^t & 1 & h^t & s + h^t k \\ 0 & 0 & 1 & k \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad h, k \in \mathrm{Mat}_{(g-r) \times r}(\mathbb{R}) \quad s \in \mathrm{Sym}_r(\mathbb{R})$$

We have the following short exact sequence

$$1 \rightarrow \mathrm{Sym}_r(\mathbb{R}) \rightarrow N_r(\mathbb{R}) \rightarrow \mathrm{Mat}_{(g-r) \times r}(\mathbb{R}) \times \mathrm{Mat}_{(g-r) \times r}(\mathbb{R}) \rightarrow 1.$$

which we will use to unfold our integral below, for compactness we define  $H_r := \mathrm{Mat}_{(g-r) \times r}$ . We will now denote  $[G] := G(\mathbb{Z}) \backslash G(\mathbb{R})$  and compute the constant term

$$\begin{aligned}
 \tilde{f}_{P_r}(m(\gamma, A)) &= \int_{[N_r]} \tilde{f}(nm(\gamma, A)) dn \\
 &= \int_{[H_r \times H_r]} \int_{[\mathrm{Sym}_{g-r}]} \tilde{f}(n(s; h, k)m(\gamma, A)) ds d(h, k)
 \end{aligned}$$

$$= \int_{[H_r]} \int_{[H_r]} \int_{[\text{Sym}_{g-r}]} \tilde{f}(n(s; h, k)m(\gamma, A)) ds dh dk. \quad (3.4.2)$$

Now we focus on simplifying the integrand. We want an explicit form of the matrix so we can relate it back to the value of the un-lifted Siegel modular form  $f$ ; simply multiply the matrices gives, where (all rings are commutative)  $A^{-t} := (A^t)^{-1}$

$$n(s; h, k)m(\gamma, A) = \begin{pmatrix} a & 0 & b & hA^{-t} \\ -k^t a + h^t c & A & -k^t b + h^t d & sA^{-t} + h^t kA^{-t} \\ c & 0 & d & kA^{-t} \\ 0 & 0 & 0 & A^{-t} \end{pmatrix}.$$

because  $a, b, c, d \in \text{Mat}_{(g-r) \times (g-r)}$ ,  $A \in \text{Mat}_{r \times r}$  we see that the  $g \times g$  blocks that we now need to take the determinant of are the  $4 \times 4$  corners of this picture, hence the matrices below should all be in  $\mathcal{H}_g \subseteq \text{Mat}_{g \times g}$

$$\begin{aligned} \tilde{f}(n(s; h, k)m(\gamma, A)) &= \det \left( \begin{pmatrix} c & 0 \\ 0 & 0 \end{pmatrix} i + \begin{pmatrix} d & kA^{-t} \\ 0 & A^{-t} \end{pmatrix} \right)^{-k} \\ &= f \left( \left( \begin{pmatrix} a & 0 \\ -k^t a + h^t c & A \end{pmatrix} i + \begin{pmatrix} b & hA^{-t} \\ -k^t b + h^t d & sA^{-t} + h^t kA^{-t} \end{pmatrix} \right) \left( \begin{pmatrix} c & 0 \\ 0 & 0 \end{pmatrix} i + \begin{pmatrix} d & kA^{-t} \\ 0 & A^{-t} \end{pmatrix} \right)^{-1} \right) \\ &= \det \left( \begin{pmatrix} ic + d & kA^{-t} \\ 0 & A^{-t} \end{pmatrix} \right)^{-k} \\ &= f \left( \begin{pmatrix} ia + b & hA^{-t} \\ -k^t(ia + b) + h^t(d + ic) & iA + sA^{-t} + h^t kA^{-t} \end{pmatrix} \begin{pmatrix} ic + d & kA^{-t} \\ 0 & A^{-t} \end{pmatrix}^{-1} \right) \\ &= \left( \frac{\det(ic + d)}{\det(A)} \right)^{-k} \\ &= f \left( \begin{pmatrix} ia + b & hA^{-t} \\ -k^t(ia + b) + h^t(d + ic) & iA + sA^{-t} + h^t kA^{-t} \end{pmatrix} \begin{pmatrix} (ci + d)^{-1} & -(ci + d)^{-1}k \\ 0 & A^t \end{pmatrix} \right) \\ &= \left( \frac{\det(A)}{\det(ic + d)} \right)^k f \left( \begin{pmatrix} \tau & -\tau k + h \\ -k^t \tau + h^t & k^t \tau k + AA^t i + s \end{pmatrix}, \quad \tau := (ai + b)(ci + d)^{-1} \right) \end{aligned}$$

So we have shown that

$$\begin{aligned} \tilde{f}_{P_r}(m(\gamma, A)) &= \int_{[H_r]} \int_{[H_r]} \int_{[\text{Sym}_{g-r}]} \left( \frac{\det(A)}{\det(ic + d)} \right)^k f \left( \begin{pmatrix} \tau & -\tau k + h \\ -k^t \tau + h^t & k^t \tau k + AA^t i + s \end{pmatrix} \right) ds dh dk \\ &= \left( \frac{\det(A)}{\det(ic + d)} \right)^k \int_{[H_r]} \int_{[H_r]} \int_{[\text{Sym}_{g-r}]} f \left( \begin{pmatrix} \tau & -\tau k + h \\ -k^t \tau + h^t & k^t \tau k + AA^t i + s \end{pmatrix} \right) ds dh dk \end{aligned}$$



Again lets focus on this integrand  $f \begin{pmatrix} \tau & -\tau k + h \\ -k^t \tau + h^t & k^t \tau k + AA^t i + s \end{pmatrix}$  and compute its Fourier expansion, see [?, 3.4]. Recall that a symmetric matrix  $n \in \text{GL}_g(\mathbb{Q})$  is called half integral if  $2n$  is integral with even diagonal entries, then a Siegel modular form has a Fourier expansion of the form

$$f(z) = \sum_{n \text{ half integral}} a(n) e^{2\pi i \text{Tr}(nz)}$$

First the space of half integral  $g \times g$  matrices,  $\text{HI}_g$ , decomposes as a direct sum via the (additive) group isomorphism

$$\text{HI}_{g-r} \oplus \frac{1}{2} \text{Mat}_{r \times (g-r)}(\mathbb{Z}) \oplus \text{HI}_r \rightarrow \text{HI}_g, \quad (n, m, l) \mapsto \begin{pmatrix} n & m \\ m^t & l \end{pmatrix},$$

thus unfolding the (discrete) integral we get

$$f \begin{pmatrix} \tau & -\tau k + h \\ -k^t \tau + h^t & k^t \tau k + AA^t i + s \end{pmatrix} = \sum_{n \in \text{HI}_{g-r}} \sum_{m \in \frac{1}{2} \text{Mat}_{r \times (g-r)}(\mathbb{Z})} \sum_{l \in \text{HI}_r} a \begin{pmatrix} n & m \\ m^t & l \end{pmatrix} \exp \left( 2\pi i \text{Tr} \begin{pmatrix} n & m \\ m^t & l \end{pmatrix} \begin{pmatrix} \tau & -\tau k + h \\ -k^t \tau + h^t & k^t \tau k + AA^t i + s \end{pmatrix} \right)$$

because all the block sizes are compatible we can "block multiply" the inner matrices and because we are taking the trace we can forget about off diagonal entries

$$\begin{pmatrix} n & m \\ m^t & l \end{pmatrix} \begin{pmatrix} \tau & -\tau k + h \\ -k^t \tau + h^t & k^t \tau k + AA^t i + s \end{pmatrix} = \begin{pmatrix} n\tau + m(-k^t \tau + h^t) & * \\ * & m^t(-\tau k + h) + l(k^t \tau k + AA^t i + s) \end{pmatrix}$$

putting this into our Fourier expansion

$$f \begin{pmatrix} \tau & -\tau k + h \\ -k^t \tau + h^t & k^t \tau k + AA^t i + s \end{pmatrix} = \sum_n \sum_m \sum_l a \begin{pmatrix} n & m \\ m^t & l \end{pmatrix} \exp (2\pi i (\text{Tr}(n\tau) + \text{Tr}(m(-k^t \tau + h^t)) + \text{Tr}(m^t(-\tau k + h)) + \text{Tr}(l(k^t \tau k + AA^t i + s))))$$

rs from Chengjing

If we denote  $T_l := \text{Tr}(l(k^t \tau k + AA^t i + s))$  and

$$T_m := \text{Tr}(m(-k^t \tau + h^t)) + \text{Tr}(m^t(-\tau k + h)) = \text{Tr}(-mk^t \tau - m^t \tau k) + \text{Tr}(mh^t + m^t h) := T_{m,k} + T_{m,h}$$

uniformly a priori on s, well I dont know if all these sums haha

we can sub this back into our constant term

$$\begin{aligned} \tilde{f}_{P_r}(m(\gamma, A)) &= \left( \frac{\det(A)}{\det(ic + d)} \right)^k \int_{[H_r]} \int_{[H_r]} \int_{[\text{Sym}_{g-r}]} \sum_n \sum_m \sum_l a \begin{pmatrix} n & m \\ m^t & l \end{pmatrix} \exp (2\pi i (\text{Tr}(n\tau) + T_m + T_l)) \, ds \, dh \, dk \\ &= \left( \frac{\det(A)}{\det(ic + d)} \right)^k \sum_n \sum_m \sum_l a \begin{pmatrix} n & m \\ m^t & l \end{pmatrix} e^{2\pi i \text{Tr}(n\tau)} \int_{[H_r]} \int_{[H_r]} \int_{[\text{Sym}_{g-r}]} e^{2\pi i (T_m + T_l)} \, ds \, dh \, dk \end{aligned}$$

$$\begin{aligned}
&= \left( \frac{\det(A)}{\det(ic+d)} \right)^k \sum_n \sum_m \sum_l a \begin{pmatrix} n & m \\ m^t & l \end{pmatrix} e^{2\pi i \text{Tr}(n\tau)} \int_{[H_r]} \int_{[H_r]} e^{2\pi i T_m} \int_{[\text{Sym}_{g-r}]} e^{2\pi i T_l} ds dh dk \\
&= \left( \frac{\det(A)}{\det(ic+d)} \right)^k \sum_n \sum_m \sum_l a \begin{pmatrix} n & m \\ m^t & l \end{pmatrix} e^{2\pi i \text{Tr}(n\tau)} \int_{[H_r]} e^{2\pi i T_{m,k}} \int_{[H_r]} e^{2\pi i T_{m,h}} \int_{[\text{Sym}_{g-r}]} e^{2\pi i T_l} ds dh dk
\end{aligned}$$

Now we use that integrating unitary characters is very controlled and the fact that

ref the lemma

$$s \mapsto e^{2\pi i T_l}$$

is a non-trivial unitary character of  $\text{Sym}_{g-r}$  whenever  $l \neq 0$  to get that

$$\int_{[\text{Sym}_{g-r}]} e^{2\pi i T_l} ds = \begin{cases} 1, & l = 0 \\ 0, & l \neq 0 \end{cases}$$

we repeat this trick with the second integral, which enforces that  $m = 0$  and end up with

$$\tilde{f}_{P_r}(m(\gamma, A)) = \left( \frac{\det(A)}{\det(ic+d)} \right)^k \sum_{n \in \text{HI}_{g-r}} a \begin{pmatrix} n & 0 \\ 0 & 0 \end{pmatrix} e^{2\pi i \text{Tr}(n\tau)}$$

but by [?, 3.5] we know that the Fourier expansion of the Siegel Phi operator is

$$(\Phi^r f)(\tau) = \sum_{n \in \text{HI}_{g-r}} a \begin{pmatrix} n & 0 \\ 0 & 0 \end{pmatrix} e^{2\pi i \text{Tr}(n\tau)}.$$

hence

$$\begin{aligned}
\tilde{f}_{P_r}(m(\gamma, A)) &= \left( \frac{\det(A)}{\det(ic+d)} \right)^k \Phi^r(f)(\tau) \\
&= \det(A)^k (\Phi^r(f))^\sim(\gamma)
\end{aligned}$$

which concludes the proof.

□

# Chapter 4

## Poles of Residual Eisenstein Series

Our goal here is to exposit and survey the work in papers such as [?][?], we were going to generalise some of these results to metaplectic covers however we found these results in [?].

### 4.1 Their Results

[?] gave an analysis of the residual poles of Eisenstein series attached to  $\mathrm{Sp}_{2n}$ , there were some minor errors that were corrected in [?] where they give essentially the same proof; theirs however works for the other classical groups. For our purposes, the case of  $\mathrm{Sp}_{2n}$ , as a group defined over  $F$  a number field, is most relevant, and we shall therefore focus exclusively on this case, however it should be noted that this limitation in the non-covering case is artificial, although it does simplify things a little by removing some casework, and we hope also in the covering case to be able to remove it in future work.

We fix an  $n \in \mathbb{N}$  and the Borel of upper triangular matrices in  $\mathrm{Sp}_{2n}$ , then we look at partitions of  $n = r + m$ , where . Then as we saw in — there corresponds a maximal standard parabolic of  $\mathrm{Sp}_{2n}$ , which we denote  $P_r = M_r N_r$ , such that the Levi component is

$$\mathrm{GL}_r \times \mathrm{Sp}_{2m}$$

As we saw in — the space of characters  $X_{M_r}^{\mathrm{Sp}_{2n}}$  is one dimensional by the maximality of  $P_r$ . If we look at the divisors of  $r = ab$  and fix a  $\tau$ , an irreducible unitary cuspidal automorphic representation of  $\mathrm{GL}_a$ , then from — we know that  $\Delta(\tau, b)$  is a residual representation of  $\mathrm{GL}_{ab} = \mathrm{GL}_r$ . Now we take an irreducible generic cuspidal automorphic representation  $\sigma$  of  $\mathrm{Sp}_{2m}$ , and so their tensor product  $\Delta(\tau, b) \otimes \sigma$  gives a representation of  $\mathrm{GL}_r \times \mathrm{Sp}_{2m}$  and hence of the Levi  $M_r$ . We now consider the Eisenstein series attached to this representation, namely if

$$\varphi \in \mathcal{A}(N_r(\mathbb{A})M_r(F) \setminus \mathrm{Sp}_{2n}(\mathbb{A}))_{\Delta(\tau, b) \otimes \sigma}$$

then we have the Eisenstein series

$$E(\varphi, s)(g) = \sum_{\gamma \in P_r(F) \backslash \mathrm{Sp}_{2n}(F)} s \cdot \varphi(\gamma g)$$

for  $g \in \mathrm{Sp}_{2n}(F) \setminus \mathrm{Sp}_{2n}(\mathbb{A})$ . Because it is induced from the residual representation  $\Delta(\tau, b)$  we call these residual Eisenstein series. The main theorem is then a statement that after some normalisation the poles of this Eisenstein series are limited to a particular set and that they are simple. Details will be given later.

explain the  $s$  action in series section.

## 4.2 Our Results

We consider an almost identical setup but we deal with the metaplectic cover of  $\mathrm{Sp}_{2n}$ , again over a number field  $F$ ,  $\mathrm{Mp}_{2n}$ . We again fix the Borel of upper triangular matrices, consider partitions of  $n = r + m$  and look at maximal standard parabolics of  $\mathrm{Sp}_{2n}$ ,  $P_r = M_r N_r$  such that

reference the section this in.

$$M_r = \mathrm{GL}_r \times \mathrm{Sp}_{2m}$$

then if  $r = ab$  we still have that  $\Delta(\tau, b) \otimes \sigma$ , for  $\tau$  irreducible unitary cuspidal automorphic representation of  $\mathrm{GL}_a$  and  $\sigma$  irreducible generic cuspidal automorphic representation of  $\mathrm{Sp}_{2m}$ , is a representation of  $M_r$ . The difference is in the parabolic induction as we now consider

$$\varphi \in \mathcal{A}(N_r(\mathbb{A})M_r(F) \setminus \mathrm{Mp}_{2n}(\mathbb{A}))_{\Delta(\tau, b) \otimes \sigma}$$

and then the Eisenstein series is defined in the same way

$$E(\varphi, s)(g) = \sum_{\gamma \in P_r(F) \backslash \mathrm{Sp}_{2n}(F)} s \cdot \varphi(\gamma g)$$

for  $g \in \mathrm{Sp}_{2n}(F) \setminus \mathrm{Mp}_{2n}(\mathbb{A})$  and  $s \in \mathbb{C} \cong X_{M_r}^{\mathrm{Mp}_{2n}}$ .

**Lemma.** *When  $b = 1$  we have the constant term*

$$E(\varphi, s)(g)_{P_a} = \varphi(g)_{P_a} + M(\omega, \tau \otimes \sigma)(\varphi)(g)$$

fill in here as theorem ever anything that I actually checking....

## 4.3 Setup

In the setup we used that  $s \in \mathbb{C} \cong X_{M_r}^{\mathrm{Mp}_{2n}}$  the first step is to make sure that this is actually true

**Lemma.**  $X_{M_r}^{\mathrm{Mp}_{2n}(\mathbb{A})}$  is at most a one dimensional  $\mathbb{C}$  vector space.

**Proof.** First of all we have that [?, I.1.4]

$$X_{M_r}^{\mathrm{Mp}_{2n}(\mathbb{A})} \subseteq X_{M_r} \cong \mathfrak{a}_{M_r}^* := \mathrm{Rat}(M_r) \otimes_{\mathbb{Z}} \mathbb{C}$$

thus it is clearly sufficient to bound the dimension of  $\mathfrak{a}_{M_r}^*$  as a  $\mathbb{C}$  vector space, moreover this dimension agrees with the dimension of  $\mathrm{Rat}(M_r)$  as a free  $\mathbb{Z}$  module.

Thus we compute  $\dim_{\mathbb{Z}}(\text{Rat}(M_r))$ :

$$\begin{aligned}
 \text{Rat}(M_r) &= \text{Rat}(\text{GL}_r \times \text{Sp}_{2m}) \\
 &= \text{Hom}(\text{GL}_r \times \text{Sp}_{2m}, \mathbb{G}_m) \\
 (2) &\cong \text{Hom}(\text{Ab}(\text{GL}_r \times \text{Sp}_{2m}), \mathbb{G}_m) \\
 (1) &\cong \text{Hom}(\text{Ab}(\text{GL}_r) \times \text{Ab}(\text{Sp}_{2m}), \mathbb{G}_m) \\
 (3) &\cong \text{Hom}(\mathbb{G}_m \times 1, \mathbb{G}_m) \\
 &\cong \mathbb{Z}
 \end{aligned}$$

in (2) we have used the universal property of the abelianization  $\text{Ab}(G) = \mathcal{D}(G) \setminus G = [G, G] \setminus G$  because  $\mathbb{G}_m$  is abelian. (1) is that the abelianization commutes with direct products (citation as comment in Tex). (3) is because  $\text{Sp}$  is a perfect group.

wn that it is not

## 4.4 Lemma 1

The representation is supposed to be of the covering of the Levi//? need to fix this

We here consider the case that  $b = 1$ , hence  $n = a + m$ . Then fixing a standard parabolic of  $\text{Sp}_{2n}$  we have the maximal standard parabolic  $P_a = M_a N_a$  where  $M_a = \text{GL}_a \times \text{Sp}_{2m}$ . Now if  $\tau$  is irreducible unitary cuspidal automorphic representation of  $\text{GL}_a$  then by definition

$$\Delta(\tau, 1)(\varphi)(g) = E(\varphi, s)(g) = s \cdot \varphi(g)$$

where the Eisenstein series is defined via the parabolic induction from the Levi  $(\text{GL}_a)^{\times b}$  to  $\text{GL}_{ab}$ . Thus we have  $\Delta(\tau, 1) = \tau$ . So for the appropriate  $\sigma$  a rep of  $\text{Sp}_{2m}$  we get a rep of the Levi of  $\text{Sp}_{2n}$ ,  $M_r = M_a = \text{GL}_a \times \text{Sp}_{2m}$  given by  $\tau \otimes \sigma$ . To this we associate the Eisenstein series for  $\varphi \in \mathcal{A}(N_r(\mathbb{A})M_r(F) \setminus \text{Mp}_{2n}(\mathbb{A}))_{\tau \otimes \sigma}$   $E(\varphi, s)$  as usual. Now we proceed to calculate the constant term of this Eisenstein series along the parabolic  $P_a = MN$ .

all others are

By our calculations, the cuspidality of the tensor and [?] we know that

$$E(\varphi, s)_P = \sum_w \sum_{m'} \int_{(w^{-1}N(\mathbb{A})w \cap M(\mathbb{A})) \setminus A} \int_{w^{-1}N(F)w \cap M(F) \setminus w^{-1}N(\mathbb{A})w \cap M(\mathbb{A})} \varphi(n_1 n_2 w^{-1} m' x) dn_1 dn_2$$

and the inner integral vanishes for all  $w \neq id, \omega$  ( $\omega$  as in [?]). Hence the first sum becomes over two elements and we have

$$E(\varphi, s)_P = E(\varphi, s)_{P, id} + E(\varphi, s)_{P, \omega}$$

where

$$E(\varphi, s)_{P, w} = \sum_{m' \in M(F) \cap wP(F)w^{-1} \setminus M(F)} \int_{N(F) \cap wP(F)w^{-1} \setminus N(\mathbb{A})} \varphi(w^{-1} n m' x) dn$$

First the identity term simplifies

$$\begin{aligned}
E(\varphi, s)(x)_{P', id} &= \sum_{m' \in M(F) \cap P(F) \backslash M(F)} \int_{N(F) \cap P(F) \backslash N(\mathbb{A})} \varphi(nm'x) dn \\
&= \sum_{m' \in M(F) \backslash M(F)} \int_{N(F) \backslash N(\mathbb{A})} \varphi(nm'x) dn \\
&= \int_{N(F) \backslash N(\mathbb{A})} \varphi(nx) dn \\
&= \varphi(x)_P
\end{aligned}$$

Considering now the  $\omega$  term

$$E(\varphi, s)_{P, \omega} = \sum_{m' \in M(F) \cap \omega P(F) \omega^{-1} \backslash M(F)} \int_{N(F) \cap \omega P(F) \omega^{-1} \backslash N(\mathbb{A})} \varphi(\omega^{-1} nm'x) dn$$

by [?, 2C]  $M(F) \cap \omega P(F) \omega^{-1} \backslash M(F)$  is isomorphic to  $P_0 \backslash \mathrm{Sp}_{2(n-a)}$  where  $P_0 \backslash \mathrm{Sp}_{2(n-a)}$  by definition, but  $P_0$  has Levi  $M_0 = \mathrm{Sp}_{2(n-a)}$  by definition and hence is itself  $\mathrm{Sp}_{2(n-a)}$ . Thus the sum is over  $\mathrm{Sp}_{2(n-a)}(F) \backslash \mathrm{Sp}_{2(n-a)}(F)$  and hence is over a point. Therefore we get by definition of the intertwining operator

$$E(\varphi, s)_{P, \omega} = \int_{N(F) \cap \omega P(F) \omega^{-1} \backslash N(\mathbb{A})} \varphi(\omega^{-1} nx) dn = M(\omega, -)(\varphi)(x)$$

because we took the constant term along the same parabolic as the definition of the Eisenstein series we know that the Levis are (the same) conjugate. Thus we have shown that

$$E(\varphi, s)_P(x) = \varphi(x)_P + M(\omega, -)(\varphi)(x)$$

Notice that the computation takes place completely at the level of the terminals which are independent of the fact that we have taken a covering group, hence we have really only reused work from [?].

I really need to fix the term computations that I dropped in the term computations.

this is not clear in the definition, just guessing this is what I mean

# Appendix A

## Motivation

We have tried to motivate the individual objects as we went, how they would help get us to the result that we were looking for. Here we try to provide a birds eye view of whats going on.

We do have to preface that much like the motivation throughout we have only understood a small piece of the story here, however we still feel obligated to communicate what we can.

### A.0.1 Reminders — Remove Later

Paul Garrett said to me something along the lines of it being arguably more important to understand the *role* of the Eisenstein series in the theory of automorphic forms rather than the technical details of its proof of meromorphicity. I think to some extent he is correct, not to say that the proof isnt important, only that its importance is given to it by its role and is hence subservient to it.

Try to elaborate the role of each introduced item in the theory, not just its technical properties.

# Todo list

<input type="checkbox"/> is this right . . . . .	4
<input type="checkbox"/> Do I need to give a reference for definitions? . . . . .	5
<input type="checkbox"/> check that this is an action, $g$ is a bijection but why does the inclusion still hold? . . . .	5
<input type="checkbox"/> this is probably known by now. . . . .	7
<input type="checkbox"/> This might all be wrong, garretts book is giving something different. . . . .	9
<input type="checkbox"/> example maybe idk . . . . .	11
<input type="checkbox"/> ref . . . . .	11
<input type="checkbox"/> what is easy to find is definitions, whaat is hard to find is long drawnout matrix algebra that has been latexed. . . . .	11
<input type="checkbox"/> maybe put the whole thing in the appendix. . . . .	11
<input type="checkbox"/> fix a global convention . . . . .	11
<input type="checkbox"/> But is the converse true? . . . . .	11
<input type="checkbox"/> perhaps a first natural question is whether or not these things can be given the structure of LAG's. That paper mentions representability. Look into it. . . . .	12
<input type="checkbox"/> reference the definition in the next chapter . . . . .	13
<input type="checkbox"/> Chengjing example of isotypic subspaces . . . . .	13
<input type="checkbox"/> make this more precise . . . . .	16
<input type="checkbox"/> is it true . . . . .	17
<input type="checkbox"/> I have lost the $\lambda$ in the below computation . . . . .	17
<input type="checkbox"/> put this in the Eisenstein series section . . . . .	19
<input type="checkbox"/> I have mixed up my $N$ 's and $U$ 's too much... . . . .	19
<input type="checkbox"/> give references for the other interpretations of the constant term, there must be some .	19
<input type="checkbox"/> reference the maximal parabolic section . . . . .	22
<input type="checkbox"/> prove it . . . . .	22
<input type="checkbox"/> prove it . . . . .	24
<input type="checkbox"/> my $T_m$ differs from Chengjing . . . . .	24
<input type="checkbox"/> Converges uniformly a priori on compact sets, well I dont know if I can swap all these sums haha . . . . .	24
<input type="checkbox"/> ref the lemma . . . . .	25
<input type="checkbox"/> what are the ranges of these . . . . .	26
<input type="checkbox"/> write this up in the classical group section . . . . .	26
<input type="checkbox"/> explain this in the Eisenstein series section . . . . .	26
<input type="checkbox"/> check the exact ranges, will depend on the range of $r$ . . . . .	26
<input type="checkbox"/> link to the residual representation section . . . . .	26



□ explain the $s$ action in Eisenstein series section. . . . .	27
□ reference the section I discuss this in. . . . .	27
□ fill in here as theorems or whatever anything that I end up actually checking.... . . . .	27
□ I havent shown that it is not trivial... . . . .	28
□ The representation is supposed to be of the covering of the Levi///? need to fix this . .	28
□ Brenner.. . . .	28
□ M+W II.1.7, all others are zero..? . . . .	28
□ red . . . . .	28
□ ref . . . . .	28
□ I really need to fix this $s$ thing that I dropped in the constant term computations. . . .	29
□ this is not clear in their paper im just guessing this is what they mean . . . . .	29

# Bibliography

- [Art79] James Arthur. Eisenstein series and the trace formula. In A. Borel and W. Casselman, editors, *Proceedings of Symposia in Pure Mathematics*, volume 33.1, pages 253–274. American Mathematical Society, Providence, Rhode Island, 1979.
- [BC79] Armand Borel and W. Casselman. *Automorphic Forms, Representations, and L-functions*. Number Volume 33 in *Proceedings of Symposia in Pure Mathematics*. American Mathematical Society, Providence (Rhode Island), 1979.
- [Boo] Jeremy Booher. VIEWING MODULAR FORMS AS AUTOMORPHIC REPRESENTATIONS.
- [Bor91] Armand Borel. *Linear Algebraic Groups*, volume 126 of *Graduate Texts in Mathematics*. Springer New York, New York, NY, 1991.
- [Bor97] Armand Borel. *Automorphic Forms on  $SL_2(R)$* . Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1997.
- [Bre09] Eliot Brenner. Notes on Analytic Properties of Residual Eisenstein Series, I, August 2009.
- [Bui] Buildings and Classical Groups - Garrett, Paul B. | 9780412063312 | Amazon.com.au | Books. <https://www.amazon.com.au/Buildings-Classical-Groups-Paul-Garrett/dp/041206331X>.
- [Bum97] Daniel Bump. *Automorphic Forms and Representations*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1997.
- [BVDGHZ08] Jan Hendrik Bruinier, Gerard Van Der Geer, Günter Harder, and Don Zagier. *The 1-2-3 of Modular Forms*. Springer Berlin Heidelberg, Berlin, Heidelberg, 2008.
- [Cli] Clifford Algebras and the Classical Groups. <https://www.cambridge.org/core/books/clifford-algebras-and-the-classical-groups/F14B7362F52D08E4F2118B28B7B22D30>.
- [Con] Brian Conrad. Standard parabolic subgroups: Theory and examples.
- [Eme] Matthew Emerton. CLASSICAL MODULAR FORMS AS AUTOMORPHIC FORMS.
- [Ess] Essays in the History of Lie Groups and Algebraic Groups. <https://bookstore.ams.org/view?ProductCode=HMATH/21>.

- [Fol16] Gerald B. Folland. *A Course in Abstract Harmonic Analysis*. Chapman and Hall/CRC, 0 edition, February 2016.
- [Gar] Paul Garrett. Modern analysis of automorphic forms by examples.
- [Gar16] Paul Garrett. Transition: Eisenstein series on adèle groups. 2016.
- [GGW17] Wee Teck Gan, Fan Gao, and Martin H. Weissman. L-groups and the Langlands program for covering groups: A historical introduction, May 2017.
- [GH24] Jayce R. Getz and Heekyoung Hahn. *An Introduction to Automorphic Representations: With a View toward Trace Formulae*, volume 300 of *Graduate Texts in Mathematics*. Springer International Publishing, Cham, 2024.
- [God] Roger Godement. Groupes linéaires algébriques sur un corps parfait.
- [Gre24] Douglas Grenier. AN ANALOGUE OF SIEGEL’S X-OPERATOR FOR AUTOMORPHIC FORMS FOR  $GL_n(\mathbb{Z})$ . 2024.
- [GS21] David Ginzburg and David Soudry. Top Fourier coefficients of residual Eisenstein series on symplectic or metaplectic groups, induced from Speh representations. *Research in Number Theory*, 8(1):10, December 2021.
- [Hal15] Brian C. Hall. *Lie Groups, Lie Algebras, and Representations: An Elementary Introduction*, volume 222 of *Graduate Texts in Mathematics*. Springer International Publishing, Cham, 2015.
- [JLZ13] Dihua Jiang, Baiying Liu, and Lei Zhang. Poles of certain residual Eisenstein series of classical groups. *Pacific Journal of Mathematics*, 264(1):83–123, July 2013.
- [Kud] Stephen S Kudla. NOTES ON THE LOCAL THETA CORRESPONDENCE.
- [Lap22] Erez Lapid. Some perspectives on Eisenstein series, April 2022.
- [Mak] Shotaro Makisumi. Structure Theory of Reductive Groups through Examples.
- [Mila] J S Milne. Basic Theory of Affine Group Schemes.
- [Milb] J S Milne. Lie Algebras, Algebraic Groups, and Lie Groups.
- [Mil17] J. S. Milne. *Algebraic Groups: The Theory of Group Schemes of Finite Type over a Field*. Cambridge University Press, 1 edition, September 2017.
- [MT] Gunter Malle and Donna Testerman. Linear Algebraic Groups and Finite Groups of Lie Type.
- [Mum99] David Mumford. *The Red Book of Varieties and Schemes*, volume 1358 of *Lecture Notes in Mathematics*. Springer, Berlin, Heidelberg, 1999.
- [MW95] C. Moeglin and J. L. Waldspurger. *Spectral Decomposition and Eisenstein Series: A Paraphrase of the Scriptures*. Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1995.

- [Not] Notes for class on algebraic groups.pdf. <https://users.math.msu.edu/users/ruiterj2/Math/Documents/S>
- [Rao93] R. Ranga Rao. On some explicit formulas in the theory of Weil representation. *Pacific Journal of Mathematics*, 157(2):335–371, January 1993.
- [RS] David Renard and Laurent Schwartz. REPRÉSENTATIONS DES GROUPES RÉDUCTIFS  $p$ -ADIQUES.
- [Ser92] Jean-Pierre Serre. *Lie Algebras and Lie Groups*, volume 1500 of *Lecture Notes in Mathematics*. Springer, Berlin, Heidelberg, 1992.
- [Sha10] Freydoon Shahidi. *Eisenstein Series and Automorphic  $L$ -Functions*, volume 58 of *Colloquium Publications*. American Mathematical Society, Providence, Rhode Island, September 2010.
- [Spr98] T. A. Springer. *Linear Algebraic Groups*. Birkhäuser, Boston, MA, 1998.
- [SZS77] Jean-Pierre Serre, Don Zagier, and Sonderforschungsbereich Theoretische Mathematik, editors. *Proceedings International Conference, University of Bonn, Sonderforschungsbereich Theoretische Mathematik, July 2-14, 1976: Ed. by J.-P. Serre and D. B. Zagier*. Number 601 in *Lecture Notes in Mathematics*. Springer, Berlin, 1977.