

STAT230 Assignment 1

Rin Meng

Student ID: 51940633

January 30, 2024

1. Let S be the sample of the given student, then we have

$$S = \{m, m, m, m, m, m, w, w, w, w\}$$

where m is men, and w is women.

- (a) Let $n = \text{sample size}$. Then there are a number of

$$n! = 10! = 3628800$$

different ways to rank the students by Theorem 2.4.

- (b) Let $n = \text{sample size}$, $w = \text{women in } n$ and $m = \text{men in } n$. Then there are a number of

$$\frac{n!}{m!w!} = \frac{10!}{6! \times 4!} = 210$$

different ways of ranking them with regards to men and women by Theorem 2.7.

2. Let S be the sample of the given flags, then we have

$$S = \{w, w, w, w, r, r, r, b, b\}$$

where w is white flag, r is red flag, and b is blue flag. Then there are a number of

$$\frac{n!}{w! \times r! \times b!} = \frac{9!}{4! \times 3! \times 2!} = 1260$$

different ways to arrange the flags if all flag in the same color is identical.

3. Let S be the set of all possible rolls of a two fair dice, and let n be the number of all possible rolls which is $6 \times 6 = 36$.

(a) Let S_6 be set of two fair dice getting a total number of 6, then we have

$$S_6 = \{(1, 5), (5, 1), (4, 2), (2, 4), (3, 3), \}$$

Let $P(S_6)$ be the probability of getting the number 6 by rolling two fair dice, then we have

$$P(S_6) = \frac{\text{Size of } S_6}{n} = \frac{5}{36}$$

(b) Let $S_{3 \vee 10}$ be the set of two fair dice getting a total number of $3 \vee 10$, then we have

$$S_{3 \vee 10} = \{(1, 2), (2, 1), (5, 5), (4, 6), (6, 4)\}$$

Let $P(S_{3 \vee 10})$ be the probability of getting the number $3 \vee 10$ by rolling two fair dice, then we have

$$P(S_{3 \vee 10}) = \frac{\text{Size of } S_{3 \vee 10}}{n} = \frac{5}{36}$$

(c) Let $S_{\text{odd} \vee \text{prime}}$ be the set of two fair dice getting a total number that is $\text{odd} \vee \text{prime}$, then we have

$$\begin{aligned} S_{\text{odd} \vee \text{prime}} = & \\ & \{(1, 1), (1, 2), (2, 1), (2, 3), (3, 2), \\ & (4, 1), (1, 4), (3, 4), (4, 3), (2, 5), \\ & (5, 2), (1, 6), (6, 1), (5, 6), (6, 5) \\ & (3, 6), (6, 3), (4, 5), (5, 4)\} \end{aligned}$$

Let $P(S_{\text{odd} \vee \text{prime}})$ be the probability of getting a number that is $\text{odd} \vee \text{prime}$ by rolling two fair dice, then we have

$$P(S_{\text{odd} \vee \text{prime}}) = \frac{\text{Size of } S_{\text{odd} \vee \text{prime}}}{n} = \frac{19}{36}$$

4. Let us denote the given data by ϕ , the size of ϕ as n which is 20 data points.

(a) Then the mean of ϕ is

$$\bar{\phi} = \frac{\sum \phi}{n} = \frac{74.4}{20} = 3.72$$

(b) Since we know that $\sum \phi^2 = 278.4196$, the standard deviation of ϕ is

$$s_{\phi} = \sqrt{\frac{\sum \phi^2 - \frac{(74.4)^2}{20}}{n - 1}} = \sqrt{\frac{278.4196 - 276.768}{20 - 1}} = 0.295$$

(c) The median value of ϕ is

$$\phi_{sorted} =$$

$$\{3.55, 3.55, 3.56, 3.56, 3.57, \\ 3.57, 3.59, 3.59, 3.59, 3.60, \\ 3.61, 3.63, 3.65, 3.66, 3.71, \\ 3.73, 3.75, 3.99, 4.15, 4.79\}$$

$$\tilde{\phi} = \frac{3.60 + 3.61}{2} = 3.605$$

(d) The *IQR* of ϕ is

$$Q1 = \frac{3.57 + 3.57}{2} = 3.57$$

$$Q3 = \frac{3.71 + 3.73}{2} = 3.72$$

$$IQR = Q3 - Q1 = 3.72 - 3.57 = 0.15$$

5. Since $P(\{1\}) = \frac{1}{2}$, we can also rewrite as $P(\{1\}) = \frac{3}{6}$ and since $P(\{1, 2\}) = \frac{2}{3}$, we can rewrite as $P(\{1, 2\}) = \frac{4}{6}$


Then we can calculate $P(\{2\})$ as follow:

$$P(\{2\}) = P(\{1, 2\}) - P(\{1\}) = \frac{4}{6} - \frac{3}{6}$$


$$P(\{2\}) = \frac{1}{6}$$

Then from this, we can conclude that $P(\{3\})$ must be equal to $\frac{2}{6}$ because $P(\{1, 2, 3\})$ cannot exceed past $\frac{6}{6}$ namely 1, by *Kolmogorov Axiom 2*.


6. Let S be the sample space, where events A, B, C can occur in S , then
- (a) This is the Venn Diagram among A, B , and C , only A occurs.




- (b) This is the Venn Diagram where at least one of the events A, B , or C occurs.



(c) This is the Venn Diagram where one of A or C occurs, but not B .



(d) This is the Venn Diagram where at most, two of the events A , B , or C occurs.



7. *Proof.* “ A and B are independent $\Leftrightarrow A^c$ and B are independent.”

“ \Rightarrow ”: Since A and B are independent, we have:

$$P(A \cap B) = P(A)P(B) \quad (\Delta)$$

by Definition 2.18 (independence). Now we want to show that:

$$P(A^c \cap B) = P(A^c)P(B)$$

We know that:

$$P(B) = P(A^c \cap B) + P(A \cap B) \quad (\theta)$$

because the probability of B , is the same as probability of A and B happening added with probability of $\neg A$ and B . We then also learn that:

$$P(A^c \cap B) = P(B) - P(A \cap B) \quad (\Theta)$$

by rearranging (θ) .

We will now substitute (Δ) into (Θ) where we will have

$$P(A^c \cap B) = P(B) - P(A)P(B)$$

$$P(A^c \cap B) = P(B)(1 - P(A)) \quad (|)$$

and by Theorem 2.19 (Complement Rule), we learn that

$$(1 - P(A)) = P(A^c)$$

then this equation,

$$P(A^c \cap B) = P(A^c)P(B)$$

is identical to (\dagger) . Hence we have proven that

$$A \text{ and } B \text{ are independent} \Rightarrow A^c \text{ and } B \text{ are independent.}$$

“ \Leftarrow ”: Now, since A^c and B are independent, we have:

$$P(A^c \cap B) = P(A^c)P(B) \quad (\phi)$$

by Definition 2.18 (independence). This proof is trivial as it follows the same structure as the last one I did. Now we want to show that:

$$P(A \cap B) = P(A)P(B)$$

We know that:

$$P(B) = P(A^c \cap B) + P(A \cap B) \quad (\gamma)$$

because the probability of B , is the same as probability of A and B happening added with probability of $\neg A$ and B . We then also learn that:

$$P(A \cap B) = P(B) - P(A^c \cap B) \quad (\Gamma)$$

by rearranging (γ) .

We will now substitute (ϕ) into (Γ) where we will have

$$P(A \cap B) = P(B) - P(A^c)P(B)$$

$$P(A \cap B) = P(B)(1 - P(A^c)) \quad (\psi)$$

and by Theorem 2.19 (Complement Rule), we learn that

$$(1 - P(A^c)) = P(A)$$

then this equation,

$$P(A \cap B) = P(A)P(B)$$

is identical to (ψ) . Hence we have proven that

$$A^c \text{ and } B \text{ are independent} \Rightarrow A \text{ and } B \text{ are independent.}$$

Therefore, We have proven that A and B are independent $\Leftrightarrow A^c$ and B are independent. \square

8. Let S be the event that the silver coin is found and let C be the event that a cabinet was selected.

We know that

$$P(C_A) = P(C_B) = \frac{1}{2}$$

because both drawers in cabinet A (C_A) contains the silver coin, we have

$$P(S|C_A) = \frac{1}{2} + \frac{1}{2} = 1$$

and because only one of the silver coin is in cabinet B (C_B), we have

$$P(S|C_B) = \frac{1}{2}$$

Using Theorem 2.20 (Rule of Probability), then we learn that

$$\begin{aligned} P(S) &= P(S|C_A)P(C_A) + P(S|C_B)P(C_B) \\ &= (1 \times \frac{1}{2}) + (\frac{1}{2} \times \frac{1}{2}) \\ &= \frac{3}{4} \end{aligned}$$

and using Theorem 2.21 (Bayes' Theorem), we can conclude that

$$P(C_A|S) = \frac{P(S|C_A)P(C_A)}{P(S)} = \frac{1 \times \frac{1}{2}}{\frac{3}{4}} = \frac{2}{3}$$

End of Assignment 1.