

**Problem §3** Suppose  $T : V \rightarrow W$  is an injective linear map between finite-dimensional vector spaces. Prove that there exists a linear map  $S : W \rightarrow V$  such that  $ST = I_V$ .

*Solution:* Let  $T \in \mathcal{L}(V, W)$  be an injective linear map. Then every  $v \in V$  maps to a unique  $w \in W$ ; that is, if  $T(v_1) = T(v_2)$ , then  $v_1 = v_2$ . Define a linear map

$$S : W \longrightarrow V, \quad S(w) = \text{the unique } v \text{ such that } T(v) = w; \text{ or, } \mathbf{0} \text{ if no such } v \text{ exists.}$$

In other words, if  $w = T(v)$  for some  $v \in V$ , then  $S(w) = v$ ; and if  $w' \neq T(v')$  for any  $v' \in V$ , then  $S(w') = \mathbf{0}$ . Then for any  $v \in V$  we have  $S(T(v)) = v = I_V(v)$ . (However, clearly under this construction  $TS \neq I_W$ , since if  $w \neq T(v)$  for any  $v \in V$ , then  $T(S(w)) = \mathbf{0} \neq I_W(w)$ ).

**Problem §4** Prove that given distinct  $a_0, \dots, a_d$  and any real numbers  $b_0, \dots, b_d$ , there exists a unique  $f \in \mathcal{P}_d(\mathbb{R})$  such that

$$f(a_0) = b_0, \dots, f(a_d) = b_d.$$

*Solution:* Let

$$T : \mathcal{P}_d(\mathbb{R}) \rightarrow \mathbb{R}^{d+1}, \quad T(f) = (f(a_0), \dots, f(a_d)),$$

and recall the definition of function addition and scalar multiplication:

$$(f + g)(x) = f(x) + g(x), \quad (cf)(x) = c(f(x)).$$

Clearly,  $T$  is a linear transformation. For  $f, g \in \mathcal{P}_d(\mathbb{R})$ ,  $c_1, c_2 \in \mathbb{R}$ , we have

$$\begin{aligned} T(c_1f + c_2g) &= ((c_1f + c_2g)(a_0), \dots, (c_1f + c_2g)(a_d)) \\ &= ((c_1f)(a_0) + (c_2g)(a_0), \dots, (c_1f)(a_d) + (c_2g)(a_d)) \\ &= (c_1f(a_0), \dots, c_1f(a_d)) + (c_2g(a_0), \dots, c_2g(a_d)) \\ &= c_1(f(a_0), \dots, f(a_d)) + c_2(g(a_0), \dots, g(a_d)) \\ &= c_1T(f) + c_2T(g). \end{aligned}$$

Now, we consider the kernel of  $T$ . In order for a function  $f \in \mathcal{P}_d(\mathbb{R})$  to have  $T(f) = (0, \dots, 0)$ , since all of  $a_i \in \mathbb{R}$  are unique,  $f$  must have each  $a_i$  as a root (i.e.  $(x - a_i)a(x) = 0$ , where  $a(x) \in \mathcal{P}_{d-1}(\mathbb{R})$ ). However, recall that a polynomial  $a_0 + \dots + a_dx^d \in \mathcal{P}_d(\mathbb{R})$  can have at max  $d$  distinct roots; thus, no polynomial in  $\mathcal{P}_d(\mathbb{R})$  can have more than  $d$  roots. Since we have  $d + 1$  unique  $a_i$ 's, it necessarily follows that no non-zero function  $f(x) \in \mathcal{P}_d(\mathbb{R})$  will satisfy  $f(x) = 0$ . Thus  $\ker(T) = \{0\}$ .

Next, observe that the dimensions of  $\mathcal{P}_d(\mathbb{R})$  and  $\mathbb{R}^{d+1}$  have the same dimension, and recall that a trivial kernel implies an injective linear transformation (and so  $T$  is injective). But we know that if two vector spaces have equal dimensions, and a map between the two is injective, then the map is also bijective.

Hence  $T$  is an isomorphism; and so for any  $(b_0, \dots, b_d) \in \mathbb{R}^{d+1}$ , we can find a  $f \in \mathcal{P}_d(\mathbb{R})$  such that  $T(f) = (f(a_0), \dots, f(a_d)) = (b_0, \dots, b_d)$ . Thus  $f(a_0) = b_0, \dots, f(a_d) = b_d$ , as required.