

**Problem §1** (6.12) Let  $G$  be a group that acts on a set  $X$ .

- (a) Suppose  $|G| = 15$  and  $|X| = 7$ . Prove there is some element in  $X$  that is fixed by every element of  $G$ .
- (b) What goes wrong if either  $|X| = 6$  or  $|X| = 8$ ?

*Solution:*

- (a) Suppose  $|G| = 15$  and  $|X| = 7$ . We know from Proposition 6.19 that  $|Gx|$  divides  $|G|$ ; furthermore, the distinct orbits  $Gx_1, \dots, Gx_k$  form a disjoint union of  $X$ .

From the Orbit-Stabilizer Counting Theorem, we know that

$$|X| = \sum_{i=1}^k |Gx_i| = \sum_{i=1}^k \frac{|G|}{|G_{x_i}|}.$$

Since  $|Gx_i|$  divides  $|G| = 15$  for all  $1 \leq i \leq k$  and  $\sum_{i=1}^k |Gx_i| = |X| = 7$ , we must have

$$|Gx_i| = 1, 3, \text{ or } 5 \text{ for all distinct orbits.}$$

With these numbers, there are thus only three possible partitions of  $X$  into distinct orbits, up to ordering (each number represents the number of elements in each distinct orbit):

- $7 = 1 + 1 + 5$
- $7 = \underbrace{1 + \dots + 1}_{7 \text{ times}}$
- $7 = 3 + 3 + 1$

In all cases, there is at least one orbit with only one element; this then implies that

$$1 = |Gx_j| = \frac{|G|}{|G_{x_j}|} = \frac{15}{|G_{x_j}|}$$

for some orbit  $Gx_j$  (from Proposition 6.19c). But then for some  $x_j \in X$ , its stabilizer has  $15 = |G|$  elements; in other words, for some element in  $X$ , it is fixed by every element of  $G$ .

- (b) Suppose instead that  $|X| = 6$  or  $8$ . There are then different possible partitions of  $X$ ; but in either case, there exists a partition of  $X$  into distinct orbits that does not consist of any orbit with only 1 element:
  - For  $|X| = 6$ ,  $X$  can be partitioned into two orbits with 3 elements each (since  $|Gx_1| + |Gx_2| = 3 + 3 = 6 = |X|$ , as required by the Orbit-Stabilizer Counting Theorem).
  - For  $|X| = 8$ ,  $X$  can be partitioned into two orbits, one with 3 elements and one with 5 ( $|Gx_1| + |Gx_2| = 5 + 3 = 8 = |X|$ ).

Thus when the group  $G$  acts on the set  $X$ , it is possible that there doesn't exist any orbit with only one element; then by Proposition 6.19, since

$$1 < |Gx_j| = \frac{|G|}{|G_{x_j}|} = \frac{15}{n},$$

where  $1 < n = 3$  or  $5$ . In other words, it is possible that no element in  $X$  is fixed by every element in  $G$  (since the number of elements in the stabilizer of any  $x$  could be less than 15).

**Problem §2** (6.14) Let  $p$  be a prime. We proved that every group with  $p^2$  elements is Abelian; now let  $G$  be a group with  $p^3$  elements.

- (a) Mimic the proof of Corollary 6.26 to try to prove that  $p^3$  is Abelian. Where does the proof go wrong?
- (b) Give two examples of non-Abelian groups with  $2^3$  elements.

(c) What sort of information can you deduce about  $G$  from the proof in (a) that failed?

*Solution:*

- (a) Let  $Z = Z(G)$  as before. Since  $Z$  is a subgroup of  $G$ , Lagrange's Theorem tells us that the order of  $Z$  divides  $|G| = p^3$ , so

$$|Z| = 1, p, p^2, \text{ or } p^3.$$

Theorem 6.25 tells us that  $Z \neq \{e\}$ , so  $|Z| \neq 1$ .

Suppose  $|Z| = p^2$ . Since the center  $Z$  of  $G$  is a normal subgroup, we form the quotient subgroup  $G/Z$ , with Lagrange telling us that

$$|G/Z| = \frac{|G|}{|Z|} = \frac{p^3}{p^2} = p.$$

Thus  $G/Z$  is of prime order, so Proposition 2.43 tells us that it is cyclic. Let  $hZ$  be a coset that generates  $G/Z$ ,

$$G/Z = \{hZ, h^2Z, \dots, h^{p-1}Z\}.$$

In particular, this implies that

$$G = Z \cup hZ \cup \dots \cup h^{p-1}Z,$$

since every element is in a coset of  $Z$ .

Let  $g_1, g_2 \in G$  be arbitrary elements. Since they're in some coset of  $Z$ , we have

$$g_1 = h^{i_1}z_1, \quad g_2 = h^{i_2}z_2 \text{ for some } z_1, z_2 \in Z \text{ and } 0 \leq i_1, i_2 \leq p-1.$$

Since  $z_1, z_2 \in Z$ , we have

$$\begin{aligned} g_1g_2 &= (h^{i_1}z_1)(h^{i_2}z_2) = (h^{i_1}h^{i_2})(z_1z_2) = h^{i_1+i_2}z_2z_1 \\ &= (h^{i_2}h^{i_1})(z_2z_1) = (h^{i_2}z_2)(h^{i_1}z_1) = g_2g_1. \end{aligned}$$

Y I K E S !!! We've shown that every element in  $G$  commutes with every other element; this means that  $Z = G$ , a contradiction of our assumption that  $|Z| = p^2 \neq p^3 = |G|$ . Thus  $|Z| \neq p^2$ .

Now, suppose  $|Z| = p$ .  $Z$  normal allows us to form the quotient subgroup  $G/Z$ , with Lagrange telling us that

$$|G/Z| = \frac{|G|}{|Z|} = \frac{p^3}{p} = p^2.$$

Thus  $G/Z$  has order  $p^2$ , so Corollary 6.26 tells us that it is Abelian. That means for two cosets  $gZ, hZ \in G/Z$ , we have  $ghZ = hgZ$ .

Like before, since every element is in a coset of  $Z$ , and  $G/Z$  is a collection of distinct cosets of  $G$ , we have

$$G = h_1Z \cup h_2Z \cup \dots \cup h_{p^2-1}Z,$$

where  $h_1, \dots, h_{p^2-1} \in G$  form distinct cosets of  $Z$ .

Let  $g_1, g_2 \in G$  be arbitrary elements of  $G$ . Then

$$g_1 = h_i z_1, \quad g_2 = h_j z_2 \text{ for some } h_i, h_j \in G \text{ and } 1 \leq i, j \leq p^2-1.$$

With  $g_1g_2$ , we have

$$g_1g_2 = (h_i z_1)(h_j z_2) = (h_i h_j)(z_1 z_2),$$

and with  $g_2g_1$ , we have

$$g_2g_1 = (h_j z_2)(h_i z_1) = (h_j h_i)(z_2 z_1) = (h_j h_i)(z_1 z_2).$$

Are these two equal?

**Not necessarily.**  $ghZ = hgZ$  means that for every  $ghz \in ghZ$ ,  $ghz \in hgZ$ ; and for every  $hgz' \in hgZ$ ,  $hgz' \in ghZ$ . However, this does **not** guarantee that  $z = z'$ . Indeed, two examples (stated below) illustrate this:  $ghZ = hgZ$  does not guarantee that  $gh = hg$  for all  $g, h \in G$ .

Therefore, since not every element necessarily commutes with every other element, it's possible for  $|Z| = p$ ; there is no contradiction.

Thus there exist non-Abelian groups of order  $p^3$ .

- (b) Two examples of non-Abelian groups of order  $2^3$  are the dihedral group  $\mathcal{D}_4$  (which one can easily verify is not Abelian), and the quaternion group  $\mathcal{Q}$  (see Example 2.18; clearly  $ji = -ij \neq ij$ , and thus is non-commutative). These examples also illustrate that  $G/Z$  being Abelian does not necessarily force  $G$  to be Abelian; both  $\mathcal{D}_4$  and  $\mathcal{Q}$  have centers of order 2 ( $Z(\mathcal{D}_4) = \{e, f\}$  where  $f$  is the flip that fixes the first and third vertices; and  $Z(\mathcal{Q}) = \{\pm 1\}$ ), so  $G/Z$  is Abelian (since  $|G/Z| = 4 = 2^2$ ), yet  $G = \mathcal{D}_4$  or  $\mathcal{Q}$  are not Abelian.
- (c) Let  $G$  be a group with order  $p^3$ . If  $G$  is Abelian, then definitionally  $Z(G) = G$ , so suppose  $Z(G) \neq G$  (i.e.  $G$  is non-Abelian). Then  $|Z(G)| \neq p^3$ ; and from the proof in (a), we see that  $|Z(G)| \neq 1$  and  $|Z(G)| \neq p^2$ ; the only possible value for  $|Z(G)|$  is  $p$ . Therefore, if  $G$  is a non-Abelian group of order  $p^3$ , then its center  $Z(G)$  has order  $p$ .

**Problem §3 (6.17)** This exercise sketches an alternative proof of a key step in the proof of Corollary 6.26.

- (a) Let  $G$  be a group, and let  $g \in G$  be an element that is not in the center of  $G$ . Prove that there is a strict inclusion

$$Z(G) \subsetneq Z_G(g);$$

i.e. prove that the centralizer of  $g$  is strictly larger than the center of  $G$ .

- (b) Let  $G$  be a finite group of prime power order, say  $|G| = p^n$ . Prove that if the center of  $G$  satisfies  $|Z(G)| \geq p^{n-1}$ , then  $Z(G) = G$ , and so  $G$  is Abelian.

*Solution:*

- (a) Let  $g \in G$  be an element not in the center of  $G$ . By definition,

$$Z(G) \subseteq Z_G(g),$$

since  $Z_G(g)$  consists of all elements that commute with any element  $g^i \in \langle g \rangle \subseteq G$ ; but every element  $z \in Z(G)$  commutes with any element in  $G$ . Moreover, since  $g \notin Z(G)$ , we know  $g \neq e$  ( $g$  is non-trivial); hence  $\langle g \rangle \neq \{e\}$  (and so  $\langle g \rangle$  has at least one non-identity element).

Clearly, any  $g^i \in \langle g \rangle$  commutes with any element in  $\langle g \rangle$ . Let  $g^i, g^j \in \langle g \rangle$ . Then

$$g^i g^j = g^{i+j} = g^{j+i} = g^j g^i.$$

Therefore  $\langle g \rangle \subseteq Z_G(g)$ .

But  $g \notin Z(G)$  and  $g \in Z_G(g)$ ; thus  $Z(G) \subsetneq Z_G(g)$ , as required.

- (b) We start with a lemma (since we didn't do 6.16):

**Lemma 1** ( $Z_G(g)$  is Subgroup). *Let  $G$  be a group, and  $g \in G$  an element. Then  $Z_G(g)$  is a subgroup of  $G$ .*

*Proof.* Clearly,  $eg = ge$  for every  $g \in G$ ; thus  $e \in Z_G(g)$ .

Let  $z_g \in Z_G(g)$ . Then  $z_g g = g z_g$ . Since  $z_g \in Z_G(g) \subseteq G$ ,  $z_g^{-1} \in G$  as well. Then

$$z_g g = g z_g \iff z_g^{-1} z_g g = z_g^{-1} g z_g \iff g z_g^{-1} = z_g^{-1} g z_g z_g^{-1} \iff g z_g^{-1} = z_g^{-1} g.$$

Hence  $z_g^{-1} \in Z_G(g)$  for any  $z_g \in Z_G(g)$ .

Finally, suppose  $z_g, z'_g \in Z_G(g)$ , and consider  $z_g z'_g$ . Then

$$z_g z'_g g = z_g g z'_g = g z_g z'_g,$$

and so  $z_g z'_g \in Z_G(g)$  as well. Therefore  $Z_G(g)$  is a subgroup of  $G$ .  $\square$

Suppose  $G$  has order  $p^n$  for some prime number  $p$ , and  $|Z(G)| \geq p^{n-1}$ . Since  $Z(G)$  is a (normal) subgroup of  $G$ , Lagrange's Theorem tells us that  $|Z(G)|$  must be either  $p^{n-1}$  or  $p^n$ .

Suppose  $Z(G) \neq G$ . Then  $|Z(G)| = p^{n-1}$ , and there exists some  $g \in G$  such that  $g \notin Z(G)$  (that is, there exists some  $g \in G \setminus Z(G)$ ). From (a), we know that

$$Z(G) \subsetneq Z_G(g);$$

that is,  $Z_G(g)$  is strictly larger than  $Z(G)$ . But we know that  $Z_G(g)$  is a subgroup of  $G$  from Lemma 1; thus Lagrange tells us that  $|Z_G(g)| = p^r$  for some  $0 \leq r \leq n$ . Since  $|Z(G)| = p^{n-1}$ , we need  $p^{n-1} = |Z(G)| < |Z_G(g)| = p^n$ . Hence  $Z_G(g) = G$ ; but since the choice of  $g \in G \setminus Z(G)$  was arbitrary, that means every element not in  $Z(G)$  commutes with every element in  $G$ . In other words,  $g \in Z(G)$ ; a contradiction.

Therefore  $|Z(G)|$  must be  $p^n$ , and so  $Z(G) = G$ , and so  $G$  is Abelian.

In Corollary 6.26, the above result can be used instead of deriving a contradiction for  $|Z(G)| = p$ . From Lagrange, we know that  $|Z(G)| = 1, p$ , or  $p^2$ . Theorem 6.25 tells us that  $Z(G) \neq \{e\}$ , so  $|Z(G)| \neq 1$ ; that is,  $|Z(G)| = p$  or  $p^2$ . In either case,  $|Z(G)| \geq p^{2-1} = p$ , so from the result above, we know that  $G$  is Abelian.