

Problem §3 (4.15) Suppose V is an F -vector space, \mathcal{A} and \mathcal{B} are subsets of V , and

- \mathcal{B} is linearly independent.
- $|\mathcal{A}| = |\mathcal{B}|$.
- $\text{span}(\mathcal{B}) \subseteq \text{span}(\mathcal{A})$.

Prove $\text{span}(\mathcal{A}) = \text{span}(\mathcal{B})$.

Solution: We start with 2 lemmas.

Lemma 1. *Let V be an F -vector space. Given a subset $\mathcal{A} \subseteq V$, $\text{span}(\mathcal{A})$ is a subspace of V ; and if \mathcal{A} is linearly independent, then \mathcal{A} forms a basis of $\text{span}(\mathcal{A})$.*

Proof. Clearly, given $v_1, v_2 \in \text{span}(\mathcal{A})$, we have $v_1 + v_2 \in \text{span}(\mathcal{A})$ (since for each component $a_i \alpha_i$ of v_1 and corresponding $a'_i \alpha_i$ of v_2 , $(a_i + a'_i) \in F$ by closure of addition in fields, so $v_1 + v_2$ is still in $\text{span}(\mathcal{A})$ by definition); and for any $c \in F$, $cv \in \text{span}(\mathcal{A})$ as well (like above, for each component $a_i \alpha_i$ of v , $(ca_i) \in F$, so $cv \in \text{span}(\mathcal{A})$ by definition). Finally, setting each a_i coefficient to 0 shows that $\mathbf{0} \in \text{span}(\mathcal{A})$. Hence $\text{span}(\mathcal{A})$ is a subspace of V .

If \mathcal{A} is linearly independent, clearly \mathcal{A} forms a basis for $\text{span}(\mathcal{A})$ (since \mathcal{A} is both linearly independent and spans $\text{span}(\mathcal{A})$. Wow!) \square

Lemma 2. *Let a set \mathcal{A} span a vector space V . If a linearly independent set \mathcal{B} has the same number of elements as \mathcal{A} , \mathcal{A} is also linearly independent, and is thus a basis for V .*

Proof. Suppose \mathcal{A} is linearly dependent. Then for some $\alpha_i \in \mathcal{A}$, $\alpha_i \in \text{span}(\mathcal{A} \setminus \{\alpha_i\})$, so $\mathcal{A} \setminus \{\alpha_i\}$ spans V as well. But then $|\mathcal{A} \setminus \{\alpha_i\}| < |\mathcal{A}| = |\mathcal{B}|$ a linearly independent set in V , a contradiction of Lemma 4.24. Thus \mathcal{A} is linearly independent in V as well, and thus is a basis for V . \square

Lemma 3. *Let \mathcal{A} be a basis for a vector space V . If a linearly independent set \mathcal{B} has the same number of elements as \mathcal{A} , then \mathcal{B} is a basis for V as well.*

Proof. Suppose \mathcal{B} is not a basis for V . Then for some $v \in V$ where $v \notin \text{span}(\mathcal{B})$, $\mathcal{B}' = \mathcal{B} \cup \{v\}$ is linearly independent. However, Lemma 4.24 tells us that the size of any linearly independent set in V is less than or equal to the length of any spanning set in V , and since \mathcal{A} spans V and $|\mathcal{A}| = |\mathcal{B}| < |\mathcal{B}'|$, this is a contradiction. Hence \mathcal{B} must span V as well, and so \mathcal{B} is a basis for V . \square

Lemma 1 tells us that $\text{span}(\mathcal{A})$ is a subspace of V (and thus also a vector space, allowing us to apply theorems about bases and dimensions of vector spaces). Since $\text{span}(\mathcal{B}) \subseteq \text{span}(\mathcal{A})$, clearly $\mathcal{B} \subseteq \text{span}(\mathcal{A})$. Thus \mathcal{B} is a linearly independent set in $\text{span}(\mathcal{A})$. Since \mathcal{A} and \mathcal{B} have the same number of elements, Lemma 2 tells us that \mathcal{A} is linearly independent in $\text{span}(\mathcal{A})$ as well, and thus is a basis. Lemma 3 then tells us that \mathcal{B} is a basis for $\text{span}(\mathcal{A})$ too; thus $\text{span}(\mathcal{A}) = \text{span}(\mathcal{B})$, as required.

Problem §4 Suppose V, W are finite-dimensional F -vector spaces. Let $L : V \rightarrow W$ be a linear transformation.

- If L injective, prove $\dim V \leq \dim W$.
- If L surjective, prove $\dim V \geq \dim W$.

Solution: Let $\{v_1, \dots, v_n\}, \{w_1, \dots, w_m\}$ be basis for V and W respectively.

- Suppose $L : V \rightarrow W$ is injective. For any $v \in V$, we can write

$$v = \sum_{i=1}^n a_i v_i, \text{ where } a_i \in F.$$

Clearly, $\mathcal{B} = \{L(v_1), \dots, L(v_n)\}$ spans range L , since for any $L(v) \in \text{range } L$, we have

$$L(v) = L(a_1v_1 + \dots + a_nv_n) = a_1L(v_1) + \dots + a_nL(v_n).$$

We claim that \mathcal{B} is a basis for range L .

Suppose $L(v), L(v') \in \text{range } L$ are different ways of representing a vector in range L ; in other words, $L(v) = L(v')$ and

$$L(v) = \sum_{i=1}^n a_i L(v_i), \quad L(v') = \sum_{i=1}^n a'_i L(v_i), \quad a_i \neq a'_i.$$

By linearity,

$$L(v) = \sum_{i=1}^n a_i L(v_i) = \sum_{i=1}^n L(a_i v_i)$$

and

$$L(v') = \sum_{i=1}^n a'_i L(v_i) = \sum_{i=1}^n L(a'_i v_i).$$

L injective then means

$$\sum_{i=1}^n a_i v_i = \sum_{i=1}^n a'_i v_i,$$

and since $\{v_1, \dots, v_n\}$ is a basis for V (and thus is linearly independent), we necessarily have

$$\sum_{i=1}^n (a_i - a'_i) v_i = 0, \quad a_i - a'_i = 0.$$

and hence $a_i = a'_i$. Equivalently, $L(v)$ and $L(v')$ are the same, and thus every $L(v) \in \text{range } L$ can be represented uniquely as

$$L(v) = \sum_{i=1}^n a_i L(v_i).$$

In other words, \mathcal{B} is a basis for range L .

Since $\text{range } L \subseteq W$ and \mathcal{B} is linearly independent in range L (and thus in W as well), by Lemma 4.24 any spanning set must have at least as many elements as \mathcal{B} . Hence any basis of W must have at least as many elements as \mathcal{B} ; and since $|\mathcal{B}| = \dim V$, we get $\dim W \leq \dim V$, as required.

- Suppose $L : V \rightarrow W$ is surjective. From before, we know that $\mathcal{B} = \{L(v_1), \dots, L(v_n)\}$ spans range L ; since L is surjective (and so $\text{range } L = W$), \mathcal{B} spans W as well. By Lemma 4.24, the size of any linearly independent set in W is less than or equal to the size of any spanning set of W . Since $\{w_1, \dots, w_m\}$ is linearly independent, we thus have $\dim W \leq \text{range } L = \dim V$. Therefore $\dim V \geq \dim W$, as required.

Problem §5 (4.18) Let V be a finite-dimensional F -vector space, and let $U \subseteq V$ be a vector subspace.

- Prove that U is finite-dimensional.
- Prove that $\dim_F U \leq \dim_F V$.
- Prove that

$$U = V \iff \dim_F U = \dim_F V.$$

Solution:

- If $U = \{0\}$, U is clearly finite-dimensional, so suppose $U \neq \{0\}$. Let $u_1 \in U$ be a non-zero vector. If $U = \text{span}(\{u_1\})$, then we are done; otherwise, continue adding non-zero vectors $u_j \in U$ such that

$$u_j \notin \text{span}(\{u_1, \dots, u_{j-1}\}),$$

until $\{u_1, \dots, u_j\}$ forms a spanning set of U . With each addition, $\{u_1, \dots, u_j\}$ is a linearly independent set by construction (since each added vector was not in the span of the previous vectors). Moreover, every linearly independent set $\{u_1, \dots, u_j\}$ is in V , since each $u_i \in U \subseteq V$.

Let $n = \dim V$. Since any basis of V is spanning, and by Lemma 4.24, the number of elements in any linearly independent set in V must be less than or equal to the length of any spanning set in V , the length of $\{u_1, \dots, u_j\}$ must be less than or equal to n . Thus the above process will eventually terminate (it cannot repeat infinitely — or past $j = n$ — since the number of elements must be less than or equal to n), and so we are left with a finite linearly independent spanning set $\{u_1, \dots, u_j\}$ of U . Thus U is finite-dimensional.

- (b) From above, we see that a basis $\{u_1, \dots, u_j\}$ of U cannot have more elements than $n = \dim V$. Hence $\dim_F U \leq \dim_F V$.

Alternatively, let $\{u_1, \dots, u_m\}$ be a basis for U . Then $\{u_1, \dots, u_m\} \subseteq U \subseteq V$ is a linearly independent set of vectors in V . By Lemma 4.24, any linearly independent set of vectors in V cannot have more elements than any spanning set of V . Since a basis $\{v_1, \dots, v_n\}$ of V is the smallest spanning set of V and has $n = \dim V$ elements, any linearly independent set cannot have more than n elements. Thus $|\{u_1, \dots, u_m\}| = \dim_F U \leq \dim_F V = n = |\{v_1, \dots, v_n\}|$.

- (c) Suppose $U = V$. Then a basis $\{v_1, \dots, v_n\}$ of U is also a basis of V , and so $\dim_F U = \dim_F V$.

Conversely, suppose $\dim_F U = \dim_F V = n$, and let $\{u_1, \dots, u_n\}$, $\{v_1, \dots, v_n\}$ be bases for U and V respectively. Since U is a subspace of V , we know that $\{u_1, \dots, u_n\}$ is a linearly independent set of vectors in V . From Lemma 3 (of Problem §3), since $\{u_1, \dots, u_n\}$ is a linearly independent set of vectors with the same number of elements as a basis $\{v_1, \dots, v_n\}$ of V , $\{u_1, \dots, u_n\}$ is a basis of V as well. Thus

$$U = \text{span}(\{u_1, \dots, u_n\}) = V,$$

and so $U = V$, as required.