

1. Say whether the following is true or false and support your answer by a

proof. $(\exists m \in \mathcal{N})(\exists n \in \mathcal{N})(3m+5n=12)$

The statement is false

Proof: list all the possible pairs of m and n

For pairs contains $m \geq 4$ or $n \geq 3$. there is no possible way to produce the equation $3m + 5n = 12$.

The reason is that:

If $m \geq 4$, then $3m + 5n > 12$, because n is a natural number.

Similarly, If $n \geq 3$, then $3m + 5n > 12$. because m is a natural number.

m must be one of 1, 2, 3 and n must be one of 1, 2

6 pairs of m and n can be formed, they are

$m=1$ and $n=1$, $3m+5n=8$

$m=1$ and $n=2$, $3m+5n=13$

$m=2$ and $n=1$, $3m+5n=11$

$m=2$ and $n=2$, $3m+5n=16$

$m=3$ and $n=1$, $3m+5n=14$

$m=3$ and $n=2$, $3m+5n=19$

Thus, no possible pairs of m and n in natural numbers can lead to the equation

$3m+5n=12$.

So the equation is false.

2. Say whether the following is true or false and support your answer by a proof: The sum of any five consecutive integers is divisible by 5 (without remainder).

The statement is true

Proof: By given arbitrary five consecutive integers

List these five numbers for smallest one to biggest one. Suppose the third a of the list is a.

Then the list can be represented as following:

$a-2$, $a-1$, a , $a+1$, $a+2$

As they are conserve integers

Their sum is $(a-2) + (a-1) + a + (a+1) + (a+2) = 5a$

since a is a integers. The sum is divisible by 5.

3. Say whether the following is true or false and support your answer by a proof: For any integer n , the number $n^2 + n + 1$ is odd.

The statement is true

Proof: By assuming n is odd or even

Suppose n is odd, n^2 is odd. Because the product of two odd numbers is still odd

$n^2 + n$ is even. Because the sum of two odd numbers is even

$n^2 + n + 1$ is odd. Because even number + 1 is odd

Suppose n is even

n^2 is even. Because the product of two even numbers is still even

$n^2 + n$ is even. Because the sum of two even numbers is even

$n^2 + n + 1$ is odd. Because the sum of one odd number, 1 in this formula.

and one even number is odd.

That is either n is even or odd, $n^2 + n + 1$ is odd. Because integers are only composed by even

numbers and odd numbers. Thus for any integer n , the number $n^2 + n + 1$ is odd

4. Prove that every odd natural number is of one of the forms $4n + 1$ or $4n + 3$, where n is an integer.

When $n=0$, $4n+1=1$, $4n+3=3$

For any arbitrary n , $n+1$: $4(n+1)+1-(4n+1)=2$, $4(n+1)+3-(4n+3)=2$, that means when n plus 1, the forms $4n+1$ and $4n+3$ will increase by 2.

As it starts with 1 (when $n=0$), so $4n+1$ and $4n+3$ can cover all odd numbers

5. Prove that for any integer n , at least one of the integers n , $n+2$, $n+4$ is divisible by 3.

Solution:

We take out n , if n is divided by 3, there are three conditions:

The remainder is 0, so we can prove that directly

The remainder is 1, so we can know that $(n+2)$ can be divided by 3

The remainder is 2, so we can know that $(n+1+3)=(n+4)$ can be divided by 3

So the statement is true

6. A classic unsolved problem in number theory asks if there are infinitely many pairs of "twin primes", pairs of primes separated by 2, such as 3 and 5, 11 and 13, or 71 and 73. Prove that the only prime triple (i.e. three primes, each 2 from the next) is 3, 5, 7.

Solution:

From question 5 we know that in the three numbers $(n, n+2, n+4)$, at least one number can be divided by 3.

If $n > 3$ so $n+2 > 3$, $n+4 > 3$, that means at least one number can be divided by 3 and itself, so this number is not a prime.

So we just can let $n=0, n=1, n=2, n=3$:

If $n=0$, 0 is not prime

If $n=1$, 1 is not prime

If $n=2$, $n+2=4$, 4 is not prime

If $n=3$, $n+2=5$, $n+4=7$ 3, 5, 7 are primes

So we can prove that

7. Prove that for any natural number n , $2 + 2^2 + 2^3 + \dots + 2^n = 2^{n+1} - 2$

Solution:

$$2 + 2^2 + 2^3 + \dots + 2^n = 2^{n+1} - 2 = 2^n * 2 - 2 = 2^n + 2^n - 2$$

$$2 + 2^2 + 2^3 + \dots + 2^{n-1} = 2^n - 2 = 2^{(n-1)} + 2^{(n-1)} - 2$$

$$2 + 2^2 + 2^3 + \dots + 2^{n-2} = 2^{(n-1)} - 2 = 2^{(n-2)} + 2^{(n-2)} - 2$$

.....

$$2 = 2 + 2 - 2 \text{ which means } 2 = 2$$

So the statement is true

8. Prove (from the definition of a limit of a sequence) that if the sequence $\{a_n\}_{n=1}^{\infty}$ tends to limit L as $n \rightarrow \infty$, then for any fixed number $M > 0$, the sequence $\{Ma_n\}_{n=1}^{\infty}$ tends to the limit ML .

Proof: Take an arbitrary real number E

Because the sequence $\{a_n\}_{n=1}^{\infty}$ tends to limit L as $n \rightarrow \infty$. For the number E/M , there always exist an n_0 that when $m > n_0$, $|a_m - L| < E/M$ which, by algebra, can be transformed to

$$|Ma_m - ML| < E$$

Combining above two points, that is for an arbitrary real number E , there always exist an n_0 that when $m > n_0$, $|Ma_m - ML| < E$. This is necessary and sufficient for the sequence $\{Ma_n\}_{n=1}^{\infty}$ tends to the limit ML . As required.