

A Brief Introduction to Topological Quantum Field Theories

SAMMS 2015


Mario L. Gutierrez Abed
Advisor: Prof. T. Kerler

August 7th, 2015



Overview

- Categories.
- Cobordisms.
- TQFT's.



Categories

Definition

A **category** \mathcal{C} consists of



Categories

Definition

A **category** \mathcal{C} consists of

- a class of **objects**, denoted $\text{Ob}(\mathcal{C})$.



Categories

Definition


A **category** \mathcal{C} consists of

- a class of **objects**, denoted $\text{Ob}(\mathcal{C})$.
- given two objects $x, y \in \mathcal{C}$, a set $\text{Hom}(x, y)$ of **morphisms**.




Properties of Morphisms

- composition




Properties of Morphisms

- composition



- identity morphism $1_x: x \rightarrow x$ such that, for any $f: x \rightarrow y$, we have $f1_x = f = 1_y f$.



Examples of categories

- Set.
 - objects are sets and morphisms are functions.



Examples of categories

- Set.
 - objects are sets and morphisms are functions.
- Vect.
 - objects are (finite-dimensional) vector spaces, and morphisms are linear operators.



Examples of categories

- Set.
 - objects are sets and morphisms are functions.
- Vect.
 - objects are (finite-dimensional) vector spaces, and morphisms are linear operators.
- Hilb.
 - objects are (finite-dimensional) Hilbert spaces, and morphisms are linear operators.



Examples of categories

- Set.
 - objects are sets and morphisms are functions.
- Vect.
 - objects are (finite-dimensional) vector spaces, and morphisms are linear operators.
- Hilb.
 - objects are (finite-dimensional) Hilbert spaces, and morphisms are linear operators.
- $n\text{Cob}$.
 - objects are $(n - 1)$ -dimensional oriented compact manifolds, and morphisms are n -dimensional cobordisms.






What are cobordisms?

Think of **space** as an $(n - 1)$ -dimensional manifold Σ , and then think of **spacetime** as an n -dimensional manifold M .




What are cobordisms?

Think of **space** as an $(n - 1)$ -dimensional manifold Σ , and then think of **spacetime** as an n -dimensional manifold M .



What are cobordisms?


Think of **space** as an $(n - 1)$ -dimensional manifold Σ , and then think of **spacetime** as an n -dimensional manifold M .



This notion of spacetime gives an intuitive representation of a **cobordism**.




What are cobordisms?



In general spaces need not be connected.



What are cobordisms?




In general spaces need not be connected.

Sooooo, what is this business of taking disjoint unions of objects in $n\text{Cob}??$



What are cobordisms?



In general spaces need not be connected.

Sooooo, what is this business of taking disjoint unions of objects in $n\text{Cob}$??

Well, it turns out that $n\text{Cob}$ is a **monoidal category**, i.e. a category equipped with a well defined operation \otimes .





Examples of monoidal categories

- Grp: The operation \otimes is the direct product of groups.



Examples of monoidal categories

- Grp: The operation \otimes is the direct product of groups.
- Vect or Hilb: the operation \otimes is the tensor product of vector (Hilbert, resp) spaces.



Examples of monoidal categories

- Grp: The operation \otimes is the direct product of groups.
- Vect or Hilb: the operation \otimes is the tensor product of vector (Hilbert, resp) spaces.
- $n\text{Cob}$: the operation \otimes , both for objects and for morphisms, is the disjoint union of manifolds.



Disjoint union on $n\text{Cob}$


Using the operation $\otimes = \sqcup$ defined on $n\text{Cob}$, we get weird universes like...



Disjoint union on $n\text{Cob}$

Using the operation $\otimes = \sqcup$ defined on $n\text{Cob}$, we get weird universes like...

$$\begin{array}{c} S \sqcup S \\ \downarrow M^* \sqcup 1_S \\ S \sqcup S \sqcup S \\ \downarrow 1_S \sqcup M \\ S \sqcup S \end{array}$$



Time to talk about TQFT's...

Definition (Sort of...)

*Given categories \mathcal{C} and \mathcal{D} , a **functor** $F: \mathcal{C} \rightarrow \mathcal{D}$ is a map that sends objects to objects, morphisms to morphisms, and preserves sources, targets, identities, and composition.*




Time to talk about TQFT's...

Definition (Sort of...)

Given categories \mathcal{C} and \mathcal{D} , a **functor** $F: \mathcal{C} \rightarrow \mathcal{D}$ is a map that sends objects to objects, morphisms to morphisms, and preserves sources, targets, identities, and composition.

Definition

A topological quantum field theory is a monoidal functor $Z: n\text{Cob} \rightarrow \text{Vect}$.



Action of the TQFT

Let's take a closer look at the situation...

$$Z: 2\text{Cob} \longrightarrow \text{Vect}$$



Action of the TQFT


Let's take a closer look at the situation...

$$\begin{array}{ccc} Z: & 2\text{Cob} & \longrightarrow & \text{Vect} \\ & \bigcirc & \mapsto & V \end{array}$$



Action of the TQFT

Let's take a closer look at the situation...

$$\begin{array}{lll} Z: & 2\text{Cob} & \longrightarrow \text{Vect} \\ & \Downarrow & \downarrow \\ & \text{Diagram} & V \\ & \Downarrow & \downarrow \\ & \text{Diagram} & V \otimes V \rightarrow V \end{array}$$




Action of the TQFT

Let's take a closer look at the situation...


$$\begin{array}{llll} Z: & 2\text{Cob} & \longrightarrow & \text{Vect} \\ & \text{[Diagram]} & \mapsto & V \\ & \text{[Diagram]} & \mapsto & V \otimes V \rightarrow V \\ & \text{[Diagram]} & \mapsto & \mathbb{F} \rightarrow V \end{array}$$

The diagrams in the 2Cob column are: a single circle, a pair of pants (three circles at the top, one at the bottom), and a cap (one circle at the top, one at the bottom).



Action of the TQFT


Let's take a closer look at the situation...



Action of the TQFT

Monoidal “products” on $n\text{Cob}$ and Vect :

$$\begin{array}{c}
 S \sqcup S \\
 \downarrow M^* \sqcup 1_S \\
 S \sqcup S \sqcup S \\
 \downarrow 1_S \sqcup M \\
 S \sqcup S
 \end{array}$$




$$\begin{array}{c}
 V \otimes V \\
 \downarrow \\
 V \otimes V \otimes V \\
 \downarrow \\
 V \otimes V
 \end{array}$$




Now things get interesting...

Consider an oriented closed n -manifold M .



Now things get interesting...

Consider an oriented closed n -manifold M .




Then we have

$$M: \emptyset_{n-1} \longrightarrow \emptyset_{n-1}$$



Now things get interesting...

Consider an oriented closed n -manifold M .




Then we have

$$M: \emptyset_{n-1} \longrightarrow \emptyset_{n-1}$$
$$Z(M): Z(\emptyset_{n-1}) \longrightarrow Z(\emptyset_{n-1})$$



Now things get interesting...

Consider an oriented closed n -manifold M .




Then we have

$$\begin{aligned}M: \emptyset_{n-1} &\longrightarrow \emptyset_{n-1} \\Z(M): Z(\emptyset_{n-1}) &\longrightarrow Z(\emptyset_{n-1}) \\Z(M): \mathbb{F} &\longrightarrow \mathbb{F}\end{aligned}$$



Now things get interesting...

Consider an oriented closed n -manifold M .



Then we have

$$\begin{aligned}M &: \emptyset_{n-1} \longrightarrow \emptyset_{n-1} \\Z(M) &: Z(\emptyset_{n-1}) \longrightarrow Z(\emptyset_{n-1}) \\Z(M) &: \mathbb{F} \longrightarrow \mathbb{F}\end{aligned}$$

$Z(M)$ is a topological invariant!




Moreover...

Let $\Sigma = \emptyset$ and consider the closed manifold $M = \Sigma \times S^1$:



Moreover...

Let $\Sigma = \bigcirc$ and consider the closed manifold $M = \Sigma \times S^1$:



Finishing up...

Hence, $\beta \circ \gamma: \mathbb{F} \rightarrow \mathbb{F}$ gives us the invariant:



Finishing up...

Hence, $\beta \circ \gamma: \mathbb{F} \rightarrow \mathbb{F}$ gives us the invariant:

$$\begin{aligned}\sum_{i,j} \gamma^{ij} \beta_{ij} &= \sum_{i,j} \gamma^{ij} \beta_{ji} && \text{(Since } (\beta_{ij}) \text{ is symmetric)} \\ &= \sum_{i=1}^n \left(\sum_{j=1}^n \gamma^{ij} \beta_{ji} \right) \\ &= \sum_{i=2}^n \left(\sum_{j=1}^n \gamma^{ij} \beta_{ji} \right) + (\gamma^{11} \beta_{11} + \gamma^{12} \beta_{21} + \cdots + \gamma^{1n} \beta_{n1}) \\ &= 1 + \sum_{i=2}^n \left(\sum_{j=1}^n \gamma^{ij} \beta_{ji} \right) \\ &= \text{Tr}(I_d(V)) \\ &= n = \dim V.\end{aligned}$$



THANK YOU!

