

# MATH 725 HW#6

MARIO L. GUTIERREZ ABED  
PROF. J. LOUSTAU

**Exercise (Exercise 1).** Let  $X$  and  $Y$  be nonempty sets. Show that  $\mathcal{F}_{X \times Y} \cong \mathcal{F}_X \otimes \mathcal{F}_Y$ , where  $\mathcal{F}_A$  is the free module (or vector space) on  $A$ .

*Proof.* Take the tensor map  $\tau: \mathcal{F}_X \times \mathcal{F}_Y \rightarrow \mathcal{F}_X \otimes \mathcal{F}_Y$ , and define the bilinear map  $\varphi: \mathcal{F}_X \times \mathcal{F}_Y \rightarrow \mathcal{F}_{X \times Y}$  by

$$\varphi \left( \sum_i \alpha_i x_i, \sum_j \beta_j y_j \right) = \sum_{i,j} \alpha_i \beta_j (x_i, y_j)$$

so that, by the universal property of tensor products, there exists a unique  $\tilde{\tau}: \mathcal{F}_X \otimes \mathcal{F}_Y \rightarrow \mathcal{F}_{X \times Y}$  that satisfies  $\tilde{\tau}\tau = \varphi$ . Notice that  $\tilde{\tau}$  is injective because  $\ker(\tilde{\tau}) = \{0\}$ .

Now if

$$\sum_{i,j} \alpha_i \beta_j (x_i, y_j) \in \mathcal{F}_{X \times Y},$$

then we have

$$\tilde{\tau} \left( \sum_{i,j} \alpha_i \beta_j (x_i \otimes y_j) \right) = \sum_{i,j} \alpha_i \beta_j (x_i, y_j),$$

which shows that  $\tilde{\tau}$  is surjective.

Thus we have shown that  $\mathcal{F}_{X \times Y} \cong \mathcal{F}_X \otimes \mathcal{F}_Y$ , as desired. □

**Exercise (Exercise 2).** Let  $S$  and  $T$  be subspaces of  $V_1$  and  $V_2$ , respectively. Then show that

$$(S \otimes V_2) \cap (V_1 \otimes T) \cong S \otimes T.$$

*Proof.* Let us choose a subspace  $T' \subset V_2$  such that  $V_2 = T \oplus T'$ . Then we have

$$\begin{aligned} (S \otimes V_2) \cap (V_1 \otimes T) &= (S \otimes (T + T')) \cap (V_1 \otimes T) \\ &= ((S \otimes T) + (S \otimes T')) \cap (V_1 \otimes T) \\ &= (S \otimes T) + \underbrace{((S \otimes T') \cap (V_1 \otimes T))}_W \quad (\text{By associativity}) \end{aligned}$$

But now notice that

$$\begin{aligned} W &= (S \otimes T') \cap (V_1 \otimes T) \subseteq (V_1 \otimes T') \cap (V_1 \otimes T) && (\text{Since } S \subseteq V_1) \\ &= V_1 \times (T' \cap T) && (\text{By exercise 2 of HW \# 5}) \\ &= \{0\}. \end{aligned}$$

Hence we have

$$(S \otimes V_2) \cap (V_1 \otimes T) = (S \otimes T) + \{0\} = S \otimes T. \quad \square$$

**Exercise (Exercise 3).** If  $T \in \mathcal{L}(V)$  is a linear transformation of a finite-dimensional vector space such that  $T^m = 0$  for some  $m \geq 1$ , then there is a basis of  $V$  of the form

$$u_1, Tu_1, \dots, T^{a_1-1}u_1, \dots, u_k, Tu_k, \dots, T^{a_k-1}u_k$$

where  $T^{a_i}u_i = 0$  for  $1 \leq i \leq k$ .

*Proof.* We work by induction on  $\dim(V)$ . For the inductive step we may assume that  $\dim(V) \geq 1$ . Clearly  $T(V)$  is properly contained in  $V$ , since otherwise  $T^m(V) = \dots = T(V) = V$ , a contradiction. Moreover, if  $T$  is the zero map then the result is trivial. We may therefore assume that  $0 \subset T(V) \subset V$ . By applying the inductive hypothesis to the map induced by  $T$  on  $T(V)$ , we may find  $v_1, \dots, v_l \in T(V)$  so that

$$v_1, Tv_1, \dots, T^{b_1-1}v_1, \dots, v_l, Tv_l, \dots, T^{b_l-1}v_l$$

is a basis for  $T(V)$  and  $T^{b_i}v_i = 0$  for  $1 \leq i \leq l$ .

For  $1 \leq i \leq l$ , choose  $u_i \in V$  such that  $Tu_i = v_i$ . Clearly  $\ker(T)$  contains the linearly independent vectors  $T^{b_1-1}v_1, \dots, T^{b_l-1}v_l$ . Now we extend these to a basis of  $\ker(T)$  by adjoining the vectors  $w_1, \dots, w_m$ . We claim that

$$u_1, Tu_1, \dots, T^{b_1}u_1, \dots, u_l, Tu_l, \dots, T^{b_l}u_l, w_1, \dots, w_m$$

is a basis for  $V$ . Linear independence may easily be checked by applying  $T$  to a given linear relation between the vectors. To show that they span  $V$ , we use dimension counting:

We know that  $\dim(\ker(T)) = l + m$  and that  $\dim(T(V)) = b_1 + \dots + b_l$ . Hence, by the rank-nullity theorem,

$$\dim(V) = (b_1 + 1) + \dots + (b_l + 1) + m,$$

which is the number of vectors in our claimed basis. We have therefore constructed a basis for  $V$  in which  $T$  is in Jordan normal form.  $\square$