



A.M.($\leq$)	Any	Moon P.	Gray
Image Quality		Seeing < 2.0"	
Cloudy Cover		Any	

Title of Proposal: Studying small Solar System bodies through stellar occultations

Telescope 1.60m Perkin-Elmer

Abstract:

An accurate characterization of small bodies in the Solar System is possible from the stellar occultation technique. This is obtained from the determination of accurate shapes/sizes, the detection of tenuous atmospheres, as well as structures (rings, etc) around those objects. Continuous observations of occultation events contributes to a broad description of small bodies and to a better understanding of the history and evolution of the Solar System. Here, we propose to observe events by one Jupiter trojan, six TNOs and the largest known Oort cloud comet. Orcus, a large TNO, and our cometary target, have not yet been observed in an occultation, while Ixion's ellipsoidal shape requires further analysis. TNOs are thought to share similarities with Jupiter/Neptune trojans, as well as cometary nuclei, making their knowledge crucial for Solar System studies. Detailed models of 2002 KX14 are necessary to confirm its density, while little is known about the other TNOs here considered.

Time requested: 13.5 hours.

Minimum time accepted: 9 hours.

Observing Mode: Classical Remote (CR)

Program Type: Regular Proposal (RP)

Principal Investigator (PI): Julio Ignacio Bueno de Camargo

PI Affiliation: ON

PI Phone/PI Email: +55 21 3504-9304 camargo@on.br

Co-authors

Name	Affiliation	E-mail
Júlio Camargo	ON	jibcamargo@yahoo.com
Marcelo Assafin	OV/UFRJ	massaf@ov.ufrj.br
Gustavo Benedetti-rossi	Outros	gugabrossi@gmail.com
Felipe Braga-ribas	UTFPR/DAFIS	felipebribas@gmail.com
Josselin Desmars	Observatoire de Paris	josselin.desmars@obspm.fr
Altair Ramos Gomes-junior	UFU	altairgomesjr@gmail.com
Roberto Vieira-Martins	ON	rvm@on.br
Bruno Eduardo Morgado	OV/UFRJ	bmorgado@ov.ufrj.br
Chrystian Luciano Pereira	ON	chryslp.fis@gmail.com
Bruno Sicardy	Observatoire de Paris	bruno.sicardy@obspm.fr
Giuliano Margoti	ON	giulianomargoti@on.br
Jonatã Arcas Silva	ON	jonatasilva@on.br
Eros Gradovski	ON	erosgradovski@gmail.com
Sérgio Santos Filho	OV/UFRJ	sergiosantos@ov.ufrj.br
Rodrigo Carlos Boufleur	Outros	rcboufleur@gmail.com

Viviane Figueiredo Peixoto	OV/UFRJ	vivianepeixoto989@gmail.com
Rafael Sfair De Oliveira	UNESP	rafael.sfair@unesp.br
Victor Lattari	UNESP	victor.lattari@unesp.br
Tiago Francisco Pinheiro	ON	franciscopinheiro@on.br
Patricia Buzzato	UNESP	patricia.buzzatto@unesp.br
Luana Liberato Mendes	UNESP	luana.l.mendes@unesp.br
Thamiris De Santana	UNESP	t.santana@unesp.br

Is there a thesis or dissertation that will benefit from the data?: YES

Student	E-mail	Degree	Defense
Jonatã Arcas Silva	jonatasilva@on.br	PhD student	03/2029
Eros Gradovski	erosgradovski@gmail.com	PhD student	04/2029
Giuliano Margoti	giulianomargoti@on.br	PhD student	11/2028
Viviane Figueiredo Peixoto	vivianepeixoto989@gmail.com	PhD student	03/2029

Optimal dates:

07/MAR, 09/APR, 30/APR, 05/MAY, 09/MAY, 14/MAY, 28/JUN, 24/JUL, 07/AGO

Impossible dates:

Any other

Instruments:

Instrument	Camera
SPARC4	4 x iXon Ultra 888 CCDs in Electron Multipliering (EM) mode

Instrument	Filter	Grating	Slit
------------	--------	---------	------

Scientific Justification

Small bodies in the Solar System are objects orbiting the Sun (natural satellites excluded) that do not have enough gravity to pull their shapes into a nearly spherical one. This includes (but not all) Main Belt Asteroids, Centaurs, Trans-Neptunian Objects (TNOs), comets, trojan asteroids.

TNOs and Centaurs have been our main targets at OPD, but also some trojans and comets. TNOs are further from the Sun than 30 AU (Neptune's heliocentric distance). Centaurs are objects in unstable orbits between those of Jupiter and Neptune, and they are as rich a class as TNOs alone [1]. They are thought to have, in addition, a common origin with the TNOs [2], so that Centaurs can be considered as brighter proxies of the more distant TNO population.

Their importance resides in the fact that distant small bodies are relatively unchanged relics of an evolved planetesimal disc [3-4]. Therefore, they are records of the original material that formed the solar nebula and a treasure trove to access the history and evolution of the outer Solar System. Given their faint magnitudes, there is still much to be learned from this cold and distant world.

In this study, we propose the use of the stellar occultation technique, where the flux of a star is recorded with a high temporal resolution when it is blocked by a passing small body. In this way, it is possible to characterize the body's physical properties on an individual basis that also helps to build a broader general scenario, giving a crucial view about the processes involved in the formation and evolution of the Solar System. Two striking features of stellar occultations are: possibility to translate high temporal resolution into high spatial resolution and to observe small sized objects beyond our current capability of detection (in a context of serendipitous observations).

On an individual basis, occultations deliver dimensions with kilometric accuracies [5-8]. Shapes are also accurately determined and related to density (therefore, mass), elongation, and rotation for large TNOs [8]. Accurate albedos (a measurement of the ratio between the incident and reflected light fluxes) are derived from accurate sizes and better constrain the nature of surfaces (for example, fresh ice or cometary type material, darkened by space weathering [9]). Sizes, from Kepler's laws, provide information on density for TNOs with satellites (Pluto, Eris, Haumea, etc). Densities tell about the internal structure of these objects, like the ice/rock ratio. Given the high temporal resolution in image acquisition, stellar occultations are unrivalled in providing details about the neighbourhoods of the occulting bodies (e.g., the discovery of rings around the Centaurs Chariklo and Chiron and the TNOs Haumea and Quaoar [6,19-22]). The presence of tenuous atmospheres (down to few nanobars) [5-6][10], as well as other interesting physical properties [11-15], are natural byproducts of an occultation light curve analysis.

In a general scenario, sizes of many TNOs lead to their size distribution (giving insights into the total mass in the transneptunian region) and the balance between accretion and collisional erosion [16]. Physical surface properties of TNOs tell us about the thermal and chemical processes in the outer protoplanetary disk, indicating when and where the objects formed [17] - generally, objects that formed closer to the Sun tend to display more neutral colors; objects that formed further from the Sun are redder due to, e.g., differing compositions and space weathering. Planetary migration (Neptune mainly) then placed these objects into their current places. Albedos, a basic quantity to the study of small bodies, are fundamental to constrain the concentration of ices, carbons, and organics, which can have widely differing reflectivities [18]. As mentioned above, accurate albedos can be determined from accurate diameters - a direct measurement from stellar occultations - and absolute magnitudes.

Here, we propose to observe events by one Jupiter trojan (1996 ST1), the largest known Oort Cloud comet (Bernardinelli-Bernstein), and six TNOs: Orcus, 2004 EW95, 2014 GU53, 2007 JH43, Ixion and 2002 KX14 (see Fig. 3). Orcus, one of the largest TNOs, and comet B-B, have never been detected in an occultation. Ixion, a moderately large TNO, needs a conclusive analysis to confirm its ellipsoidal three-dimensional shape (suggested by previous occultations). A genetic dynamical relationship is believed to exist between TNOs and Trojans (Jupiter/Neptune ones), that have compositional similarities with TNOs and cometary nuclei [23]. Therefore, a comprehensive knowledge about these objects' sizes becomes relevant to improve our understanding about the evolution of our Solar System. A more detailed model of the shape of 2002 KX14 is relevant to confirm its density [24]. 2004 EW95, 2014 GU53 and 2007 JH43 have diameters ranging from 300-550 km and little is known about them from occultations. Observations of all these events with OPD/PE provide invaluable pieces of information on sizes and shapes, along with exciting discoveries of satellites, rings, and other features.

References

1. Peixinho, N., Audrey, T., Tegler, S.C. et al. 2020, "The Trans-Neptunian Solar System", Elsevier, 307
2. Fernández, Y.R., Jewitt, D.C., Sheppard, S.S. 2002, AJ, 123, 1050
3. Levison, H. F. & Morbidelli, A. 2003, Nature, 426, 419
4. Gladman, B., Marsden, B. G., Vanlaerhoven, C. 2008, "The Solar System Beyond Neptune", University of Arizona Press, Tucson, 43
5. Sicardy, B., Ortiz, J. L., Assafin, M., et al. 2011, Nature, 478, 493
6. Ortiz, J. L., Sicardy, B., Braga-Ribas, F., et al. 2012, Nature, 491, 566
7. Gomes-Junior, Giacchini, B.L., Braga-Ribas, F., et al. 2015, MNRAS, 451, 2295
8. Braga-Ribas, F., Sicardy, B., Ortiz, J. L., et al. 2013, ApJ, 773, 26
9. Elliot, J. L., Person, M. J., Zuluaga, C. A., et al. 2010, Nature, 465, 897
10. Widemann, T., Sicardy, B., Dusser, R., et al. 2009, Icarus, 199, 458
11. Ortiz, J.L., Santos-Sanz, P., Sicardy, B., et al. 2017, Nature, 550, 219
12. Sicardy, B., Talbot, J., Meza, E., et al. 2016, ApJ, 819, 38
13. Dias-Oliveira, A., Sicardy, B., Lellouch, E., et al. 2015, ApJ, 811, 53
14. Dias-Oliveira, A., Sicardy, B., Ortiz, J. L., et al. 2017, AJ, 154, 22
15. Rommel, F. L., Braga-Ribas, F.; Ortiz, J. L., et al. 2023, A&A, 678, A167
16. Petit, J.-M., Kavelaars, J.J., Gladman, B., Lored, T. 2008, "The Solar System Beyond Neptune", University of Arizona Press, Tucson, 71
17. Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A. 2008, "The Solar System Beyond Neptune", University of Arizona Press, Tucson, 3
18. Barucci, M. A., Brown, M. E., Emery, J. P., Merlin, F. 2008, "The Solar System Beyond Neptune", University of Arizona Press, Tucson, 143
19. Morgado, B.E., Sicardy, B., Braga-Ribas, F., et al. 2023, Nature, 614, 239
20. Pereira, C.L., Sicardy, B., Morgado, B.E. et al. 2023, A&A, 673L, 4
21. Braga-Ribas, F., Sicardy, B., Ortiz, J. L. et al. 2014, Nature, 508, 72
22. Pereira, C.L., Braga-Ribas, F., Sicardy, B. et al. 2025, ApJL, DOI: 10.3847/2041-8213/ae0b6d
23. Dotto, E., Emery, J.P., Barucci, M. A. et al. 2008, "The Solar System Beyond Neptune", University of Arizona Press, Tucson, 383
24. Rivos, J.L., Ortiz, J.L., Rommel, F.L. et al. 2025, A&A, 697, A62, DOI: 10.1051/0004-6361/202554154

Experimental Design

Sample and target selection: Targets are selected following basic criteria: visibility from OPD (do predictions indicate that the event shadow will cross the telescope site?); magnitude limit (magnitude of the occulted star and contrast between it and the occulting object); scientific interest (has the object ever been observed?; does it need more observations (to pin down size/shape, ring density/dynamics, is it targeted by a space mission, etc)?; is there any special feature - presence of ring, satellite, atmosphere, cometary activity, topography - that needs confirmation or deeper characterization?

Observing strategy: The basic point of the observational strategy is founded on the sampling of the target star flux with the highest possible temporal resolution. In other words, we want to observe the occultation event (before, during and after it) with short exposure times and ideally no dead time between consecutive images. Filters are useful to help, for instance, characterize atmospheres and ring particle size. SPARC4 has been a very useful instrument at OPD for occultations.

Observables: The observable is the flux variation of the event so that the variation of this flux, as a function of time, can be investigated. Stellar occultation translates temporal resolution into spatial resolution. Therefore, the shorter the exposure and dead times the more details about the occulted object and its neighborhoods will be contained in the light curve. Since light curves from different observatories are combined to recover the object's size and shape (and, consequently, its center), clocks should present no offsets. In this way, not only should a clock be precise but accuracy is also of the highest importance. GPS or internet protocols are frequently used to correct clocks from different sites for any offsets.

Data analysis: Achieving kilometric accuracy in determining the sizes of small solar system bodies from stellar occultations requires a meticulous approach to analyze the light curves. This involves fitting a model to the observational data that accounts for the finite exposure time, the apparent angular size of the occulted star at the distance of the occulting body and Fresnel diffraction effects (particularly important for precisely measuring features like rings).

Once the precise instants of the disappearances and reappearances of a star (ingress and egress times) are determined from the light curves of a given event, the projected shape of the object onto the sky plane can be reconstructed and, from some assumptions, the ellipsoidal figure and orientation can be determined. Here, the position of the occulted star, the object's ephemeris, the positions of the observing sites on the Earth as well as a model of the Earth's shape, are necessary.

Timing issues are frequently present in a few of the observations. These include constant time offsets, which can be corrected using a reliable site as a reference, and truncated timestamps (showing only the integer part of the second), which can be fixed - assuming that the acquisition cycle is regular - by performing a linear fit on the image-sequence $\times$ time to determine the acquisition cycle (the slope from the fit gives the cycle time). Combining data from multiple telescopes at one site can also help compensate for data gaps caused by long camera readout times.

Quantifications: SNR determination in the present context takes into consideration the magnitude drop of the event, determined when both star and small solar system body are no longer resolved, and the magnitude of the occulted star. The magnitude drop is given by the combined fluxes of the star+object and that of the object only (in the absence of an atmosphere, the total obstruction of the stellar flux is an accepted simplification). This is given by $\Delta m = -2.5 * \log(1 + 10^{(m_a - m_e) * 0.4})$, where m_a and m_e are the solar system body and star magnitudes respectively. Obviously, although it is not necessary to detect the object's flux during the observation, the flux of the occulted star and that of the sky are of fundamental importance (if the star is too faint, the magnitude drop may not be detected regardless the occurrence of the event and the magnitude of the occulting body). The value of Δm for the events proposed in this work ranges from -1.3 to -6.7. Considering that the magnitude uncertainty can be estimated from $\frac{2.5}{\ln(10)} \times \frac{1}{SNR}$, a SNR of 10, for instance, is enough. The speed of the occultation shadow on the ground is also of relevance, as slow speeds accept longer exposure times without losses to spatial resolution. In practice, exposure times ranging from fractions of a second to few seconds are used. Since occultations are serendipitous and transient events, observations are done regardless seeing, moon or clouds presence (limited to the safety of the telescope). Also, if necessary but at the cost of time (and spatial) resolution, images can be co-added to improve SNR.

Technical Description

Instrument configuration: High temporal resolution with adequate SNR can be obtained from a combination between the telescope aperture and a fast readout EM camera. The latter is easily obtained from SPARC4 iXon EM detectors. For the events proposed here, the PE is the most suitable one at OPD (most of them can be considered faint - 3 stars brighter than $G=15$ can profit from higher temporal resolution with good SNR).

Sky conditions: Since occultations are serendipitous and transient events, observations are done regardless seeing, moon or clouds presence (limited to the safety of the telescope). Objects may potentially hide features that only high spatial resolution can unveil and stellar occultation is the unrivaled and cheapest ground-based technique for this task. With little telescope time and simple instrumentation, moons, rings, close binary systems (where we thought to exist a single object), atmospheres and other features can be discovered at very large distances from the Sun. Therefore, image acquisition, even under poor conditions, is our preferred option.

Exposure times and SNR: The exposure time depends on factors like the magnitude of the star (frequently, the asteroid flux has little influence due to its faint magnitude and short exposure times) and the expected duration of the event. We want the best compromise between time resolution (sequence of images with short exposure times and virtually with no dead time), high SNR to accurately model the occultation light curve, and sampling inside the event. From our experience, considering clear skies and given the magnitudes of the occulted stars (see Sect. "Object List"), exposure times from < 1 s to 2-3 seconds are enough for our purposes. **Please note: We request, to the observation of the occultations, the use of the EMCCD up to 100 for 30 minutes and a readout rate of 10MHz. Exposures as short as 0.1s may also be necessary.**

Total observing time and overheads: In general, a total of 30 minutes per event is enough. In practice, we must add time to properly place the target in the FOV (in case of low density stellar fields or of a too bright star in the field) and set the instrument configuration (exposure time may be increased depending on the sky conditions, etc). In this way, 2 hours/event are enough to comfortably have everything set. Nine events involving eight objects, listed below (event web pages are also provided), are proposed here. Since the minimum observing time unit offered by the OPD is one night, we request nine nights. A relevant backup program, that contributes to this one, is presented later in this section.

1. Bernardinelli-Bernstein: 2026APR09T22:53:00 UTC (<https://shorturl.at/YxQNa>)
2. Orcus: 2026MAY01T00:46:00 UTC (<https://shorturl.at/DIQiP>)
3. 2007 JH43: 2026MAY15T01:32:00 UTC (<https://h1.nu/1hBCK>)
4. Ixion: 2026JUN28T23:18:00 UTC (<https://h1.nu/1cTbv>)
5. Ixion: 2026JUL25T04:31:00 UTC (<https://h1.nu/1hBCV>)
6. 2002 KX14: 2026AUG07T21:55:00 UTC (<https://h1.nu/1cTbL>)
7. (39639) 1996 ST1: 2026MAR08T04:29:00 UTC (<https://shorturl.at/885dO>)
8. 2004 EW95: 2026MAY10T05:32:00 UTC (<https://shorturl.at/WUMWA>)
9. 2014 GU53: 2026MAY06T04:07:00 UTC (<https://h1.nu/1cTbk>)

Minimum observing time: The last three targets above can be considered as lower priorities. We would like to observe as many occultations as possible, taking into account the relevance of the events proposed here and the number of nights granted by the TAC. However, we stress that stellar occultations offer excellent results with relatively small telescope time.

Backup program: Once the occultation is recorded, the remaining of the night will be used to photometric and astrometric observations. Astrometry is of utmost importance to improve the accuracy of predictions. Some of the targets of this proposal are natural targets for astrometry. The need for such astrometric task should significantly decrease as a consequence of the observations from the Vera Rubin Observatory's survey - the LSST. Photometric measurements to measure the rotational period of the occulting objects is also important. With their help, occultations observed in different epochs when the object is probed in different rotational phases, can be combined into a 3-D shape.

Coordination with other observatories: The events by 2004 EW95, 2014 GU53 and 2007 JH43 are in coordination with the VLT. The event by Ixion (25/JUL) is in coordination with SOAR and VLT.

Previous missions of this proposal to this telescope

Here, we report the last five proposals of this one.

- **OP2023B-002:** observations in September, October and December of 5 events by TNOs, a Jupiter trojan and the rings of Neptune. Three nights were lost due to the weather, one event (1998 SN165 was negative) and one event (by the Centaur Chiron) was positive! **This event is reported, along with other observations of this same event from different sites, in The Astrophysical Journal Letters and also in various press releases** (see Fig. 2)!
- **OP2024A-006:** observations in April and July of two TNOs. One night was lost due to the weather (08/APR). The event on 19/JUL by the TNO 2002 KX14 was negative.
- **OP2024B-003:** observations in October, November and December of 2 TNOs, a comet (Bernardinelli-Bernstein) and Miranda (moon of Uranus). Three nights were lost due to bad weather, and the events by Miranda and the Neptune rings are under analysis in collaboration with a french team (Paris Observatory).
- **OP2025A-005:** observations in March, May, June and July by three TNOs, one Centaur and one Jupiter trojan. Two observations were lost due to technical problems. The events by Pholus (Centaur, see, Fig. 1) and 2002 KX14 (TNO) were negative. The event by Dzwinnia (TNO) was also probably negative but, given the presence of clouds, a careful analysis is underway before a conclusion.
- **OP2025B-002:** observations in September, November and December. The only run so far, aiming at an occultation event by the large Oort Cloud comet Bernardinelli-Bernstein, was lost due to the weather.

Previous missions of other proposals to this telescope

- **OP2024B-002:** "*Astrometry and photometry of small bodies in the Solar System*" - Ten nights granted, 65% lost due bad weather. All data reduced. Positions were obtained and sent to <https://lesia.obspm.fr/lucky-star/predictions.php>. The results from proposals like this are important to improve the predictions of stellar occultations.
- **OP2025A-006:** "*Astrometry and photometry of small bodies in the Solar System*" - Eight nights granted, 35% lost due to bad weather. All data reduced. Positions were obtained and sent to <https://lesia.obspm.fr/lucky-star/predictions.php>. The results from proposals like this are important to improve the predictions of stellar occultations.
- **2025A-P009:** "*Mutual events of the Saturnian satellites*": Eight nights granted. We could not observe in one of those nights because the event occurred too early, after the technician's working hours (06:43 Local Time). Of the remaining nights, we failed to obtain data for only one night due to weather. The data is still under analysis.
- **2025B-P008:** "*Mutual events of the Saturnian satellites*" - Four nights were granted, of which two have already been observed, with one of them being lost due to weather. The data is still under analysis.
- **2025B-P005:** "*Multiband rotational photometry of small bodies in the Solar System*" - Four observation nights have been granted, which will be conducted in person between October 23rd and 27th.

Previous results in the field by the Principal Investigator

1. "The Rings of (2060) Chiron: Evidence of an Evolving System", Pereira, C.L. et al. (2025), ApJL, DOI: 10.3847/2041-8213/ae0b6d
2. "Size and Shape of Jupiter Trojan (2207) Antenor from Stellar Occultations", Ferreira, F.S. et al. (2025), AJ, DOI: 10.3847/1538-3881/ade0dc
3. "The trans-Neptunian object (119951) 2002 KX14 revealed via multiple stellar occultations", Rizos, J.L. et al. (2025), A&A, DOI: 10.1051/0004-6361/202554154
4. "Centaur 29P/Schwassmann-Wachmann 1 and its near-nucleus environment from a stellar occultation", Pereira, C.L. et al. (2025), Philosophical Transactions A, DOI: 10.1098/rsta.2024.0189
5. "A study of centaur (54598) Bienor from multiple stellar occultations and rotational light curves", Kretlow, M. et al. (2024), A&A, DOI: 10.1051/0004-6361/202450833
6. "Constraints on the evolution of the Triton atmosphere from occultations: 1989-2022", Sicardy, B. et al. (2024), A&A, DOI: 10.1051/0004-6361/202348756
7. "Physical properties of Centaur (60558) 174P/Echeclus from stellar occultations", Pereira, C.L. et al. (2024), MNRAS, DOI: 10.1093/mnras/stad3318
8. "Kilometer-precise (Ull) Umbriel physical properties from the multichord stellar occultation on 2020 September 21", Assafin, M. et al. (2023), MNRAS, DOI: 10.1093/mnras/stad3093
9. "A large topographic feature on the surface of the trans-Neptunian object (307261) 2002 MS4 measured from stellar occultations", Rommel, F. et al. (2023), A&A, DOI: 10.1051/0004-6361/202346892
10. "Changing material around (2060) Chiron revealed by an occultation on December 15, 2022", Ortiz, J.L. et al. (2023), A&A, DOI: 10.1051/0004-6361/202347025
11. "Constraints on (2060) Chiron's size, shape, and surrounding material from the November 2018 and September 2019 stellar occultations", Braga-Ribas, F. et al. (2023), A&A, DOI: 10.1051/0004-6361/202346749
12. "The two rings of (50000) Quaoar", Pereira, C.L. et al. (2023), A&A, DOI: 10.1051/0004-6361/202346365
13. "A dense ring of the trans-Neptunian object Quaoar outside its Roche limit". Morgado, B.E. et al. (2023), Nature, DOI: 10.1038/s41586-022-05629-6
14. "The multichord stellar occultation by the centaur Bienor on January 11, 2019", Fernández-Valenzuela, E. et al. (2023), A&A, DOI: 10.1051/0004-6361/202243214
15. "Physical properties of the trans-Neptunian object (38628) Huya from a multi-chord stellar occultation", Santos-Sanz, P. et al. (2022), A&A, DOI: 10.1051/0004-6361/202141546
16. "The multichord stellar occultation on 2019 October 22 by the trans-Neptunian object (84922) 2003 VS₂", Vara-Lubiano, M. et al. (2022), A&A, DOI: 10.1051/0004-6361/202141842
17. "Constraints on the structure and seasonal variations of Triton's atmosphere from the 5 October 2017 stellar occultation and previous observations", Marques Oliveira, J. et al. (2022), A&A, DOI: 10.1051/0004-6361/202141443
18. "Refined physical parameters for Chariklo's body and rings from stellar occultations observed between 2013 and 2020", Morgado, B.E. et al. (2021), A&A, DOI: 10.1051/0004-6361/202141543
19. "The 2017 May 20 stellar occultation by the elongated centaur (95626) 2002 GZ32", Santos-Sanz, P. et al. (2021), MNRAS, DOI: 10.1093/mnras/staa3881
20. "Stellar occultations enable milliarcsecond astrometry for Trans-Neptunian objects and Centaurs", Rommel, F.L. et al. (2020), A&A, DOI: 10.1051/0004-6361/202039054
21. "A multi-chord stellar occultation by the large trans-Neptunian object (174567) Varda", Souami, D. et al. (2020), A&A, DOI: 10.1051/0004-6361/202038526
22. "The large trans-Neptunian object 2002 TC₃₀₂ from combined stellar occultation, photometry, and astrometry data". Ortiz, J.L. et al. (2020), A&A, DOI: 10.1051/0004-6361/202038046

PI and Co-Is publications relevant to this proposal

Publications that resulted from observations of our occultation proposals

1. "The Rings of (2060) Chiron: Evidence of an Evolving System", Pereira, C.L. et al. (2025), ApJL, DOI: 10.3847/2041-8213/ae0b6d
2. "Centaur 29P/Schwassmann-Wachmann 1 and its near-nucleus environment from a stellar occultation", Pereira, C.L. et al. (2025), Philosophical Transactions A, DOI: 10.1098/rsta.2024.0189
3. "Physical properties of Centaur (60558) 174P/Echeclus from stellar occultations", Pereira, C.L. et al. (2024), MNRAS, DOI: 10.1093/mnras/stad3318
4. "A large topographic feature on the surface of the trans-Neptunian object (307261) 2002 MS4 measured from stellar occultations", Rommel, F. et al. (2023), A&A, DOI: 10.1051/0004-6361/202346892
5. "The two rings of (50000) Quaoar", Pereira, C.L. et al. (2023), A&A, DOI: 10.1051/0004-6361/202346365
6. "Refined physical parameters for Chariklo's body and rings from stellar occultations observed between 2013 and 2020", Morgado, B.E. et al. (2021), A&A, DOI: 10.1051/0004-6361/202141543
7. "The first observed stellar occultations by the irregular satellite Phoebe (Saturn IX) and improved rotational period", Gomes-Júnior et al. (2020), MNRAS, DOI: 10.1093/mnras/stz3463
8. "Size and shape of Chariklo from multi-epoch stellar occultation", Leiva, R., Sicardy, B., Camargo, J. I. B., et al. (2017), AJ, DOI:10.3847/1538-3881/aa8956.
9. "The Structure of Chariklo's Rings from Stellar Occultations", Bérard, D., Sicardy, B., Camargo, J. I. B. et al. (2017), AJ, DOI:10.3847/1538-3881/aa830d.




Figure 1: Light curves (one for each SPARC4 channel) from the occultation event by Pholus (negative) on 2025JUN29. Time axis is in UTC. The vertical black line indicates where we should have seen the flux drop in case the event were positive.





Figure 2: Occultation profile from the PE observation on September 10, 2023 event of the normal optical depth as a function of radial distance in the ring plane shows the extent of the broad ring and the arrangement of the dense rings. The coloured regions indicate the boundaries of the spin-orbit resonances (SOR) 1/2 resonance (green), SOR 1/3 resonance (red), and the Roche limit (blue). The vertical dotted lines indicate the respective ring centers. See also **press release note** in <https://shorturl.at/T3Uae> and also <https://www.youtube.com/live/DKYDGWKFvk>. Pereira et al. (2025), ApJL DOI: 10.3847/2041-8213/ae0b6d

2007JH43, GaiaDR3+pmGaiaDR3, NIMAv10
 updated: 2025-08-22 by Lucky Star


Offset: 0.0mas 0.0mas



yyyy mm dd hh:mm:ss.s	RA_star_J2000	DE_star_J2000	C/A	P/A	vel	Delta	G*	RP*	H*
2026-05-15 01:29:20.7	17 14 29.8302	-27 54 39.379	0.011	177.97	-22.07	39.6057	16.6	15.7	13.4

Ixion, GaiaDR3+pmGaiaDR3, NIMAv13
 updated: 2025-08-24 by Lucky Star

Offset: 0.0mas 0.0mas



yyyy mm dd hh:mm:ss.s	RA_star_J2000	DE_star_J2000	C/A	P/A	vel	Delta	G*	RP*	H*
2026-07-25 04:35:04.8	18 32 34.6778	-32 32 26.356	0.004	358.36	-21.91	36.5285	15.9	15.1	13.3

Figure 3: Two of the events in this proposal. Links to the web pages of all events are provided in Technical Description.

Objects List.

Object	A.R.(h,m)	Dec.(deg)	Mag.	Band	Type	Comments
1996 ST1	12:29:58.05	-06:00:38.28	13.4	G	Sci	Occultation by 1996 ST1 on 2026MAR08T04:29:00 UTC
Comet B-B	05:02:31.90	-69:57:23.19	16.3	G	Sci	Occultation by comet B-B on 2026APR09T22:53:00 UTC
Orcus	10:36:47.44	-13:20:48.07	17.7	G	Sci	Occultation by Orcus on 2026MAY01T00:46:00 UTC
2014 GU53	17:01:56.59	+00:39:10.52	17.8	G	Sci	Occultation by 2014 GU53 on 2026MAY06T04:06:00 UTC
2004 EW95	16:20:13.68	-43:55:18.76	14.5	G	Sci	Occultation by 2005 EW95 on 2026MAY10T05:32:00 UTC
2007 JH43	17:14:29.83	-27:54:39.38	16.6	G	Sci	Occultation by 2007 JH43 on 2026MAY15T01:32:00 UTC
Ixion	18:35:15.22	-32:29:45.33	17.5	G	Sci	Occultation by Ixion on 2026JUN28T23:18:00 UTC
Ixion	18:32:34.68	-32:32:26.36	15.9	G	Sci	Occultation by Ixion on 2026JUL25T04:31:00 UTC
2002 KX14	17:58:00.13	-23:33:11.24	14.8	G	Sci	Occultation by 2002 KX14 on 2026AUG07T21:55:00 UTC