# **Reduce Function** ★



Q

Your Reduce Function submission got 30.00 points. Share Tweet

Try the next challenge | Try a Random Challenge

Problem Submissions Leaderboard Editorial △

Given a list of rational numbers, find their product.

#### Concept

The reduce() function applies a function of two arguments cumulatively on a list of objects in succession from left to right to reduce it to one value. Say you have a list, say [1,2,3] and you have to find its sum.

```
>>> reduce(lambda x, y : x + y,[1,2,3])
6
```

You can also define an initial value. If it is specified, the function will assume initial value as the value given, and then reduce. It is equivalent to adding the initial value at the beginning of the list. For example:

```
>>> reduce(lambda x, y : x + y, [1,2,3], -3)
3
>>> from fractions import gcd
>>> reduce(gcd, [2,4,8], 3)
1
```

#### **Input Format**

First line contains **n**, the number of rational numbers.

The  $i^{ ext{th}}$  of next n lines contain two integers each, the numerator(  $N_i$  ) and denominator(  $D_i$  ) of the  $i^{ ext{th}}$  rational number in the list.

## Constraints

- $1 \le n \le 100$
- $1 \le N_i, D_i \le 10^9$

## **Output Format**

Print only one line containing the numerator and denominator of the product of the numbers in the list in its simplest form, i.e. numerator and denominator have no common divisor other than 1.

# Sample Input 0

3 1 2

3 4

10 6

# Sample Output 0

5 8

## **Explanation 0**


Required product is  $\frac{1}{2}\frac{3}{4}\frac{10}{6}=\frac{5}{8}$ 



You have earned 30.00 points! 92/115 challenges solved.

#### 80%





⊗ Test case 4 🖰	Unlock
⊗ Test case 5 🖰	
⊗ Test case 6 🖰	
· ·	

Contest Calendar | Blog | Scoring | Environment | FAQ | About Us | Support | Careers | Terms Of Service | Privacy Policy | Request a Feature