

Course: Machine Learning - Foundations
Week 11 Questions

PRACTICE QUESTIONS

1. (1 point) The continuous random variable X represents the amount of sunshine in hours between noon and 8 pm at a skiing resort in the high season. The probability density function, $f(x)$, of X is modelled by

$$f(x) = \begin{cases} kx^2, & \text{for } 0 \leq x \leq 8 \\ 0, & \text{otherwise} \end{cases}$$

Find the probability that on a particular day in the high season there is more than two hours of sunshine between noon and 8 pm.

Answer: 0.98

Solution:

We know that $\int_{-\infty}^{\infty} f(x)dx = 1$

$$\Rightarrow \int_0^8 kx^2 dx = 1$$

value of k can be calculated while solving the above equation.
and then calculate $P(2 \leq X \leq 8) \Rightarrow \int_2^8 kx^2 dx$

2. (1 point) Let X be a continuous random variable with PDF

$$f_X(x) = \begin{cases} 6x + bx^2 & \text{for } 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

Calculate Value of b

Answer: -6

Solution:

We know that $\int_{-\infty}^{\infty} f(x)dx = 1$

$$\Rightarrow \int_0^1 6x + bx^2 dx = 1$$

value of b can be calculated while solving the above equation.

3. (1 point) Let X be a continuous random variable with PDF

$$f_X(x) = \begin{cases} 4x^k & \text{for } 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

find $E(X)$.

Answer: 0.8

We know that $\int_{-\infty}^{\infty} f(x)dx = 1$

$$\Rightarrow \int_0^1 4x^k = 1$$

value of k can be calculated while solving the above equation.i.e $k = 3$

$$E(X) = \int_0^1 x \times 4x^3 dx$$

4. (1 point) The time that passenger train will reach the station is uniformly distributed between 2:00 PM and 4:00 PM. What is the probability that the train reaches station exactly at 04:00 PM?

Answer: 0

The probability that train reaches exactly at 4:30 PM is $\int_{4:30}^{4:30} f(x)dx = 0$

Since the area under curve at a particular instant of x value is zero.

5. (1 point) If X is an exponential random variable with rate parameter λ then which of the following statement(s) is(are) correct.

- A. $P(X > x + k | X > k) = P(X > x)$ for $k, x \leq 0$.
- B. $P(X > x + k | X > k) = P(X > k)$ for $k, x \leq 0$.
- C. $P(X > x + k | X > k) = P(X > x)$ for $k, x \geq 0$.
- D. $P(X > x + k | X > k) = P(X > k)$ for $k, x \geq 0$.

Answer: C

Solution :

$$\begin{aligned} P(X > x + k | X > k) &= \frac{P((X > x + k) \cap (X > k))}{P(X > k)} \\ &\Rightarrow \frac{P(X > x + k)}{P(X > k)} = \frac{e^{-\lambda \times (x+k)}}{e^{-\lambda k}} \end{aligned}$$

$$\Rightarrow e^{-\lambda x}$$

Hence, Option C is correct

6. (1 point) The lifetime of a electric bulb is exponentially distributed with a mean life of 18 months. If there is 60% chances that a electric bulb will last mostly for t months, then what is the value of t ?

Solution:

Given mean of exponential random variable (life of light bulb)=18 months.

$$\Rightarrow \frac{1}{\lambda} = 18, \Rightarrow \lambda = \frac{1}{18}.$$

$$\text{Given } P(X \leq t) = 0.6$$

$$\Rightarrow 1 - e^{-\lambda t} = 0.6$$

$$\Rightarrow e^{-\lambda t} = 0.4$$

$$\lambda t = \ln 2.5$$

$$\frac{1}{18}t = \ln 2.5$$

$$t = 18 \ln 2.5$$

7. (1 point) **(Multiple Select)** Let X be uniformly distributed with parameters a and b , then which of the following is/are true:

A. $E(X^2) = \frac{(b^2 + a^2 + ab)}{3}$

B. $f(x) = \frac{-1}{(a-b)} \quad ; \quad a \leq x \leq b$

C. $E(X) = \frac{(b+a)}{2}$

D. $V(X) = \frac{(b-a)^2}{12}$

Answer: A, B, C, D

Solution:

From the properties of uniform distribution, options (a), (b), (c) and (d) are correct.

8. (1 point) **(Multiple Select)** Which of the following option is/are correct?

A. The shape of the Normal Curve is bell curved.

B. Normal Distribution is symmetric is about mean.

C. For a standard normal variate, the value of mean is 0.

D. The area under a standard normal curve is 1.

E. The standard normal curve is symmetric about the value 1.

Answer: A, B, C, D, E

9. (1 point) Let X and Y be continuous random variables with joint density

$$f_{XY}(x, y) = \begin{cases} cxy & \text{for } 0 < x < 1, 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Calculate $P(0 < X < \frac{1}{2}, 0 < Y < \frac{1}{2})$

Answer: 0.015

Solution:

We know that $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x, y) dy dx = 1$

$$\int_0^1 \int_0^1 cxy dy dx = 1$$

c can be calculated from above equation.

10. (points) Let X is a uniformly distributed random variable with $\mu_x = 15$ and $\sigma_x^2 = \frac{25}{3}$.
Calculate $P(X > 17)$

Answer: 0.5

$$E[X] = \frac{b+a}{2} = 15.$$

$$Var(X) = \frac{(b-a)^2}{12} = 12$$

a and b can be calculated from above equation.

Then calculate $P(X \geq 17)$

11. (1 point) Let X be a continuous random variable with PDF

$$f_X(x) = \begin{cases} ax & \text{for } 0 < x < 3 \\ a(6-x) & \text{for } 3 \leq x \leq 6 \\ 0 & \text{otherwise} \end{cases}$$

Calculate $P(0 \leq x \leq 4)$

Answer: 0.77

We know that $\int_{-\infty}^{\infty} f(x)dx = 1$

Solving above equation taking required $f(x)$ we will get, $a = \frac{1}{9}$

12. Let X be exponentially distributed with parameter λ , then which of the following is/are true about the variance of X :

- a. $V(X) = E[X^2] - (E[X])^2$
- b. $V(X) = E[X - E[X]]^2$
- c. $V(X) = (E[X])^2$
- d. $V(X) = E[X^2]$

Answer: A, B, C

Solution: From definition A and B are correct and for exponential distribution option C are correct.