



IIT Madras
ONLINE DEGREE

MACHINE LEARNING - FOUNDATIONS

REVISION (WEEK 2)

IIT Madras Online Degree

- Continuity
- Differentiability
- Linear Approximation
- Higher order approximations
- Multivariate Linear Approximation
- Directional Derivative

A function $f(x)$ is continuous at $x = a$ if:

$$f(a) = \lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x)$$

A function $f(x)$ is differentiable at $x = a$ if:

$$f'(a) = \lim_{x \rightarrow a^-} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a^+} \frac{f(x) - f(a)}{x - a}$$

Linear Approximation

The linear approximation $L(x)$ of a function $f(x)$ at point a is given by:

$$L(x) = f(a) + f'(a)(x - a)$$

Linear Approximation

The linear approximation $L(x)$ of a function $f(x)$ at point a is given by:

$$L(x) = f(a) + f'(a)(x - a)$$

This is indeed the equation of a tangent line:

$$y - y_1 = m(x - x_1)$$

$$y = y_1 + m(x - x_1)$$

Linear Approximation

The linear approximation $L(x)$ of a function $f(x)$ at point a is given by:

$$L(x) = f(a) + f'(a)(x - a)$$

This is indeed the equation of a tangent line:

$$y - y_1 = m(x - x_1)$$

$$y = y_1 + m(x - x_1)$$

If $x_1 = a$, $y_1 = f(a)$ and $m = f'(a)$, we get,

$$y = f(a) + f'(a)(x - a)$$

Higher order approximations

Linear Approximation

$$L(x) = f(a) + f'(a)(x - a)$$

Higher order approximations

Linear Approximation

$$L(x) = f(a) + f'(a)(x - a)$$

Quadratic Approximation

$$L(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2$$

Higher order approximations

Linear Approximation

$$L(x) = f(a) + f'(a)(x - a)$$

Quadratic Approximation

$$L(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2$$

Higher-order Approximations

$$L(x) = f(a) + f^{(1)}(a)(x - a) + \frac{f^{(2)}(a)}{2}(x - a)^2 + \\ + \frac{f^{(3)}(a)}{3 \cdot 2}(x - a)^3 + \frac{f^{(4)}(a)}{4 \cdot 3 \cdot 2}(x - a)^4 \dots$$

Multivariate linear approximation: Linear approximation of functions involving multiple variables

The linear approximation of a function f of two variables x and y in the neighborhood of (a, b) is:

$$L(x, y) = f(a, b) + \frac{\partial f}{\partial x}(a, b)(x - a) + \frac{\partial f}{\partial y}(a, b)(y - b)$$

DIRECTIONAL DERIVATIVES

Directional Derivative

$$\cdot f_x(x, y) = \frac{\partial f}{\partial x}(x, y)$$

Directional Derivative

- $f_x(x, y) = \frac{\partial f}{\partial x}(x, y)$ = Rate of change of f as we vary x (keeping y fixed).

Directional Derivative

- $f_x(x, y) = \frac{\partial f}{\partial x}(x, y)$ = Rate of change of f as we vary x (keeping y fixed).
- $f_y(x, y) = \frac{\partial f}{\partial y}(x, y)$

Directional Derivative

- $f_x(x, y) = \frac{\partial f}{\partial x}(x, y)$ = Rate of change of f as we vary x (keeping y fixed).
- $f_y(x, y) = \frac{\partial f}{\partial y}(x, y)$ = Rate of change of f as we vary y (keeping x fixed).

Directional Derivative

- $f_x(x, y) = \frac{\partial f}{\partial x}(x, y)$ = Rate of change of f as we vary x (keeping y fixed).
- $f_y(x, y) = \frac{\partial f}{\partial y}(x, y)$ = Rate of change of f as we vary y (keeping x fixed).
- Directional derivative of $f(x, y)$

Directional Derivative

- $f_x(x, y) = \frac{\partial f}{\partial x}(x, y)$ = Rate of change of f as we vary x (keeping y fixed).
- $f_y(x, y) = \frac{\partial f}{\partial y}(x, y)$ = Rate of change of f as we vary y (keeping x fixed).
- Directional derivative of $f(x, y)$ = Rate of change of f if we allow both x and y to change simultaneously (in some direction (u)).

Directional Derivative

- $f_x(x, y) = \frac{\partial f}{\partial x}(x, y)$ = Rate of change of f as we vary x (keeping y fixed).
- $f_y(x, y) = \frac{\partial f}{\partial y}(x, y)$ = Rate of change of f as we vary y (keeping x fixed).
- Directional derivative of $f(x, y)$ = Rate of change of f if we allow both x and y to change simultaneously (in some direction (u)).

$$\begin{aligned} D_{\vec{u}}f(x, y) &= \nabla f \cdot u \\ &= \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right] \cdot [u_1, u_2] \\ &= u_1 \frac{\partial f}{\partial x} + u_2 \frac{\partial f}{\partial y} \end{aligned}$$

Directional Derivative

- $f_x(x, y) = \frac{\partial f}{\partial x}(x, y)$ = Rate of change of f as we vary x (keeping y fixed).
- $f_y(x, y) = \frac{\partial f}{\partial y}(x, y)$ = Rate of change of f as we vary y (keeping x fixed).
- Directional derivative of $f(x, y)$ = Rate of change of f if we allow both x and y to change simultaneously (in some direction (u)).

$$\begin{aligned} D_{\vec{u}}f(x, y) &= \nabla f \cdot \vec{u} \\ &= \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right] \cdot [u_1, u_2] \\ &= u_1 \frac{\partial f}{\partial x} + u_2 \frac{\partial f}{\partial y} \end{aligned}$$

Directional derivative can be considered to be a weighted sum of partial derivatives.

Thank you.