

Relation Web

- Let D be the set of descriptors.
- Let R be the set of possible relations.
- Let $G = (V, E)$ be a directed graph where $V = D$ and E is a set of edges such that each $v \in V$ has $|R| \times |D|$ edges, one per relation per descriptor (exhaustively listing all possible relations between this descriptor and all other descriptors).
- Goal: assign each edge $e \in E$ a weight $P(e) \in [0, 1]$ that represents the probability of that being a valid relation.
- Idea: as triples are processed, adjust the weight of not only the “explicit” relation between the relevant descriptors that is directly stated with that triple, but also the “implicit” relation(s).

Algorithm

- Assume that we have a stream S of verified confidence measures for specific relations between two descriptors.
- We want to sequentially construct a relation web, processing one $s \in S$ at a time, updating the web for each new piece of information.
- Let L be the set of labels.
- Let D be the set of descriptors.
- Let L_p be the set of all unique ordered pairwise combinations of $l \in L$.
- Let $C(l_p)$, where $l_p = (l_1, l_2) \in L_p$, be the set of all possible paths between l_1 and l_2 , where each path is generalized to the labels on the vertices.
- Let $C_p(l_p)$, be the set of all unique unordered pairwise combinations of possible paths between l_1 and l_2 , where each path is generalized to the labels on the vertices.
- Let $L(d)$, where $d \in D$, be the label of d .
- Let $L(c^*)$, where c^* is a path in the graph, be a path with the descriptors replaced with their labels.
- Let $Corr_{l_p}(c_p)$, where $c_p = (c_1, c_2) \in C_p(l_p)$, be an assignment of a correlation score between paths c_1 and c_2 .
- When processing a new measure $s \in S$:
Extrapolation Stage:
 - Assign edge e specified by s .
 - For each path $d_1 \xrightarrow{r_{1,2}} \dots \xrightarrow{r_{n-1,n}} d_n = c_1^*$ that e is a part of:
 - * For each path c_2^* starting at d_1 and ending at d_n :
 - (Recursively?) update weights along c_2^* according to $Corr_{(L(d_1), L(d_2))}((L(c_1^*), L(c_2^*)))$.

Correlation Reassessment Stage:

- Reassess correlations $(Corr_{l_p}(c_p) \forall l_p, c_p)$ based on current state of graph.
- For each pair of labels (l_1, l_2) :
 - * Initialize list of vectors V .
 - * For each pair of descriptors (d_1, d_2) with labels (l_1, l_2) :
 - Construct a vector v where each entry corresponds to a label-generalized path between the descriptors, and store the sum of the edge weights along that path.
 - Add v to V .
 - * Using V , determine pairwise correlations between all label-generalized paths c_1, c_2 , ignoring paths with uninitialized values, storing results in $Corr_{(l_1, l_2)}(c_1, c_2)$.

Event: Initialization

X_1

X_2

Y_1

Z_1

Y_2

Z_2

X_3

Y_3

Z_3

Event: Recieved from S : Confidence of 1.0 for relation 1 between X_1 and Y_1 .

Extrapolation Stage:


- Because there are not any correlations yet, no other edges are updated.

Correlation Reassessment Stage:

- Correlation vectors:

$X \rightarrow Y$				$Y \rightarrow Z$			$X \rightarrow Z$		
$x \perp\!\!\!\rightarrow y$	$x_1 \rightarrow y_1$ 1	$x_2 \rightarrow y_2$	$x_3 \rightarrow y_3$	$y_1 \rightarrow z_1$	$y_2 \rightarrow z_2$	$y_3 \rightarrow z_3$	$x_1 \rightarrow z_1$	$x_2 \rightarrow z_2$	$x_3 \rightarrow z_3$

- No correlations can be drawn.



Event: Recieved from S : Confidence of 1.0 for relation 2 between Y_1 and Z_1 .

Extrapolation Stage:


- Because there are not any correlations yet, no other edges are updated.

Correlation Reassessment Stage:

- Correlation vectors:

$X \rightarrow Y$				$Y \rightarrow Z$				$X \rightarrow Z$			
$X \xrightarrow{1} Y$	$X_1 \rightarrow Y_1$ 1	$X_2 \rightarrow Y_2$	$X_3 \rightarrow Y_3$	$Y \xrightarrow{2} Z$	$Y_1 \rightarrow Z_1$ 1	$Y_2 \rightarrow Z_2$	$Y_3 \rightarrow Z_3$	$X \xrightarrow{1} Y \xrightarrow{2} Z$	$X_1 \rightarrow Z_1$ 2	$X_2 \rightarrow Z_2$	$X_3 \rightarrow Z_3$

- No correlations can be drawn.



Event: Recieved from S : Confidence of 1.0 for relation 3 between X_1 and Z_1 .

Extrapolation Stage:


- Because there are not any correlations yet, no other edges are updated.

Correlation Reassessment Stage:

- Correlation vectors:

$X \rightarrow Y$				$Y \rightarrow Z$				$X \rightarrow Z$			
$X \xrightarrow{1} Y$	$X_1 \xrightarrow{1} Y_1$	$X_2 \rightarrow Y_2$	$X_3 \rightarrow Y_3$	$Y \xrightarrow{2} Z$	$Y_1 \xrightarrow{1} Z_1$	$Y_2 \rightarrow Z_2$	$Y_3 \rightarrow Z_3$	$X \xrightarrow{3} Z$ $X \xrightarrow{1} Y \xrightarrow{2} Z$	$X_1 \xrightarrow{1} Z_1$	$X_2 \rightarrow Z_2$	$X_3 \rightarrow Z_3$

- No correlations can be drawn.



Event: Recieved from S : Confidence of 1.0 for relation 1 between X_2 and Y_2 , confidence of 1.0 for relation 3 between X_2 and Y_2 .

Extrapolation Stage:


- Because there are not any correlations yet, no other edges are updated.

Correlation Reassessment Stage:

- Correlation vectors:

$X \rightarrow Y$				$Y \rightarrow Z$				$X \rightarrow Z$			
$X \xrightarrow{1} Y$	$X_1 \xrightarrow{1} Y_1$	$X_2 \xrightarrow{1} Y_2$	$X_3 \rightarrow Y_3$	$Y \xrightarrow{2} Z$	$Y_1 \xrightarrow{1} Z_1$	$Y_2 \xrightarrow{1} Z_2$	$Y_3 \rightarrow Z_3$	$X \xrightarrow{3} Z$ $X \xrightarrow{1} Y \xrightarrow{2} Z$	$X_1 \xrightarrow{1} Z_1$	$X_2 \rightarrow Z_2$	$X_3 \rightarrow Z_3$
	1	1			1	1			1 2	2	

- No correlations can be drawn.



Event: Recieved from S : Confidence of 1.0 for relation 1 between X_2 and Z_2 .

Extrapolation Stage:


- Because there are not any correlations yet, no other edges are updated.

Correlation Reassessment Stage:

- Correlation vectors:

$X \rightarrow Y$				$Y \rightarrow Z$				$X \rightarrow Z$			
$X \xrightarrow{1} Y$	$X_1 \rightarrow Y_1$ 1	$X_2 \rightarrow Y_2$ 1	$X_3 \rightarrow Y_3$	$Y \xrightarrow{2} Z$	$Y_1 \rightarrow Z_1$ 1	$Y_2 \rightarrow Z_2$ 1	$Y_3 \rightarrow Z_3$	$X \xrightarrow{3} Z$ $X \xrightarrow{1} Y \xrightarrow{2} Z$	$X_1 \rightarrow Z_1$ 1 2	$X_2 \rightarrow Z_2$ 1 2	$X_3 \rightarrow Z_3$

- A positive correlation now exists between $X \xrightarrow{3} Z$ and $X \xrightarrow{1} Y \xrightarrow{2} Z$. That is, $Corr_{(X,Z)}(X \xrightarrow{3} Z, X \xrightarrow{1} Y \xrightarrow{2} Z) > 0$.



Event: Recieved from S : Confidence of 1.0 for relation 1 between X_3 and Y_3 .

Extrapolation Stage:




- Because the path $X_3 \xrightarrow{1} Y_3 \xrightarrow{2} Z$ is incomplete, no extrapolation occurs.

Correlation Reassessment Stage:

- Correlation vectors:

$X \rightarrow Y$				$Y \rightarrow Z$				$X \rightarrow Z$			
$X \xrightarrow{1} Y$	$X_1 \rightarrow Y_1$	$X_2 \rightarrow Y_2$	$X_3 \rightarrow Y_3$	$Y \xrightarrow{2} Z$	$Y_1 \rightarrow Z_1$	$Y_2 \rightarrow Z_2$	$Y_3 \rightarrow Z_3$	$X \xrightarrow{3} Z$	$X_1 \rightarrow Z_1$	$X_2 \rightarrow Z_2$	$X_3 \rightarrow Z_3$
	1	1	1		1	1		$X \xrightarrow{1} Y \xrightarrow{2} Z$	1 2	1 2	

- $Corr_{(X,Z)}(X \xrightarrow{3} Z, X \xrightarrow{1} Y \xrightarrow{2} Z) > 0$.
- No new correlations can be drawn.



Event: Recieved from S : Confidence of 1.0 for relation 2 between X_3 and Y_3 .

Extrapolation Stage:

- Because the path $X_3 \xrightarrow{1} Y_3 \xrightarrow{2} Z_3$ is complete, extrapolation occurs between it and the path $X_3 \xrightarrow{3} Z_3$, using $Corr_{(X,Z)}(X \xrightarrow{3} Z, X \xrightarrow{1} Y \xrightarrow{2} Z)$. A weight σ is applied to the correlated path $X_3 \xrightarrow{3} Z_3$.

Correlation Reassessment Stage:

- Correlation vectors:

$X \rightarrow Y$				$Y \rightarrow Z$				$X \rightarrow Z$			
$X \xrightarrow{1} Y$	$X_1 \rightarrow Y_1$	$X_2 \rightarrow Y_2$	$X_3 \rightarrow Y_3$	$Y \xrightarrow{2} Z$	$Y_1 \rightarrow Z_1$	$Y_2 \rightarrow Z_2$	$Y_3 \rightarrow Z_3$	$X \xrightarrow{3} Z$	$X_1 \rightarrow Z_1$	$X_2 \rightarrow Z_2$	$X_3 \rightarrow Z_3$
	1	1	1		1	1	1	$X \xrightarrow{1} Y \xrightarrow{2} Z$	1	1	σ
									2	2	2

- $Corr_{(X,Z)}(X \xrightarrow{3} Z, X \xrightarrow{1} Y \xrightarrow{2} Z) > 0$.
- $Corr_{(X,Z)}(X \xrightarrow{3} Z, X \xrightarrow{1} Y \xrightarrow{2} Z)$ is adjusted.

