(Saathi)

1534 Written Assignment 1 19 E: ~N(0,0;) = 4: -W'X; +E; P(Y/X;N) = T 1 = -(y, - Wx,)2/45,2 * 7 = L(W) N(W) = log (I(W)) - log | T 1 e - (4-4/x,) =] $=\frac{1}{\sqrt{20}}\sum_{i,j}\log\left(\frac{1}{\sigma_{i}}\right) \neq -\left(\frac{1}{2}-W_{i}\right)$ b argmax $I(\omega) = 1 \int log(1) - lgi - Wx, 1$ $Van = 1 \int log(\sigma_i) - lgi - Wx, 1$ Constant constant= argmax $\frac{N}{I-I} - \left(\frac{y_i - W^T x_i}{2\sigma_i^2}\right)^2$ = argmin 1 5 1 (Nx-y). - 1 5 ai (NTxi-yi)2

where ai . 1

C. argmax (l(w)) = argmin - 5 ai (yi-Wx) THE Saily-WXi). (-xi) = \(\int ai \left(w \text{xi} - yi \right) \text{xi} = \(\int ai \left(\hat{y} - yi \right) \text{xi} \) d. Maxii Let \(\times_{00} \times_{01} \cdot \times_{0d-1} \) \\
\times_{10} \times_{11} \cdot \times_{10} \cdot \times_{11} \cdot \times_{1d-1} $\begin{bmatrix} N_0 \\ W_1 \end{bmatrix} = W \qquad \begin{bmatrix} y_0 \\ y_1 \end{bmatrix} - y$ $\begin{bmatrix} W_{d-1} \end{bmatrix}$ 4MA WWW Let Ta, a. and A = asgmin Allxw-yll = argmin AO (XW-Y) O= dot operation Since A is a constant it will not affect the minimum => argmin 1XW-YII = [WTXTXW-QWTXTY+YTY] $\gg W = (X^T X) X^T Y.$

Page No.

(Saathi) Date $-\frac{1}{N}$ $\frac{1}{N}$ $\frac{1}{N}$ log(L(W)) = 5 (5 log P(y = K/Xi) I(y = K) = 5 (5 I(y=k) log P(y=k/xi)) = 5 (5](y;=k). log (k x)) = 5 5 7 (y=k) (log(e WE'x) - log(5 e WE'x)) Since I(y=k) is 1 only for a particular value of k= ko, we can say $\frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} = \frac{$ $\nabla \ell(w) = \frac{1}{2} \frac{1$ $= \chi \left[\frac{\sum_{k=1}^{K} e^{W_{k}} \times - e^{W_{k}} \times - e^{W_{k}} \times }{\sum_{k=1}^{K} e^{W_{k}} \times } \right]$

Page No.

 $\frac{P(x_1) \cdot P(x_2) \cdot \cdot P(x_n)}{P(x_1) \cdot (1-\alpha)}$ $= \frac{P(x_1) \cdot P(x_2) \cdot \cdot P(x_n)}{P(x_1)}$ $= \frac{P(x_1) \cdot P(x_2) \cdot \cdot P(x_n)}{P(x_n)}$

Date ___ /___ /



NON Since P(O(x, xn) is a PDF $\int P(O(x, xn) = 1$

 $\frac{p(\theta|X)}{p(\theta|X)} = \frac{e^{-1+\sum_{i=1}^{N}(i-x_i)}}{e^{-1+\sum_{i=1}^{N}(i-x_i)}}$

B(d, p). To P(Xi)

Comparing this to the Beta Function for P(0) we see $P(0|X) = Beta(X + \sum Xi, \beta + \sum (1-Xi))$ And $P(Xi) = B(X + \sum Xi, \beta + \sum (1-Xi))$ $P(Xi) = B(X + \sum Xi, \beta + \sum (1-Xi))$ $P(Xi) = B(X + \sum Xi, \beta + \sum (1-Xi))$

B(x,B)

b- P(0) = Beta (2,2)

If me observe 5# 2 Heads out of 5 tosses:

P(0 | x) = Beta (2+2, 2+3) = Beta (4,5)

If m observe \$20 Heads out of 50 forses

P(O(X) = Beta(22, 32)