



भारत सरकार

Government of India

कृषि और किसान कल्याण मंत्रालय

Ministry of Agriculture & Farmers Welfare

कृषि एवं किसान कल्याण विभाग

Department of Agriculture & Farmers Welfare

वनस्पति संरक्षण, संगरोध एवं संग्रह निदेशालय

Directorate of Plant Protection, Quarantine & Storage

केंद्रीय कीटनाशी बोर्ड एवं पंजीकरण समिति

Central Insecticides Board & Registration Committee

एनएच IV, फरीदाबाद-121 001 (हरियाणा)

N.H. IV, Faridabad-121 001 (Haryana)

Major Uses of Pesticides

(Registered under the Insecticides Act, 1968)

(UPTO – 31/03/2026)

(Based on certificates issued)

Disclaimer: The document has been compiled on the basis of available information for guidance and not for legal purposes.

PLANT GROWTH REGULATORS (PGR)

Plant Growth Regulators (PGR): (Page No. 2 to 12)

APPROVED USES OF REGISTERED PGR

PLANT GROWTH REGULATORS (PGR)

Name of PGR & approved Crops	Time of application / purpose	Dosage/ha		Dilution In Water (Litres) / Preparation of solution	Waiting period / PHI between last application & harvest (days)
		a.i. (ppm/gm/%)	Formulation (ml/gm/Ltr/kg/%)		
Alpha Naphthyl Acetic Acid 4.5% SL (Na salt)					
Tomato	At the time of flowering two spray.	45 ppm	-	-	-
Chillies	1 st spray during flowering & 2 nd spray 20 -30 days later.	10 ppm	-	-	-
Mango	1 st spray when tender fruits one of pea size. 2 nd spray when fruits one of marble size (about 2 cm diameter)	20 ppm	-	2 ml in 4.5litre.	-
	<u>To control Mango malformation</u> - Before fruit bud differentiations approx. 3 months before flowering	200 ppm	-	20 ml in 4.5 ltrs.	-
Grapes	(a)To increase size & weight of berries. – 1 st sprays at pruning time. – 2 nd spray when flowering shoot appear	10 ppm	-	2 ml in 49 ltrs.	-
	(b)To control berry drop (spray on matured grape bunches) 10-15 days before harvesting.	100 ppm	-	20 ml. in 49 Litres	-
Pineapple	(a) To induce flowering and uniform growth	10 ppm (In dry weather half strength solution i.e. 5 ppm may be used)	-	1 ml in 4.5 Litres (pour 30-50 ml of solution in to the head of each plant)	-
	(b) To increase fruit size.	199 ppm	-	10 ml in 4.5 Litres (spray to wet the whole	-

	To delay maturity - Two weeks before harvest.	100 ppm	-	plant) 10 ml in 4.5 Litres (Wet the whole fruit 2 weeks before harvest)	-
Cotton	To prevent shedding of flower squares & bolls (3 sprays at 15 days interval from square formation stage	10-20 ppm	222-444 ml	1000 ltr.	-
Chlormequat Chloride 50% SL					
Cotton (American)	Square formation of early flowering (one spray)	20-40 gm a.i./ha	40-80 ml/ha	High Volume 375-600 Low volume 125-187	-
Cotton (Desi)	Square formation of early flowering (one spray)	75 a.i. gm/ha	150 ml/ha	High volume 375-600	-
Brinjal	Seed soaking for 24 hours (before sowing)	50 ppm	100ppm	1ml/ 10L water	-
Potato	Dipping of cut pieces for 10 minutes	100 ppm	200ppm	2.0ml/ 10 L water	
Grapes 1 st spray:	3-5 leaf stage after April pruning	500 g a.i./ha	1000ml	1000 L	91
2 nd Spray:	5-7 leaf stage after April Pruning	1000 g a.i./ha	2000 ml		
3 rd Spray:	3-5 leaf stage after October Pruning	250 g a.i./ha	500ml		
Chlorpropham 50% HN					
Potato	Antisprouting agent for stocked potatoes under cold storage condition Temp= 10±2°C R.H.= 90±5%	18-20 gm/MT	36-40 ml/MT	Formulation is to be applied as such with fogging applicator	20
Chlorpropham 55.37% w/w (624 g/l) HN					
Potato	Anti-sprouting agent for stocked potatoes under cold storage condition Temp= 10±2°C R.H.= 90±5%	18-20 gm/MT	29-32 ml/MT	Formulation is to be applied as such with fogging applicator	20
Chlorpropham 300 g/l EC					
Potato	Anti-sprouting agent for	18	60	1 L	Not applicable

	potatoes in heap storage conditions				
Ethephon 10% Paste					
Rubber	For renewed bark 4 times bark swabbing. During March, August, September & November below the tapping panel after 4 cm scrap of the bark /above the tapping panel/on the tapping cut after removing the lace.	10%	50 ml. formulation per tree directly used without dilution.	-	-
Ethephon 39 % SL					Method of preparation of required concentration
Mango	a) For breaking alternate bearing tendencies	200 ppm	770-1025	1500-2000	5 ml in 10 lit of water
	b) For Flower induction in juvenile mango	1000 ppm	3846-5128	1500-2000	26 ml in 10 lit of water
	c) Post-harvest treatment (For Uniform Ripening)	500 ppm	1923-2564	1500-2000	13 ml in 10 lit of water
Pineapple	For flower induction	100 ppm	385-513	1500-2000	2.5 ml in 10 lit of water
Coffee (Arabica)	For uniform ripening of berries, One spray at fly pricking stage, when 10-15% berries are ripened.	192 ppm	738-985	1500-2000	5 ml in 10 lit of water
Coffee (Robusta)	For uniform ripening of berries, one spray at fly pricking stage, when 10-15% berries are ripened.	96 ppm	215-287	1500-2000	2.5 ml in 10 lit of water
Tomato	Post-harvest treatment (for Uniform Ripening)	2500 ppm	-	-	65 ml in 10 lit of water
Rubber	Yielding rubber latex	1000 ppm	-	-	26 ml in 10 lit of water
Pomegranate	Defoliation for better flowering and fruit yield	390-487.5 g	1000-1250 ml	500	135 days (2-2.5 ml/lit water)
Grape	Defoliation for better flowering and fruit yield	487.5-682.5 g	1250-1750	750	2.5 ml in 1 lit of water
Ethephon 20% WP					
Mango	Post-harvest Ripening	1.2	6 gm	NA	NA
Forchlorfenuron 0.1% L (w/v)					
Grapes	Two dipping applications. 1 st When size of berry is 3-4 mm diameter and 2 nd When size of berry is 6-	2 ppm.	1 ltrs.	500	60 days

	7 mm diameter,				
Forchlorfenuron 0.12% EC w/w					
Grapes	To enhance the fruit size in seedless grapes single directed spray on berries at 4-6 mm berry size	3 ppm	1.5 liter	500 liter/ha.	20
Pigeon pea (Tur)	Single directed spray at the time of 100% flowering	2.5ppm	1.125 Litres/ha	Spray Volume- 450 l/ha. Mix 250 ml of Sitofex in 100 l water	30 days
Gibberellic Acid Technical (90% w/w)					
Grape fruit	a) At full bloom (for fruit set)-single spray b) 1 st week of May (For June fruit drop) –single spray c) 1 st week of October (For pre-harvest drop)-single spray	500-1000 ppm	-	-	-
Sweet cherry	When more than 60% buds opened fully.	40-80 ppm	-	-	-
Grapes	Two directed spray 1 st at full bloom & 2 nd at fruit set stages.	100 ppm	-	-	-
Grape (Seedless)	Two blanket spray at 1 st full bloom & 2 nd at post bloom stage.	15-60 ppm	-	-	-
Brinjal	a) seed treatment (dipping)	10 ppm	-	-	-
	b) When 4 weeks old - weekly spray	50 ppm	-	-	-
Gibberellic Acid 0.001% L					
Paddy	To increase the yield and quality of the crop produce				
	Short duration varieties 20-25DAT Medium duration varieties 30-35 DAT Long duration varieties 40-45 DAT	0.018 gm	180 ml	450-500	-
Sugarcane (Planted crops)	a) First spray 40-45 DAP b) Second spray 70-80 DAS	0.018 gm	180 ml	450-500	-
Cotton	a) First spray 40-45 DAP b) Second spray: At the time of ball formation	0.018 gm	180 ml	450-500	-
Groundnut	a) First spray at flowering	0.018 gm	180 ml	450-500	-

	(30-35 AS) b) Second spray at the time of flowering				
Banana	a) First spray 3 rd month b) Second spray 5 th month Third spray at the time of fruit formation	0.027 gm	270 ml.	450-500	-
Tomato / Potato / Cabbage / Cauliflower	a) First spray 45 DAS b) Second spray 65 DAS	0.018 gm	180 ml.	450-500	-
Onion	25-30 days after planting	0.018 g	180	450-500	-
Grapes	a) First spray 30-35 days after pruning b) Second during the match head stage	0.018 gm	180 ml.	450-500	-
Brinjal, Bhindi	a) First spray 34 DAP b) Second spray 70 DAP c) Third spray 105 DAP	0.045 gm	450 ml.	450-500	-
Tea	Five spray at monthly interval	-	270ml	450-500	-
Mulberry	First spray: 15-20 days after harvest	0.045	450	450-500	-
Gibberellic Acid 0.1% GR					
Rice	Broadcast (Manual by hand with rubber gloves or through mechanical dispenser) at 15-20 days after transplanting	12.5-15 g	12.5-15.0 kg	-	76
Gibberellic Acid 0.186% SP					
Cotton	To improve fiber quality one spray at square formation or early flowering stage	142 ppm.	71 gm	450-500	-
Gibberellic Acid 40% WSG					
Grape	Pre-Bloom Elongation	40	50	500	-
	Fruit Setting Thinning				
	6-7 mm berry size-enlargement				
Rice	20-25 Days After Transplanting	20-25	50-62.5	500	
	At Panicle emergence	20-25	50-62.5	500	
Wheat	20-25 Days After sowing	10-15	25-37.5	500	-
	10% ear emergence				
Maize	Knee high stage (25-30 DAS)	20	50	500	-
Gibberellic acid 0.45% SL					
Apple	Time of application 1. Pink	7.2 ppm	2ml/Liter	8-12 L (As per	24 hours

	bud stage 2. Pea nut Stage 3. Walnut Stage			plant canopy)	
Hydrogen Cyanamide 50% SL (Import)					
Grapes	For breaking bud dormancy Single application as spray Just after pruning.	1-1.5%	2-3%	375-500	90-120 days
Hydrogen Cyanamide 50% SL (Indigenous manufacture)					
Grapes	For breaking dormancy of fruiting buds Just after pruning, single application by swabbing.	1.5%	1.5 ltrs.	Mix with 200- 300 ml. of product in 10 litres of water.	120 days
Hydrogen Cyanamide 49% AS					
Grapes	For breaking bud dormancy One directed spray, just after pruning.	1.0-1.5%	2-3%	50 ltrs.	110 days
Sugarcane	Dipping of setts	0.50	1.00%	Mix 1000 ml of the product per 100 litres of water	319 days
Apple	For early initiation of bud sprouting and improving the fruit quality and enhance the yield of Apple. One-time single application four weeks prior to expected bloom	1.00%	2.00%	Mix 2000 ml of the product per 100 liters of water	188 days
Mepiquat chloride 5% AS					
Potato	One spray 45 DAP To restrict the excessive vegetative growth of potato and increasing its yield	62.5-75 g	1.25-1.50 ltr	500-600	60-90 days
Cotton	Single spray at flowering stage to control excessive vegetative growth and to increase crop yield in cotton	50-62.5 g	1.00-1.25 ltr	500-600	57
Groundnut	Single spray at flowering stage to control excessive vegetative growth and to increase crop yield in groundnut	50-62.5 g	1.00-1.25 ltr	500	60
Chickpea	Single spray at flowering stage to control excessive vegetative growth and to	62.5 g	1.25 ltr	500	56

	increase crop yield in chickpea				
Soybean	Single spray at flowering stage to control excessive vegetative growth and to increase crop yield in soybean	62.5 g	1.25 ltr	500	54
Brinjal	Single spray at flowering stage to control excessive vegetative growth and to increase crop yield in brinjal	62.5 g	1.25 ltr	500	7
Onion	Single spray at flowering stage to control excessive vegetative growth and to increase crop yield in onion	62.5 g	1.25 ltr	500	48 (bulb) 7 (green leaves)
Rice	Sprayed at mid tillering stage. For optimum plant height with more no. of tillers per hill. To enhance panicle length for better quality grain and yield in rice.	62.5	1.25	500	39
Wheat	Sprayed at vegetative growth stage/ majority of tillers. For more number of tillers and increase in panicle length. For better quality grains and improved grain and straw yield.	62.5	1.25	500	47
Cumin	Sprayed at vegetative growth stage. To reduce plant height and increase the no. of branches. For better quality grains and yield.	62.5	1.25	500	45
Okra	Spray at initiation of flowering. To increase number of fruits and fruit weight. For better quality of fruits and yield.	62.5	1.25	500	7
Nitrobenzene 20% w/w EW					
Tomato	Plant Growth Regulator	200 g	1000	500	Zero days
1-Methylcyclopropene 3.3% VP (Vapour Releasing Product)					

Apple fruit (Under ambient and cold condition)	Applied as soon as possible after harvest, within a maximum of 7 days after harvest on fruits kept at ambient and cold temperature away from source of external ethylene.	2.24 mg	68 mg (1000 PPB)	-	1
1-methylcyclopropene 0.014% VP					
Apple	-	0.0875 g (2000 ppb)	625 mg (/10 kg apple)	20 L	2 days
Paclobutrazol 23% w/w (25% w/v) SC (Import Source:- ZENECA Agrochemicals, Fernhurst, Haslemere, Surrey, UK)					
Mango	To reduce the inter node length of new shoots and earlier formation of terminal bud. Favourably, influence the fruit bud production, fruit colour and harvest yield 7-15 yrs old 16-25 yrs old >25 yrs old Application after the harvest of fruits (Any time from July to Oct)	- - -	15 ml. Per tree 20 ml. Per tree. 25-40 ml. Per tree (Note: If the soil is sandy the rate of application may be reduced to 75 % of the recommended . For repeat use the rate of application can be 50 to 75 % of the rate used in the 1st year)	Recommended quantity diluted in clean water of 5-10 lit. and applied in furrow 5 to 10 cm deep about 30 cm away from the trunk. Fill up with soil after application or apply as soil – collar drench. -	
Paclobutrazol 23% SC (W/W) / (25% W/V) (Import Source:- PGR International Pty. Ltd., 4 Dairy road, Werribee Vic. 3030 Australia)					
Mango	To reduce the inter node length of new shoots and earlier formation of	4.0 gm per tree	16 ml. Per tree	Make a round furrow about 5 to 10 cms deep at	Waiting Period - NIL as the chemical is

	terminal bud. increase fruit bud production, and improve fruit yield texture 16-25 yrs old Application after the harvest of fruits (Any time from July to Oct)		(Note: If the soil is sandy the rate of application be reduced to 75 % of the recommended . For repeat use the rate of application can be 50 to 75 % of the rate used in 1 st year)	least 30 cm away from the trunk. Mix the recommended dose with about 5-10 litres of clean water and apply to the furrow. Fill up with soil after application and irrigate once or twice a month subsequently	applied 8 months before harvest of fruits
Paclobutrazol 23% SC (w/w) / (25% w/v) (Indigenous manufacture)					
Mango	To reduce the inter node length of new shoots and earlier formation of terminal bud. Favourably, influence the fruit bud production, fruit colour and harvest yield 7-15 yrs old 16-25 yrs old >25 yrs old Application after the harvest of fruits (Any time from July to Oct)	3.45 4.6 5.75-9.2	15 ml per tree 20 ml per tree 30 ml per tree (Note: If the soil is sandy the rate of application may be reduced to 75 % of the recommended . For repeat use the rate of application can be 50 to 75 % of the rate used in the 1 st year)	Recommended quantity diluted in clean water of 5 litres and applied in furrow 5 to 10 cm deep about 30 cm away from the trunk. Fill up with soil after application or apply as soil – collar drench.	-
Pomegranate	To induce flowering and enhance yield	0.69 g a.i./ha	3.0 ml/tree	2L Recommended as soil drench	83

				(single application) ring form furrow to be made at a depth of 5-7 cm around plants and soil drenching to be done in active root zone and covered with soil.	
Apple	To induce flowering and enhance yield	2.3 g a.i./tree	10 ml/tree	5L Recommended as soil drench (single application) Treatment should be drenched in soil in circular area 25 cm away from tree stem. (Dormant stage)	155
	To induce flowering and enhance yield	0.46 g a.i./ Litre water (460 ppm)	2 ml/Litre (2000 ppm)	3L Recommended as foliar spray (single application) with the help of high volume knap sack sprayer (at green tip stage)	134
Cotton	To restrict vegetative growth, prevent shedding of squares/bolls & enhance yield	34.50g a.i./ha	150ml/ha	500 L/ha	42
Groundnut	To enhance yield by restricting vegetative growth	28.75 g a.i./ha	125 ml/ha	500 L/ha	70
Paclobutrazol 40% SC					
Pigeon Pea	Plant growth regulator	30	75	500	48
Soybean	Plant growth regulator	30	75	500	56
Prohexadione-Ca 10% WG					

Apple	Two split applications: 1 st application: at 3-5 leaves/ shoot 2 nd application 4 weeks after 1 st application	125 150	50 gm per 100 liter 60 gm per 100 liter	2500 2500	94
Sodium Para –Nitrophenolate 0.3% SL					
Cotton	Flower bud initiated stage and fruit set stage	0.5%	5 ml	800	16
Tomato	Flowering and fruit stages	0.5%	4 ml	200	7
Triacntanol 0.05% EC					
Cotton	To increase the yield Three sprays at 45, 65 and 85 days after planting	0.125 gm	0.25ltr	400-500	-
Rice	Three sprays at 25, 45 and 65 days after transplanting	0.125 gm	0.25ltr	400-500	-
Chilli	Three sprays at 25, 45 and 65 days after planting	0.125 gm	0.25ltr	400-500	-
Tomato	Three sprays at 25, 45 and 65 days after planting	0.125 gm	0.25 ltr	400-500	-
Groundnut	Three sprays at 25, 45 and 65 days after planting	0.125 gm	0.25 ltr	400-500	-
Potato	Two sprays at 30 and 45 days after planting	0.250 gm	0.50 ltr	500-600	-
Triacntanol 0.05% w/w min. GR					
Cotton	To increase the yield Broadcast & mix the desired quantity of granules in soil 2-3 days before sowing.	12.5 gm	25 kg.	-	-
Rice	Broadcast & mix the desired quantity of granules in soil 2-3 days before transplanting.	12.5 gm	25 kg.	-	-
Chilli	Broadcast & mix the desired quantity of granules in soil 2-3 days before sowing.	12.5 gm	25 kg.	-	-
Tomato	Broadcast & mix the desired quantity of granules in soil 2-3 days before sowing.	12.5 gm	25 kg.	-	-
Groundnut	Broadcast & mix the desired quantity of granules in soil 2-3 days before sowing.	12.5 gm	25 kg.	-	-
Triacntanol 0.1% EW					
Cotton	To increase the yield	0.25 g	0.25 ltr.	400-500	-

	Three sprays at 45, 65 and 85 days after sowing				
Rice	Three sprays at 25, 45 and 65 days after transplanting	0.25 g	0.25 ltr.	400-500	-
Chilli	Three sprays at 25, 45 and 65 days after transplanting	0.25 g	0.25 ltr.	400-500	-
Tomato	Three sprays at 25, 45 and 65 days after transplanting	0.25 g	0.25 ltr.	400-500	-
Groundnut	Three sprays at 25, 45 and 65 days after sowing	0.25 g	0.25 ltr.	400-500	-
Tea	Three sprays: 1 st spray on mature plants, 2 nd spray one month after 1 st spray, 3 rd spray one month after 2 nd spray	0.25 g	0.25 ltr	400-500	-
Cyclanilide 2.10% w/w +Mepiquat Chloride 8.40% w/w SC					
Cotton	First spray should be applied at square formation stage or after 45-55 days of sowing. 2 nd and 3 rd spray should be applied at an interval of 15 days.	4.40 + 17.60 to 4.95 + 19.80 gm	200 - 225	500	21
Gibberellic acid 1.8% + 6-Benzyladenine 1.8% L					
Apple	To increase the yield through enhancement of fruit size and weight, to improve the shape and development of prominent calyx lobes (typiness). To increase lateral bud break and shoot growth (branching) & improving branch angle of nursery stock young apple trees	30-60 ppm	840-1680 ppm	1000	-
