

### HINTS & SOLUTIONS


- |          |          |          |          |          |
|----------|----------|----------|----------|----------|
| Q.1 (C)  | Q.2 (D)  | Q.3 (A)  | Q.4 (B)  | Q.5 (B)  |
| Q.6 (B)  | Q.7 (C)  | Q.8 (B)  | Q.9 (C)  | Q.10 (A) |
| Q.11 (A) | Q.12 (C) | Q.13 (A) | Q.14 (C) | Q.15 (B) |
| Q.16 (B) | Q.17 (A) | Q.18 (C) | Q.19 (D) | Q.20 (B) |
| Q.21 (B) | Q.22 (A) | Q.23 (A) | Q.24 (C) | Q.25 (B) |
| Q.26 (A) | Q.27 (D) | Q.28 (D) | Q.29 (B) | Q.30 (B) |
| Q.31 (C) | Q.32 (C) | Q.33 (C) |          |          |

Q.34 (A)

(A) Fractional distillation is used when boiling point difference is present in between gangue and ore.

- |                                    |                                       |            |            |
|------------------------------------|---------------------------------------|------------|------------|
| Q.35 (ACD)                         | Q.36 (ABCD)                           | Q.37 (ABD) | Q.38 (AB)  |
| Q.39 (ABC)                         | Q.40 (ABC)                            | Q.41 (BC)  | Q.42 (ABD) |
| Q.43 (AB)                          | Q.44 (BC)                             | Q.45 (AD)  | Q.46 (C)   |
| Q.47 (BCD)                         | Q.48 (BCD)                            | Q.49 (AD)  |            |
| Q.50 (A) P (B) R (C) Q (D) P,Q,S   | Q.51 (A) P, R ; (B) P ; (C) Q ; (D) S |            |            |
| Q.52 (A) P; (B) Q; (C) P,R (D) P,S |                                       |            |            |

Q.53 (4055)




$$w = 4$$

$$x = 0$$

$$y = 5 \text{ (linkage isomers)}$$

$$z = 5$$



Q.54 (0005)

Q.55 (0003)

Pb, Hg, Cu so Ans. is 3