


**WORK SHEET - 07**

- Q.1 (C)
LAH reduces esters to alcohols but do not affect C=C

- Q.2 (C)
Q.3 (D)



- Q.4 (B)
Esters which give stable carbocation reacts with A_{AL}^1 Mech.

- Q.5 (C)




- Q.6 (C)

- Q.7 (C)


- Q.8 (B)

- Q.9 (A) (C)
 NaBH_4 will not reduce ketone

Q.10 (B)(C)(D)







So, one mol per mol CHI_3 is formed.



$\text{NaOH} + \text{I}_2$ is a good oxidising agent but $\text{NH}_4\text{OH} + \text{I}_2$ is not

Q.11 (A) (C) (D)



Q.12 (A) (C)




Q.13 (C)

Q.14 (B)


Q.15 (D)

Q.16 (A) T (B) P (C) Q (D) T


Q.17 0006




β-Keto acid gives $\text{CO}_2 \uparrow$




β-Nitro acid gives $\text{CO}_2 \uparrow$




β-imino acid gives $\text{CO}_2 \uparrow$



β-γ unsaturated $\text{CO}_2 \uparrow$




1,6 dioic acid gives $\text{CO}_2 \uparrow$




1,2,dioic acid gives $\text{CO}_2 \uparrow$


Q.18 0005




(more acid than B.A.)




(SIR)
More acidic than BA




(-I effect)
more acidic




(+Hyperconjugation +I)
less acidic




Less acidic




Less acidic




Less acidic



More acidic



more acidic



Less acidic

Acidic strength order : $\text{H}-\text{C}(=\text{O})-\text{OH} > \text{Ph}-\text{C}(=\text{O})-\text{OH} > \text{CH}_3-\text{COOH}$

-I < +M
slightly

+ I

Q.19 4

Q.20 2