



**Sri Chaitanya**  
Educational Institutions

**Infinity**  
**Learn**



**JEE ADVANCED**



**Sri Chaitanya** IIT Academy.,India.

☆ A.P ☆ T.S ☆ KARNATAKA ☆ TAMILNADU ☆ MAHARASTRA ☆ DELHI ☆ RANCHI

*A right Choice for the Real Aspirant*

ICON Central Office - Madhapur - Hyderabad

Sec: **Sr.Super60\_STERLING\_BT**

**Paper -2(Adv-2021-P2-Model)**

**Date: 28-07-2024**

**Time: 02.00Pm to 05.00Pm**

**CTA-02**

**Max. Marks: 180**

**28-07-2024\_Sr.Super60\_STERLING\_BT\_Jee-Adv (2021-P2)\_CTA-02\_Syllabus**

**PHYSICS**

: Thermal physics, Heat Transfer, Ray Optics, Geometrical Optics,  
Experiments

**CHEMISTRY**

: Nomenclature, Isomerism, GOC, Alkanes, Alkene & Alkyne

**MATHEMATICS**

: Function and Inverse Trigonometric Function, LCD, AOD

**Name of the Student:** \_\_\_\_\_

**H.T. NO:**

|  |  |  |  |  |  |  |
|--|--|--|--|--|--|--|
|  |  |  |  |  |  |  |
|--|--|--|--|--|--|--|

**JEE-ADVANCE-2021-P2-Model**

Time: 3:00Hr's

**IMPORTANT INSTRUCTIONS**

Max Marks: 180

**PHYSICS:**

| Section                  | Question Type                                            | +Ve Marks | - Ve Marks | No.of Qs  | Total marks |
|--------------------------|----------------------------------------------------------|-----------|------------|-----------|-------------|
| Sec – I(Q.N : 1 – 6)     | Questions with Multiple Correct Choice with Partial mark | +4        | -2         | 6         | 24          |
| Sec – II(Q.N : 7 – 12)   | Paragraph Questions with Numerical Value Answer Type     | +2        | 0          | 6         | 12          |
| Sec – III(Q.N : 13 – 16) | Paragraph Questions with Single Answer Type              | +3        | -1         | 4         | 12          |
| Sec – IV(Q.N : 17 – 19)  | Questions with Non-negative Integer Value Type           | +4        | 0          | 3         | 12          |
| <b>Total</b>             |                                                          |           |            | <b>19</b> | <b>60</b>   |

**CHEMISTRY:**

| Section                  | Question Type                                            | +Ve Marks | - Ve Marks | No.of Qs  | Total marks |
|--------------------------|----------------------------------------------------------|-----------|------------|-----------|-------------|
| Sec – I(Q.N : 20 – 25)   | Questions with Multiple Correct Choice with Partial mark | +4        | -2         | 6         | 24          |
| Sec – II(Q.N : 26 – 31)  | Paragraph Questions with Numerical Value Answer Type     | +2        | 0          | 6         | 12          |
| Sec – III(Q.N : 32 – 35) | Paragraph Questions with Single Answer Type              | +3        | -1         | 4         | 12          |
| Sec – IV(Q.N : 36– 38)   | Questions with Non-negative Integer Value Type           | +4        | 0          | 3         | 12          |
| <b>Total</b>             |                                                          |           |            | <b>19</b> | <b>60</b>   |

**MATHEMATICS:**

| Section                  | Question Type                                            | +Ve Marks | - Ve Marks | No.of Qs  | Total marks |
|--------------------------|----------------------------------------------------------|-----------|------------|-----------|-------------|
| Sec – I(Q.N : 39 – 44)   | Questions with Multiple Correct Choice with Partial mark | +4        | -2         | 6         | 24          |
| Sec – II(Q.N : 45 – 50)  | Paragraph Questions with Numerical Value Answer Type     | +2        | 0          | 6         | 12          |
| Sec – III(Q.N : 51 – 54) | Paragraph Questions with Single Answer Type              | +3        | -1         | 4         | 12          |
| Sec – IV(Q.N : 55 – 57)  | Questions with Non-negative Integer Value Type           | +4        | 0          | 3         | 12          |
| <b>Total</b>             |                                                          |           |            | <b>19</b> | <b>60</b>   |

Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 2

**Sri Chaitanya**  
Educational Institutions**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1  
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023****JEE MAIN  
2023**SINGARAJU  
VENKAT KOUNDINYA  
HOD, Sr.Chaitanya  
Super60 Class  
**300**  
AIR**RANK  
1****JEE Advanced  
2023**VANILALA  
CHIDVILAS REDDY  
HOD, Sr.Chaitanya  
Super60 Class  
**341**  
AIR**RANK  
1****NEET  
2023**BORA VARUN  
CHAKRAVARTHI  
HOD, Sr.Chaitanya  
Super60 Class  
**720**  
AIR**RANK  
1**

## PHYSICS


Max. Marks: 60

## SECTION-1(Maximum Marks: 24)

One or More Type

- This section contains SIX (06) questions.
- Each question has FOUR options (A), (B), (C) and (D). ONE OR MORE THAN ONE of these four option(s)
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:  
 Full Marks : +4 If only (all) the correct option(s) is(are) chosen;  
 Partial Marks : +3 If all the four options are correct but ONLY three options are chosen;  
 Partial Marks : +2 If three or more options are correct but ONLY two options are chosen, both of which are correct;  
 Partial Marks : +1 If two or more options are correct but ONLY one option is chosen and it is a correct option;  
 Zero Marks : 0 If unanswered;  
 Negative Marks : -2 In all other cases.

01. Two bodies A and B have thermal emissivities of 0.01 and 0.81 respectively. The outer surface areas of the two bodies are the same. The two bodies radiate energy at the same rate. The wavelength  $\lambda_B$ , corresponding to the maximum spectral radiance in the radiation from B, is shifted from the wavelength corresponding to the maximum spectral radiance in the radiation from A by 1.00 mm. If the temperature of A is 5802 K, then
- A) the temperature of B is 1934 K      B)  $\lambda_B = 1.5 \text{ mm}$   
 C) the temperature of B is 11604 K      D) the temperature of B is 2901 K
02. The temperature Vs time graph for two materials A & B has been plotted when the energy has been given to both at same rate. Initially both the materials are in solid state (below their Melting point) and have equal mass. Given  $L_f$  = latent heat of fusion and  $L_v$  = latent heat of vapourisation. Then



- A)  $L_{fA} = L_{fB}$   
 B)  $L_{vA} < L_{vB}$   
 C) Specific heat of A in liquid state > specific heat of B in liquid state  
 D) Specific heat of A in solid state > specific heat of B in solid state

Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 3

**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1**  
**IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN 2023**  
 SINGARAJU  
 VENKAT KOUNDINYA  
 Rank 1  
 300  
 300



**RANK 1**

**JEE Advanced 2023**  
 VANILALA  
 CHIDVILAS REDDY  
 Rank 1  
 341  
 350



**RANK 1**

**NEET 2023**  
 BORA VARUN  
 CHAKRAVARTHI  
 Rank 1  
 720  
 720




**RANK 1**






03. Four rods A, B, C, D of same length and material but of different radii  $r, r\sqrt{2}, r\sqrt{3}$  and  $2r$  respectively are held between two rigid walls. The temperature of all rods is increased by same amount. If the rods do not bend, then
- the stress in the rods are in the ratio 1 : 2 : 3 : 4
  - the force on the rod exerted by the wall are in the ratio 1 : 2 : 3 : 4
  - the energy stored in the rods due to elasticity are in the ratio 1 : 2 : 3 : 4
  - the strains produced in the rods are in the ratio 1 : 2 : 3 : 4
04. 1 mole of an ideal monoatomic gas is confined in a cylinder fitted with a piston as shown in figure. Take mass of the piston is 0.5 kg, spring constant,  $K = 1000 \text{ N/m}$ , atmospheric pressure  $P_0 = 10^5 \text{ N/m}^2$ , cross sectional area of piston  $A = 50 \text{ cm}^2$ . If energy of 50J has been supplied to system then piston moves up by 5 cm (take  $g = 10 \text{ m/s}^2$ ). Initially spring is in its natural position. Mark the correct statement(s)



- Increase in internal energy of the system is 23.50 J
  - Initial pressure of the gas is  $1.01 \times 10^5 \text{ N/m}^2$
  - For mentioned change, the temperature of gas increases by 1.88K
  - For mentioned change, the temperature of gas decreases by 1.88K
05. A transparent thin film of uniform thickness and refractive index  $n_1 = 1.4$  is coated on the convex spherical surface of radius R at one end of a long solid glass cylinder of refractive index  $n_2 = 1.5$ , as shown in the figure. Rays of light parallel to the axis of the cylinder traversing through the film from air to glass get focused at distance  $f_1$  from the film, while rays of light traversing from glass to air get focused at distance  $f_2$  from the film, Then




- $|f_1| = 3R$
- $|f_1| = 2.8R$
- $|f_2| = 2R$
- $|f_2| = 1.4R$





06. One mole of a monoatomic ideal gas goes through a thermodynamic cycle, as shown in the volume versus temperature (V-T) diagram. The correct statement(s) is/are:  
[R is the gas constant]



- A) Work done in this thermodynamic cycle ( $1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 1$ ) is  $|W| = \frac{1}{2} RT_0$   
 B) The above thermodynamic cycle exhibits only isochoric and adiabatic processes.  
 C) The ratio of heat transfer during processes  $1 \rightarrow 2$  and  $2 \rightarrow 3$  is  $\left| \frac{Q_{1 \rightarrow 2}}{Q_{2 \rightarrow 3}} \right| = \frac{5}{3}$   
 D) The ratio of heat transfer during processes  $1 \rightarrow 2$  and  $3 \rightarrow 4$  is  $\left| \frac{Q_{1 \rightarrow 2}}{Q_{3 \rightarrow 4}} \right| = \frac{1}{2}$

## SECTION-2(Maximum Marks: 12)

### Paragraph with Numerical

- This section contains THREE (03) question stems.
- There are TWO (02) questions corresponding to each question stem.
- The answer to each question is a NUMERICAL VALUE.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, truncate/round-off the value to TWO decimal places.
- Answer to each question will be evaluated according to the following marking scheme:  
 Full Marks : +2 If ONLY the correct numerical value is entered at the designated place;  
 Zero Marks : 0 In all other cases.

### Question Stem for Question Nos. 7 and 8

#### Question Stem

A cubical box of side 1m contains nitrogen gas (molecular weight 28g) at a pressure of  $100 \text{ N/m}^2$ . During an observation time of one second, a molecule travelling with the root-mean-square speed parallel to one of the edges of the cube, was found to make 500 hits per second with a particular wall, without any collision with other molecules.

Take  $R = \frac{25}{3} \text{ J/mol-K}$  and  $k = 1.38 \times 10^{-23} \text{ J/K}$

7. The temperature of the gas is \_\_\_\_\_ (in kelvin)  
 8. The average translational kinetic energy per molecule of gas is \_\_\_\_\_ ( $\times 10^{-22} \text{ J}$ ).  
 Round off to nearest integer



JEE MAIN

2023  
SINGARAJU  
VENKAT KOUNDINYA  
Rank 1  
Sri Chaitanya  
JEE 12th Class  
300  
300  
MAINS



RANK

1

JEE Advanced

2023  
VAVILALA  
CHIDVILAS REDDY  
Rank 1  
Sri Chaitanya  
JEE 12th Class  
341  
350  
MAINS



RANK

1

NEET

2023  
BORA VARUN  
CHAKRAVARTHI  
Rank 1  
Sri Chaitanya  
JEE 12th Class  
720  
720  
MAINS



RANK


1



### Question Stem for Question Nos. 9 and 10

#### Question Stem

Five rods of same material and same length are joined as shown. Cross-section of rods ab, ad and bc are A, 2A and 3A respectively. Ends a and c are maintained at temperatures  $200^{\circ}\text{C}$  and  $0^{\circ}\text{C}$  respectively.




9. For cross-section  $(x)A$  of rod dc there will be no heat flow through rod bd, then  $x = \underline{\hspace{2cm}}$
10. If  $x$  has the value calculated in above question. Then temperature of junction b is:  $\underline{\hspace{2cm}}^{\circ}\text{C}$

### Question Stem for Question Nos. 11 and 12

#### Question Stem

In the arrangement shown, the angle of thin prism is  $1^{\circ}$ . The image of the object kept at origin O is formed at a point  $\left(10\text{ cm}, -\frac{\pi}{24}\text{ cm}\right)$  for paraxial rays. Use the fact that small angled prism produces only lateral shift for paraxial rays



11. Radius of curvature of the mirror is  $\underline{\hspace{2cm}}$  (in cm)
12. Refractive index of the prism is  $\underline{\hspace{2cm}}$

### SECTION-3(Maximum Marks: 12) Paragraph with Single Answer Type

- This section contains TWO (02) paragraphs. Based on each paragraph, there are TWO (02) questions.
- Each question has FOUR options (A), (B), (C) and (D). ONLY ONE of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer
- Answer to each question will be evaluated according to the following marking scheme:

Full Marks : +3 If ONLY the correct option is chosen;

Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks : -1 In all other cases.

Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 6



**Sri Chaitanya**  
Educational Institutions



**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1  
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN  
2023**  
SINGARAJU  
VENKAT KOUNDINYA  
Rank 1  
300  
300  
MAHAR



**RANK  
1**

**JEE Advanced  
2023**  
VANILALA  
CHIDVILAS REDDY  
Rank 1  
341  
350  
MAHAR



**RANK  
1**

**NEET  
2023**  
BORA VARUN  
CHAKRAVARTHI  
Rank 1  
720  
720  
MAHAR



**RANK  
1**



## Paragraph-I


Most materials have the refractive index,  $n > 1$ . So, when a light ray from air enters a naturally occurring material, then by Snell's law,  $\frac{\sin \theta_1}{\sin \theta_2} = \frac{n_2}{n_1}$ , it is understood that the

refracted ray bends towards the normal. But it never emerges on the same side of the normal as the incident ray. According to electromagnetism, the refractive index of the medium is given by the relation,  $n = c/v = \pm \sqrt{\epsilon_r \mu_r}$ , where  $c$  is the speed of electromagnetic waves in vacuum,  $v$  its speed in the medium,  $\epsilon_r$  and  $\mu_r$  are the relative permittivity and permeability of the medium respectively.

In normal materials, both  $\epsilon_r$  and  $\mu_r$ , are positive, implying positive  $n$  for the medium.

When both  $\epsilon_r$  and  $\mu_r$  are negative, one must choose the negative root of  $n$ . Such negative refractive index materials can now be artificially prepared and are called meta-materials. They exhibit significantly different optical behavior, without violating any physical laws. Since  $n$  is negative, it results in a change in the direction of propagation of the refracted light. However, similar to normal materials, the frequency of light remains unchanged upon refraction even in meta-materials.

13. For light incident from air on a meta-material, the appropriate ray diagram is






14. Choose the correct statement.
- A) The speed of light in the meta-material is  $v = c|n|$
- B) The speed of light in the meta-material is  $v = \frac{c}{|n|}$
- C) The speed of light in the meta-material is  $v = c$ .
- D) The wavelength of the light in the meta-material ( $\lambda_m$ ) is given by  $\lambda_m = \lambda_{air}|n|$ , where  $\lambda_{air}$  is wavelength of the light in air.

## Paragraph-II

In the figure, a container is shown to have a movable (without friction) piston on top. The container and the piston are all made of perfectly insulated material allowing no heat transfer between outside and inside the container. The container is divided into two compartments by a rigid partition made of a thermally conducting material that allows slow transfer of heat. The lower compartment of the container is filled with 2 moles of an ideal monoatomic gas at 700K and the upper compartment is filled with 2 moles of an ideal diatomic gas at 400K. The heat capacities per mole of an ideal monoatomic gas are  $C_V = \frac{3}{2}R$ ,  $C_P = \frac{5}{2}R$ , and those for an ideal diatomic gas are  $C_V = \frac{5}{2}R$ ,  $C_P = \frac{7}{2}R$ .



15. Consider the partition to be rigidly fixed so that it does not move. When equilibrium is achieved, the final temperature of the gases will be
- A) 550 K      B) 525 K      C) 513 K      D) 490 K
16. Now consider the partition to be free to move without friction so that the pressure of gases in both compartments is the same. The total work done by the gases till the time they achieve equilibrium will be
- A) 250 R      B) 200 R      C) 100 R      D) - 100 R

JEE MAIN  
2023

SINGARAJU  
VENKAT KOUNDINYA  
sri chaitanya  
Super 60 Class  
300  
300  
AIR

RANK  
1JEE Advanced  
2023

VAIVILALA  
CHIDVILAS REDDY  
sri chaitanya  
Super 60 Class  
341  
350  
AIR

RANK  
1NEET  
2023

BORA VARUN  
CHAKRAVARTHI  
sri chaitanya  
Super 60 Class  
720  
720  
AIR

RANK  
1






### SECTION-4(Maximum Marks: 12) Non-Negative Integer Answer Type

- This section contains THREE (03) questions.
- The answer to each question is a NON-NEGATIVE INTEGER.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:

Full Marks : +4 If ONLY the correct integer is entered;

Zero Marks : 0 In all other cases.

17. A diatomic ideal gas is compressed adiabatically to  $\frac{1}{32}$  of its initial volume. If the initial temperature of the gas is  $T_i$  (in Kelvin) and the final temperature is  $aT_i$ , the value of  $a$  is \_\_\_\_\_
18. Two spherical stars A and B emit blackbody radiation. The radius of A is 300 times that of B and A emits  $10^4$  times the power emitted from B. The ratio  $\left(\frac{\lambda_A}{\lambda_B}\right)$  of their wavelengths  $\lambda_A$  and  $\lambda_B$  at which the peaks occur in their respective radiation curves is  $\sqrt{N}$ . Then  $N =$  \_\_\_\_\_
19. The apparatus shown in the figure consists of four glass columns connected by horizontal sections. The height of two central columns B and C are 49 cm each. The two outer columns A and D are open to the atmosphere. A and C are maintained at a temperature of  $95^\circ\text{C}$  while the columns B and D are maintained at  $5^\circ\text{C}$ . The height of the liquid in A and D measured from the base are 52.8 cm and 51 cm respectively. The coefficient of thermal volume expansion of the liquid is  $N \times 10^{-4} (^\circ\text{C})^{-1}$ , then  $N =$  -----



Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 9



**Sri Chaitanya**  
Educational Institutions



**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1  
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN  
2023**

SINGARAJU  
VENKAT KOUNDINYA  
AIR 1  
Sri Chaitanya  
JEE Prep Class  
**300**  
MAINS



**RANK  
1**

**JEE Advanced  
2023**

VAHILA  
CHIDVILAS REDDY  
AIR 1  
Sri Chaitanya  
JEE Prep Class  
**341**  
MAINS  
**350**  
EXAM



**RANK  
1**

**NEET  
2023**

BORA VARUN  
CHAKRAVARTHI  
AIR 1  
Sri Chaitanya  
JEE Prep Class  
**720**  
MAINS  
**720**  
EXAM



**RANK  
1**



## CHEMISTRY

Max. Marks: 60




## SECTION-1(Maximum Marks: 24)

One or More Type

- This section contains SIX (06) questions.
- Each question has FOUR options (A), (B), (C) and (D). ONE OR MORE THAN ONE of these four option(s)
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:  
 Full Marks : +4 If only (all) the correct option(s) is(are) chosen;  
 Partial Marks : +3 If all the four options are correct but ONLY three options are chosen;  
 Partial Marks : +2 If three or more options are correct but ONLY two options are chosen, both of which are correct;  
 Partial Marks : +1 If two or more options are correct but ONLY one option is chosen and it is a correct option;  
 Zero Marks : 0 If unanswered;  
 Negative Marks : -2 In all other cases.



(C is C-14 isotope)






D) HBr

21. Which of the following is/are correct order of basic strength.

A)  $\text{NH}_3 > \text{pyridine}$

B) Imidazole > pyridine




A) F

B) Cl

C) Br

D) I



What is/are the suitable match of x.

A)  $\text{Cr}_2\text{O}_3$

B)  $\text{V}_2\text{O}_5$

C)  $\text{KMnO}_4$

D)  $\text{Mo}_2\text{O}_3$

Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 10



**Sri Chaitanya**  
Educational Institutions

**Infinity**  
Learn



**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1**  
**IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

JEE MAIN

2023

SINGARAJU  
VENKAT KOUNDINYA

Rank 1  
Sri Chaitanya  
Super Class

300  
300  
MAINS

RANK

1

JEE Advanced

2023

VARILALA  
CHIDVILAS REDDY

Rank 1  
Sri Chaitanya  
Super Class

341  
350  
MAINS

RANK

1

NEET

2023

BORA VARUN  
CHAKRAVARTHI

Rank 1  
Sri Chaitanya  
Super Class





720  
720  
MAINS

RANK




1



24. Which of the following is the correct combination of reactant, catalyst/reagent and product  
(Given reactions are not the balanced equations)



25. Which of the following is/are the correct match of structure and its property or nature.









**SECTION-2(Maximum Marks: 12)****Paragraph with Numerical**

- This section contains THREE (03) question stems.
- There are TWO (02) questions corresponding to each question stem.
- The answer to each question is a NUMERICAL VALUE.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, truncate/round-off the value to TWO decimal places.
- Answer to each question will be evaluated according to the following marking scheme:  
Full Marks : +2 If ONLY the correct numerical value is entered at the designated place;  
Zero Marks : 0 In all other cases.

**Question Stem for Question Nos. 26 and 27****Question Stem**


Carbonyl compounds which contain  $\alpha - H$  participate in Tautomerism. Tautomers exist in dynamic equilibrium. The stability of enol form depends on H-bond, steric factors, electronic factors, resonance etc.

|    | Column – I<br>Compound                                                              |    | Column – II<br>% Enol content |
|----|-------------------------------------------------------------------------------------|----|-------------------------------|
| A. |    | P. | 89.2                          |
| B. |   | Q. | $1.1 \times 10^{-6}$          |
| C. |  | R. | 80                            |
| D. |  | S. | 8.4                           |


The order of enol content in column II is not in match of order of column-I

26. What is the % enol content value corresponds to compound C is \_\_\_\_\_
27. What is the % enol content value corresponds to compound D is \_\_\_\_\_

**JEE MAIN**  
**2023**SINGARAJU  
VENKAT KOUNDINYA  
HOD, Chemistry  
Sri Chaitanya  
JEE 12th Class**300**  
**300**  
RANK**RANK****1****JEE Advanced**  
**2023**VANILALA  
CHIDVILAS REDDY  
HOD, Chemistry  
Sri Chaitanya  
JEE 12th Class**341**  
**350**  
RANK**RANK****1****NEET**  
**2023**BORA VARUN  
CHAKRAVARTHI  
HOD, Chemistry  
Sri Chaitanya  
JEE 12th Class**720**  
**720**  
RANK**RANK****1**

**Question Stem for Question Nos. 28 and 29****Question Stem**


The IUPAC name of x is cyclohexa-p,q-diene



Y can be detected by Tollens reagent

28. What is the value of "q - p"

29. Number of hybrid orbitals involved in the structure of y is \_\_\_\_ (y contain elements C & H only)

**Question Stem for Question Nos. 30 and 31****Question Stem**

30. What is the position of Br in the IUPAC name (IUPAC numbering) of the compound x

31. What is the position of Cl in the IUPAC name (IUPAC numbering) of the compound y

**SECTION-3(Maximum Marks: 12)****Paragraph with Single Answer Type**

- This section contains TWO (02) paragraphs. Based on each paragraph, there are TWO (02) questions.
- Each question has FOUR options (A), (B), (C) and (D). ONLY ONE of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer
- Answer to each question will be evaluated according to the following marking scheme:

Full Marks : +3 If ONLY the correct option is chosen;


Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks : -1 In all other cases.

**Paragraph-I:**

Resonance stabilise the molecule as the energy of the resonance hydride is less than the energy of any single cannoncial structure. Resonance average the bond characteristics as whole. the molecule as such has a single structure which is the resonance hydride of the canonical forms.

32. Which of the following pair exist in real equilibrium?



Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 13



**Sri Chaitanya**  
Educational Institutions

**Infinity**  
Learn



**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1**  
**IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN**

**2023**

SINGARAJU  
VENKAT KOUNDINYA

Rank 1

300

300

MAHAR

**RANK**

**1**

**JEE Advanced**

**2023**

VARILALA  
CHIDVILAS REDDY

Rank 1

341

350

MAHAR

**RANK**

**1**

**NEET**

**2023**

BORA VARUN  
CHAKRAVARTHI

Rank 1

720

720


MAHAR

**RANK**

**1**



33. How many of the following structures have no real existence for cyclopropenyl cation (Aromatic)




- A) 1 B) 2 C) 3 D) 4

### Paragraph-II:

The optical isomers are constitutionally identical but differ in the direction of rotation of plane polarised light. Meso isomer is optically inactive due to internal compensation. The extent of rotation and nature of rotation of plane polarised light depends on many factors.

34. The optical rotation of A is  $+45^{\circ}$ . What is the optical rotation of B at same set of conditions.




A is

B is

- A)  $+45^{\circ}$  B)  $-45^{\circ}$  C)  $+90^{\circ}$  D) 0

35. Which of the following is optically inactive optical isomer.






### SECTION-4(Maximum Marks: 12)

#### Non-Negative Integer Answer Type


- This section contains THREE (03) questions.
- The answer to each question is a NON-NEGATIVE INTEGER.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:

Full Marks : +4 If ONLY the correct integer is entered;

Zero Marks : 0 In all other cases.





X is the most stable enol form of given structure. Number of  $\pi$  – bonds under conjugation with  $-OH$  group of stable enol group



What is the degree of unsaturation of compound x.

38. Number of resonance structures possible for A is x and for B is y. The value of " $x - y$ " is \_\_\_\_\_.

JEE MAIN  
2023

SINGARAJU  
VENKAT KOUNDINYA  
AIR 1st Rank  
Sri Chaitanya  
Super 60 Class  
300  
300  
MAINS

RANK  
1JEE Advanced  
2023

VAIVILALA  
CHIDVILAS REDDY  
AIR 1st Rank  
Sri Chaitanya  
Super 60 Class  
341  
350  
MAINS

RANK  
1NEET  
2023

BORA VARUN  
CHAKRAVARTHI  
AIR 1st Rank  
Sri Chaitanya  
Super 60 Class  
720  
720  
MAINS

RANK  
1



**MATHEMATICS****Max. Marks: 60****SECTION-1(Maximum Marks: 24)****One or More Type**

- This section contains SIX (06) questions.
- Each question has FOUR options (A), (B), (C) and (D). ONE OR MORE THAN ONE of these four option(s)
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:  
 Full Marks : +4 If only (all) the correct option(s) is(are) chosen;  
 Partial Marks : +3 If all the four options are correct but ONLY three options are chosen;  
 Partial Marks : +2 If three or more options are correct but ONLY two options are chosen, both of which are correct;  
 Partial Marks : +1 If two or more options are correct but ONLY one option is chosen and it is a correct option;  
 Zero Marks : 0 If unanswered;  
 Negative Marks : -2 In all other cases.

39. 
$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{\int_{\frac{\pi}{2}}^x \sin\left(2t^{\frac{1}{3}}\right) + \cos\left(t^{\frac{1}{3}}\right) dt}{\left(x - \frac{\pi}{2}\right)^2} = \frac{k_1 \pi^2}{k_2}$$
 where  $k_1, k_2$  are co-prime integers then

A)  $k_1 + k_2 = 17$

B)  $k_1 + k_2 = 21$

C)  $k_1 - k_2 = 1$

D) Sum of even factors of  $k_1 \cdot k_2 = 182$

40. Which of the following is (are) TRUE?

A) If the function  $f(x) = \frac{\sin 3x + \alpha \sin x - \beta \cos 3x}{x^3}$ ,  $x \in \mathbb{R} - \{0\}$  is continuous at  $x = 0$  then  $f(0)$  is equal to '-4'

B) Let  $f(x) = x^5 + 2x^3 + 3x + 1$ ,  $x \in \mathbb{R}$  and  $g(x)$  be a function such that

$g(f(x)) = x$  for all  $x \in \mathbb{R}$ , then  $\frac{g'(7)}{g(7)} = \frac{1}{14}$  (Here  $g'(x)$  means first derivate of  $g(x)$  w.r.t  $x$ )

C) Let  $f: [-1, 2] \rightarrow \mathbb{R}$  be given by  $f(x) = 2x^2 + x + \left[x^2\right] - [x]$ , where  $[t]$  is greatest integer function  $\leq t$ . The no. Of points where  $f(x)$  is not continuous is '4'

D) Let  $a > 0$ , be a root of the equation  $2x^2 + x - 2 = 0$ . If

$$\lim_{x \rightarrow \frac{1}{a}} \frac{16(1 - \cos(2 + x - 2x^2))}{(1 - ax)^2} = \alpha + \beta\sqrt{17} \text{ where } \alpha, \beta \in \mathbb{Z}, \text{ then } \alpha - \beta = 136$$

Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 16

**Sri Chaitanya**  
Educational Institutions**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1  
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023****JEE MAIN  
2023**SINGARAJU  
VENKAT KOUNDINYA  
Rank 1  
Sri Chaitanya  
JEE 12th Class  
**300**  
AIR 1**RANK****1****JEE Advanced  
2023**VANILALA  
CHIDVILAS REDDY  
Rank 1  
Sri Chaitanya  
JEE 12th Class  
**341**  
AIR 1**RANK****1****NEET  
2023**BORA VARUN  
CHAKRAVARTHI  
Rank 1  
Sri Chaitanya  
JEE 12th Class  
**720**  
AIR 1**RANK****1**



41. If the function  $f : (-\infty, -1] \rightarrow (a, b]$  is defined by  $f(x) = e^{x^3 - 3x + 1}$  is one to one and onto. Then which of the following is/are ?
- A)  $a = 0$
- B)  $e^3 = b$
- C) The perpendicular distance of the point  $p(2b + 6, a + 2)$  from the line  $x + e^{-3}y = 6$  is  $2\sqrt{1 + e^6}$
- D)  $\lim_{x \rightarrow 0} \frac{e - (1 + 2x)^{(1/2x)}}{x} = b.e^{-2}$
42. Let  $x = \frac{m}{n}$  ( $m, n$  are co-prime natural numbers) be a solution of the equation  $\cos(2\sin^{-1}x) = \frac{1}{9}$  and  $\alpha, \beta (\alpha > \beta)$  be the roots of the equation  $mx^2 - nx - m + n = 0$ , then which of the following is are CORRECT?
- A)  $m + n = 5$
- B)  $\alpha = 1$  and  $\beta = \frac{1}{2}$
- C) The point  $(\alpha, \beta)$  lies on the line  $7x + 8y = 9$
- D) The range of  $f(\theta) = \frac{\sin^4 \theta + 3\cos^2 \theta}{\sin^4 \theta + \cos^2 \theta} (\theta \in \mathbb{R})$  is  $[\alpha, 6\beta]$
43. Let  $g(x) = f(x) + f(1-x)$  and  $f''(x) > 0, \forall x \in (0, 1)$ . If  $g(x)$  is decreasing in  $(0, \alpha)$  and increasing in  $(\alpha, 1)$  then (w.r.t value of ' $\alpha$ ' as defined in this question) which of the following is/are correct
- A)  $\alpha = \frac{1}{2}$
- B) The value of  $\tan^{-1}(2\alpha) + \tan^{-1}\left(\frac{1}{\alpha}\right) + \tan^{-1}\left(\frac{\alpha+1}{\alpha}\right) = \pi$
- C) If  $h(x) = \frac{(\tan 2^\circ)x + \log_e(123)}{x \log_e(2025) - \tan 2^\circ} (x > 0)$  then the least value of  $h(h(x)) + h\left(h\left(\frac{4}{x}\right)\right)$  is ' $8\alpha$ '
- D) The number of points, where the curve  $y = x^5 - 20x^3 + 50x + 2$  crosses x axis is ' $10\alpha$ '



JEE MAIN 2023

SINGARAJU  
VENKAT KOUNDINYA  
AIR 1  
Sri Chaitanya  
Super 60 Class  
300  
300  
MAINS

RANK

1

JEE Advanced 2023

VANILALA  
CHIDVILAS REDDY  
AIR 1  
Sri Chaitanya  
Super 60 Class  
341  
350  
MAINS

RANK

1

NEET 2023

BORA VARUN  
CHAKRAVARTHI  
AIR 1  
Sri Chaitanya  
Super 60 Class  
720  
720  
MAINS

RANK

1



44. Let  $f(x)$  be a polynomial degree of '5' such that  $\lim_{x \rightarrow 0} \frac{f(x)}{x^2} = 1, f(-1) = -1, f(x) - 14x$  has an extremum at  $x = 1$  and  $f(x) - 10x$  has extremum at  $x = -1$  then
- A)  $f(2) = 92$
- B) The equation of tangent at  $(-1, -1)$  on the curve  $y = f(x)$  is  $x + 10y + 11 = 0$
- C) The slope of normal at  $(2, 92)$  on the curve  $y = f(x)$  is  $\frac{-1}{232}$
- D)  $y = f(x)$  has local minimum at  $x = 0$  is '0'.

**SECTION-2(Maximum Marks: 12)****Paragraph with Numerical**

- This section contains THREE (03) question stems.
- There are TWO (02) questions corresponding to each question stem.
- The answer to each question is a NUMERICAL VALUE.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, truncate/round-off the value to TWO decimal places.
- Answer to each question will be evaluated according to the following marking scheme:  
Full Marks : +2 If ONLY the correct numerical value is entered at the designated place;  
Zero Marks : 0 In all other cases.

**Question Stem for Question Nos. 45 and 46****Question Stem**

$$\text{Let } y(\theta) = \frac{2\cos\theta + \cos 2\theta}{\cos 3\theta + 4\cos 2\theta + 5\cos\theta + 2},$$

$$f(x) = \begin{cases} x^3 \sin\left(\frac{1}{x}\right) & \text{if } (x \neq 0) \\ 0 & \text{if } x = 0 \end{cases}$$

45. At  $\theta = \frac{\pi}{2}$ ,  $y'' + y' + y$  is equal to (Where  $y''$  = second derivative of 'y' w.r.t to  $\theta$   
 $y'$  = derivative of 'y' w.r.t to ' $\theta$ ') \_\_\_\_\_
46.  $f''\left(\frac{2}{\pi}\right) = \frac{K_1 + K_2\pi^2}{K_3\pi}$  then  $\frac{K_1}{K_2 + K_3}$  equal to \_\_\_\_\_ ( $K_1, K_2, K_3 \in \mathbb{Z}$  &  $K_3 \neq 0$ )

**JEE MAIN**  
**2023**SINGARAJU  
VENKAT KOUNDINYA  
Rank 1  
Sri Chaitanya  
Super 60 Class  
**300**  
AIR 1**RANK****1****JEE Advanced**  
**2023**VANILALA  
CHIDVILAS REDDY  
Rank 1  
Sri Chaitanya  
Super 60 Class  
**341**  
AIR 1**RANK****1****NEET**  
**2023**BORA VARUN  
CHAKRAVARTHI  
Rank 1  
Sri Chaitanya  
Super 60 Class  
**720**  
AIR 1**RANK****1**

**Question Stem for Question Nos. 47 and 48****Question Stem**

Let  $S \in (-1, \infty)$  and  $f: S \rightarrow \mathbb{R}$  be defined as

$$f(x) = \int_{-1}^x (e^t - 1)^{11} \cdot (2t - 1)^5 \cdot (t - 2)^7 \cdot (t - 3)^{12} \cdot (2t - 10)^{61} dt, \text{ Let } p = \text{sum of squares of the}$$

values of  $x$ , where  $f(x)$  attains local maxima on  $S$ , and  $q = \text{sum of the values of } x$ , where

$f(x)$  attains local minima on  $S$ ,

Let  $h(x) = \left( \sqrt{8x - x^2 - 12} - 4 \right)^2 + (x - 7)^2$  ( $x \in \mathbb{R}$ ). Let the maxima and minimum values of  $h(x)$  be  $M$  and ' $m$ ' respectively

47.  $M^2 + m^2 = \underline{\hspace{2cm}}$

48.  $p^2 + 4q = \underline{\hspace{2cm}}$

**Question Stem for Question Nos. 49 and 50****Question Stem**

Let  $[t]$  is greatest integer function  $\leq t$ , Let  $A$  be set of all prime factors of 2310 and

$$f: A \rightarrow \mathbb{Z} \text{ be the function defined by } f(x) = \left\lfloor \log_2 \left( x^2 + \left\lceil \frac{x^3}{5} \right\rceil \right) \right\rfloor$$

49. Sum of elements in range of  $f(x)$  is  $\underline{\hspace{2cm}}$

50. The number of one-one functions from  $A$  to the range of ' $f$ ' is  $\underline{\hspace{2cm}}$

**SECTION-3(Maximum Marks: 12)**  
**Paragraph with Single Answer Type**

- This section contains TWO (02) paragraphs. Based on each paragraph, there are TWO (02) questions.
- Each question has FOUR options (A), (B), (C) and (D). ONLY ONE of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer
- Answer to each question will be evaluated according to the following marking scheme:

Full Marks : +3 If ONLY the correct option is chosen;

Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks : -1 In all other cases.

Sec: Sr.Super60\_STERLING\_BT

Space for rough work

Page 19



**Sri Chaitanya**  
Educational Institutions



**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1**  
**IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN**  
**2023**

SINGARAJU  
VENKAT KOUNDINYA  
Rank 1  
Sri Chaitanya  
Super 60 Class  
**300**  
AIR 1



**RANK**

**1**

**JEE Advanced**  
**2023**

VAHILALA  
CHIDVILAS REDDY  
Rank 1  
Sri Chaitanya  
Super 60 Class  
**341**  
AIR 1



**RANK**

**1**

**NEET**  
**2023**

BORA VARUN  
CHAKRAVARTHI  
Rank 1  
Sri Chaitanya  
Super 60 Class  
**720**  
AIR 1



**RANK**

**1**



**Paragraph-I:**

Let  $f(x) = \left[ \sin^{-1} x \right] + \left[ \cos^{-1} x \right], x \in [0, 1]$  where  $[t]$  denotes the greatest integer  $\leq t$  and

$\alpha$  = Number of discontinuities of  $f(x)$

$\beta$  = Number of points at which  $f(x)$  is not differentiable

$\gamma$  = The least value of  $f(x)$

$\delta$  = The greatest value of  $f(x)$  then

51.  $\frac{\alpha + \beta}{\delta - \gamma} =$

A) 2

B) 4

C) 1

D) 0

52. The solution set of the equation  $f(x) = 0$  is an interval of length

A)  $\sqrt{2} \cos\left(1 + \frac{\pi}{4}\right)$  B)  $\sqrt{2} \sin\left(1 + \frac{\pi}{4}\right)$  C)  $\sqrt{2} \sin\left(1 - \frac{\pi}{4}\right)$  D)  $\sqrt{2} \cos\left(1 - \frac{\pi}{4}\right)$

**Paragraph-II:**

Let  $\alpha = \sum_{k=1}^{\infty} \cos^{2k}\left(\frac{\pi}{3}\right)$ , Let  $g: [0, 1] \rightarrow \mathbb{R}$  be the function defined by  $g(x) = 2^{\alpha x} + 2^{\alpha(1-x)}$

and  $h(x) = 3^{\left(x^2 - 2\right)^3 + 4}, (x \in \mathbb{R})$

53. Which of the following is FALSE about  $g(x)$

A) The minimum of  $g(x)$  is  $2^{\frac{7}{6}}$

B) The maximum of  $g(x)$  is  $1 + 2^{\frac{1}{3}}$

C) The function  $g(x)$  attains its maximum at more than one point

D) The function  $g(x)$  attains its minimum at more than one point

54. Which of the following is FALSE about  $h(x)$

A)  $x = 0$  is a point of local minimum of  $h(x)$

B)  $x = \sqrt{2}$  is a point of inflection of  $h(x)$

C)  $h(x)$  is increasing for  $x > \sqrt{2}$

D)  $h(x)$  is decreasing for  $x > \sqrt{2}$



JEE MAIN  
2023

SINGARAJU  
VENKAT KOUNDINYA  
All India Rank  
Sri Chaitanya  
Super 60 Class  
300  
300  
MAINS



RANK  
1

JEE Advanced  
2023

VARILALA  
CHIDVILAS REDDY  
All India Rank  
Sri Chaitanya  
Super 60 Class  
341  
350  
MAINS



RANK  
1

NEET  
2023

BORA VARUN  
CHAKRAVARTHI  
All India Rank  
Sri Chaitanya  
Super 60 Class  
720  
720  
MAINS



RANK  
1



### SECTION-4(Maximum Marks: 12)

#### Non-Negative Integer Answer Type

- This section contains THREE (03) questions.
- The answer to each question is a NON-NEGATIVE INTEGER.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:

Full Marks : +4 If ONLY the correct integer is entered;

Zero Marks : 0 In all other cases.

55. The domain of the function  $f(x) = \cos^{-1}\left(\frac{x^2 - 3x + 2}{x^2 + 2x - 1}\right)$ , ( $x$  is real) is  $[k_1, \infty)$  and

$\lim_{x \rightarrow 0} (1 + 3x)^{\frac{x+2}{x}}$  is  $e^{k_2}$  then  $15K_1 + K_2$  is equal to

56. If  $[t]$  denotes the greatest integer  $\leq t$ , then the number of points, at which the function

$f(x) = [x + x^3] + |x - x^3| + \left|x + \frac{1}{2}\right|$  is not differentiable in the open interval  $(-10, 10)$  is  $k$

then the sum of the digits of  $k =$  \_\_\_\_\_

57. Let  $\tan^{-1} x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$  for  $x \in \mathbb{R}$  then the number of real solutions of the equation

$\sqrt{1 + \cos 2x} = \sqrt{2} \tan^{-1}(\tan x)$  in the set

$\left(-\frac{5\pi}{2}, -\frac{3\pi}{2}\right) \cup \left(-\frac{3\pi}{2}, -\frac{\pi}{2}\right) \cup \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$  is equal to 'K' then  $K + 10$  equal to

JEE MAIN  
2023

SINGARAJU  
VENKAT KOUNDINYA  
AIR 1  
Sri Chaitanya  
JEE Prep Class  
300  
300  
MAINS

RANK  
1JEE Advanced  
2023

VAIVILALA  
CHIDVILAS REDDY  
AIR 1  
Sri Chaitanya  
JEE Prep Class  
341  
350  
MAINS

RANK  
1NEET  
2023

BORA VARUN  
CHAKRAVARTHI  
AIR 1  
Sri Chaitanya  
JEE Prep Class  
720  
720  
MAINS

RANK  
1





# Sri Chaitanya

Junior Colleges & Coaching Centres

Infinity  
Learn BY SRI CHAITANYA



**VAVILALA CHIDVILAS REDDY**  
HT.NO. 236165088  
Sri Chaitanya  
6<sup>th</sup> - 12<sup>th</sup> Class

**RANK**

WITH ALL INDIA **RANK 1** IN JEE ADVANCED 2023

# SRI CHAITANYA

STANDS AT THE TOP

SEIZES 5 RANKS IN TOP 10 IN ALL-INDIA OPEN CATEGORY

ANDHRA PRADESH STATE TOPPER



8<sup>th</sup> - 12<sup>th</sup> Class

Sri Chaitanya

**2**

RANK



AITS Student

**3**

RANK



9<sup>th</sup> - 12<sup>th</sup> Class

Sri Chaitanya

**4**

RANK



4<sup>th</sup> - 12<sup>th</sup> Class

Sri Chaitanya

**7**

RANK

32 TOP RANKS BELOW 100 IN ALL-INDIA OPEN CATEGORY



RANK

**12**

DASARI SAKETH NAIDU  
HT.No. 236060148



RANK

**14**

GUTHIKONDA ABHIRAM  
HT.No. 239129121



RANK

**17**

P GNANA KOUSIK REDDY  
HT.No. 236133320



RANK

**18**

D VENKATA YUGESH  
HT.No. 236043353



RANK

**19**

MOULIK JINDAL  
HT.No. 237021004



RANK

**22**

D PRATAP SINGH  
HT.No. 232006138



RANK

**23**

SHAIK SAHIL CHANDA  
HT.No. 236123370



RANK

**26**

SONI MAYANK  
HT.No. 232097025



RANK

**33**

GOYAL NAMAN  
HT.No. 232029345



RANK

**37**

SUMEDH S  
HT.No. 231023088



RANK

**38**

V DEERAJ SATHVIK REDDY  
HT.No. 238957161



RANK

**44**

V VERGIYA KAUSHAL  
HT.No. 231004058



RANK

**46**

SUTHAR HARSHAL  
HT.No. 231005060



RANK

**58**

ABHI JAIN  
HT.No. 232001036



RANK

**61**

RIDAM JAIN  
HT.No. 232006021



RANK

**62**

SUTHAR HARSH  
HT.No. 232000028



RANK

**63**

SOJITRA DIPEN  
HT.No. 231071001



RANK

**65**

D PANEENDRANADHA REDDY  
HT.No. 236060288



RANK

**67**

GUNDA SUSHANTH  
HT.No. 231212972



RANK

**68**

HARSH TAYA  
HT.No. 237021718



RANK

**69**

KRISH GUPTA  
HT.No. 232008189



RANK

**73**

KUNCHAKURTHY VIVEK  
HT.No. 236134083



RANK

**81**

N DHARMA TEJA REDDY  
HT.No. 236050138



RANK

**84**

S VENKAT KOUNDINYA  
HT.No. 231030679



RANK

**87**

MRUNAL SHRIKANT  
HT.No. 231041138



RANK

**92**

SAMOTA APURVA  
HT.No. 232099210



RANK

**97**

N PRINCE BRANHAM REDDY  
HT.No. 236064112

BELOW  
**20**  
All India Open  
Category Ranks

BELOW  
**100**  
All India Open  
Category Ranks

BELOW  
**32**  
All India Open  
Category Ranks

BELOW  
**1000**  
All India Open  
Category Ranks

BELOW  
**181**  
All India Open  
Category Ranks

BELOW  
**100**  
All India Open  
Category Ranks

BELOW  
**89**  
All India Open  
Category Ranks

BELOW  
**1000**  
All India Open  
Category Ranks

BELOW  
**699**  
All India Open  
Category Ranks

BELOW  
**3621**  
All India Open  
Category Ranks

NUMBER OF RANKS QUALIFIED  
**3,621**

ADMISSIONS OPEN » JEE ADVANCED LONG-TERM 2024

040 66 06 06 06



Scan QR Code for  
JEE Advanced Offers