

Sec: XI_IIT_IC
Time: 3 Hrs.

FTM – 01

Date:03-06-2024
Max.Marks:300

JEE MAIN 2022 Model

PHYSICS

Section	Question Type	+Ve Marks	-Ve Marks	No.of Qs.	Total marks
Sec – I (Q.N : 1 – 20)	Questions with Single Correct Choice	4	–1	20	80
Sec–II (Q.N : 21 – 30)	Questions with Numerical Answer Type(+ / –Number)	4	–1	10	20
Total				30	100

CHEMISTRY

Section	Question Type	+Ve Marks	-Ve Marks	No.of Qs.	Total marks
Sec – I (Q.N : 31 – 50)	Questions with Single Correct Choice	4	–1	20	80
Sec–II (Q.N : 51 – 60)	Questions with Numerical Answer Type (+ / – Number)	4	–1	10	20
Total				30	100

MATHEMATICS

Section	Question Type	+Ve Marks	-Ve Marks	No.of Qs.	Total marks
Sec–I (Q.N : 61 – 80)	Questions with Single Correct Choice	4	–1	20	80
Sec–II (Q.N : 81 – 90)	Questions with Numerical Answer Type (+ / – Number)	4	–1	10	20
Total				30	100

Exam Syllabus

PHYSICS	50%: MATHEMATICAL PHYSICS : Application of differentiation in physics, Maxima & Minima Integration and its application in physics. 50%: Unit and Dimension : Introduction of physics, physical quantities and physical laws Fundamental and derived quantities and ways to write dimension, Unit and Dimension Uses of dimensional analysis (Principle of homogeneity)
CHEMISTRY	50% : Nature of Matter, Properties of matter and their measurement SI unit, Mass weight, Volume temperature, Need for standard reference, Scientific notation, Precision and accuracy, Significant figure and calculation involving significant figure. Law of chemical combination : (i) Law of conservation of mass, (ii) Law of constant combination, (iii) Law of multiple proportion, (v) Gay Lussac's law, (vi) Avagadro law, (vii) Dalton's atomic theory Atomic mass 50% : Mole concept: Calculation based on mole concept, concept of gram atom, gram molecule Percentage by mass (w/w), percentage by volume (v/v)
MATHEMATICS	50%: SEQUENCE AND SERIES Introduction of Sequence and Series, nth term of AP. Properties of A.P. , Sum of n term of an AP, Arithmetic Mean. 50%: nth term of G.P., sum of n term of GP. Properties of G.P., Geometric mean, A.G.P.

Rough Work

PHYSICS**Max.Marks:100****SECTION – I****(SINGLE CORRECT ANSWER TYPE)**

This section contains 20 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which **ONLY ONE** option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 if not correct.

- A metallic disc is being heated its area A (in m^2) at any time t (in second) is given by $A = 5t^2 + 4t + 8$. Calculate the rate of increase in area at $t = 3$ sec.
(A) $34 \text{ m}^2/\text{s}$ (B) $40 \text{ m}^2/\text{s}$ (C) $80 \text{ m}^2/\text{s}$ (D) $36 \text{ m}^2/\text{s}$
- If $y = 5x^2 - 10x + 5$. Find the value of x for which given function give minimum value.
(A) $x = 2$ (B) $x = 1$ (C) $x = 4$ (D) $x = 6$
- The volume of a sphere is given by $V = \frac{4}{3}\pi R^3$. Where R is the radius of the sphere find the rate of change of volume with respect to R
(A) πR^2 (B) $3\pi R^2$ (C) $4\pi R^2$ (D) $4\pi R^3$
- If $y = \cos^2 x$ then find $\frac{dy}{dx}$
(A) $\sin 2x$ (B) $\cos 2x$ (C) $-\cos 2x$ (D) $-\sin 2x$
- Dimension of work is
(A) $[M L^2 T^{-2}]$ (B) $[M^0 L^2 T^{-2}]$ (C) $[M L^0 T^{-2}]$ (D) $[M L^2 T^{-3}]$
- A force is given by $F = at + bt^2$. Where t is time what are dimension of a and b ?
(A) $[MLT^{-3}]$ and $[MLT^{-4}]$ (B) $[MLT^{-1}]$ and $[MLT^0]$
(C) $[MLT^{-4}]$ and $[MLT]$ (D) $[MLT^{-3}]$ and $[ML^2T^{-4}]$
- Dimension of electrical resistance is
(A) $[ML^{-1}T^3A^{-2}]$ (B) $[ML^3T^{-3}A^{-2}]$ (C) $[ML^2T^{-3}A^{-2}]$ (D) $[ML^2T^{-3}A^{-1}]$
- The dimension formula of relative density is
(A) $[ML^{-3}T^0]$ (B) $[M^0LT^{-1}]$ (C) $[MLT^{-2}]$ (D) Dimensionless
- Which of following pairs have same dimensions?
(A) Angular momentum and work (B) Torque and work
(C) Energy and young modulus (D) Force and mass
- If the Planck's constant h , velocity of light C and time T are taken as fundamental units, then the dimension of force in the new system will be
(A) $hc^{-1}T^{-1}$ (B) $hc^{-1}T^2$ (C) $hc^{-1}T^{-2}$ (D) $h^{-1}c^{-1}T^{-2}$.
- The work done 'W' by a body varies with displacement 'x' as $W = Ax + \frac{B}{(C-x)^2}$. The dimensional formula for 'B' is
(A) $[ML^2T^{-2}]$ (B) $[ML^4T^{-2}]$ (C) $[MLT^{-2}]$ (D) $[MLT^{-2}]$

Rough Work

12. If the velocity v (in cm/s) of a particle is given in terms of time t (in sec) by the equation $v = at + \frac{b}{t+c}$, then the dimensions of a , b and c are
- | a | b | c |
|-----------------|--------|----------|
| (A) $[L^2]$ | $[T]$ | $[LT^2]$ |
| (B) $[LT^2]$ | $[LT]$ | $[L]$ |
| (C) $[LT^{-2}]$ | $[L]$ | $[T]$ |
| (D) $[L]$ | $[LT]$ | $[T^2]$ |
13. If energy E , Velocity v and time T are taken as fundamental quantities, the dimensional formula for surface tension is
- (A) $[Ev^{-2}T^{-2}]$ (B) $[E^2vT^{-2}]$ (C) $[E^2vT^{-2}]$ (D) $[E^{-2}v^{-2}T^{-1}]$
14. The Richardson equation is given by $I = AT^2e^{-B/KT}$, where A and B are constants. The dimensional formula for AB^2 is same as
- (A) IT^{-2} (B) kT (C) Ik^2 (D) Ik^2/T
15. If $\eta = \frac{A}{B} \log(Bx + C)$ is dimensionally true, then (here η is the coefficient of viscosity and x is the distance)
- (A) C is dimensionless constant (B) B has dimension $[M^0L^{-1}T^0]$
 (C) **A has dimension $[ML^{-2}T^{-1}]$** (D) All are true
16. If $y = 2x^3 + 3x^2 + 6x + 1$, then $\frac{dy}{dx}$ will be
- (A) $6(x^2 + x + 1)$ (B) $6(x^2 + x + 2)$ (C) $6x^2 + 3x$ (D) $(x^2 + 6x + 1)$
17. If $\int \left[(x)^{1/3} - \frac{1}{(x)^{1/3}} \right] dx$ is equal to
- (A) $x^{4/3} - x^{2/3} + c$ (B) $\frac{4}{3}x^{2/3} - \frac{2}{3}x^{2/3} + c$
 (C) $\frac{3}{4}x^{4/3} - \frac{2}{3}x^{1/3} + c$ (D) $\frac{3}{4}(x)^{4/3} - \frac{3}{2}(x)^{2/3} + c$
18. If $Q = 4v^3 + 3v^2$, then the value of ' v ' such that, there exist maxima of ' Q '
- (A) 0 (B) $-\frac{1}{2}$ (C) $\frac{1}{2}$ (D) None
19. $\int 2\sin(x) dx$ is equal to:
- (A) $-2\cos x + C$ (B) $2\cos x + C$ (C) $-2\cos x$ (D) $2\cos x$

Rough Work

20. If $y = \sin(x) + \ln(x^2) + e^{2x}$ then $\frac{dy}{dx}$ will be
- (A) $\cos x + \frac{2}{x} + 2e^{2x}$ (b) $\cos x - \frac{2}{x} + 2e^{2x}$ (C) $\cos x + \frac{2}{x} - 2e^{2x}$ (D) $\cos x - \frac{2}{x} - 2e^{2x}$

SECTION-II

(Numerical Value Answer Type)

This section contains 10 questions, but candidates have to attempt any 5 questions out of 10. The answer to each question is a Numerical values comprising of positive or negative decimal numbers (place value ranging from Thousands Place to Hundredths Place).

Eg: 1234.56, 123.45, -123.45, -1234.56, -0.12, 0.12 etc.

Marking scheme: +4 for correct answer, -1 in all other cases.

21. If $y = \sin \theta$ then $\frac{dy}{d\theta}$ at $\theta = 60^\circ$ is $\frac{1}{x}$ where x is
22. Integrate the functions: $\int_1^2 (2t - 2) dt$:
23. Find the value of x for which function $y = x^3 - 3x^2 + 6$ will give minimum value
24. If $x = 5 + 4t + 3t^2$ find $\frac{dx}{dt}$ at $t = 0$
25. If velocity (v) of a body in time t is given by $v = AT^3 + BT^2 + CT + D$ and dimension of C is LT^{-x} where x is
26. The speed of a wave produced in water is given by $v = \lambda^a g^b \rho^c$. Where λ, g and ρ are wavelength of wave, acceleration due to gravity and density of water respectively. Then find the value of $a + b + c$.
27. The air bubble formed by explosion inside water performed oscillation with time period T that is directly proportional to $P^a d^b E^c$. Where p is pressure, d is density & E is the energy due to explosion if Value of b is $1/x$. Where x is
28. $\int_0^1 (3x^2 + 4) dx = I$. Find the value of I.
29. $y = x^4 + 3x^2 + \pi + 2$. the $\left. \frac{dy}{dx} \right|_{x=1} = 2n$ then n is
30. Frequency of sound that can be produced by a pipe depends on length (l) of the pipe atmospheric pressure (p) and density (d) of air, according to relation $v = \frac{p^b d^c}{l^a}$. The value of (a + b + c) is

Rough Work

CHEMISTRY**Max.Marks:100****SECTION – I****(SINGLE CORRECT ANSWER TYPE)**

This section contains 20 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which **ONLY ONE** option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

31. The percentage composition of carbon by mole in methane is
(A) 75% (B) 80% (C) 25% (D) 20%
32. Which of the following set of compounds illustrate law of multiple proportions
(A) CO_2 , CH_4 , H_2O (B) H_2O , D_2O , HOD (C) N_2O , N_2O_5 , N_2O_3 (D) All of these
33. Caffeine has a molecular weight of 194. If it contains 28.9% by mass of nitrogen, number of atoms of nitrogen in one molecule of caffeine is:
(A) 4 (B) 6 (C) 2 (D) 3
34. The formula of an acid is HXO_2 . The mass of 0.0242 moles of the acid is 1.657 gm. What is the atomic weight of 'X'?
(A) 35.5 (B) 28.1 (C) 128 (D) 19
35. The number of neutron in 8.5 gm of NH_3 are
(A) $50 N_A$ (B) $5 N_A$ (C) $3.5 N_A$ (D) $35 N_A$
36. If we consider that $1/6$ in place of $1/12$ mass of carbon atom as the relative atomic mass unit, the mass of one mole of the substance will
(A) be a function of the molecular mass of the substance
(B) remain unchanged
(C) increase two fold
(D) decrease twice
37. Number of atoms in 635.46 gm of Cu (At. Wt. of Cu = 63.54) is
(A) twice that in 60 gm of carbon (B) 6.023×10^{23}
(C) half that in 8 gm He (D) $635.46 \times 6.023 \times 10^{23}$
38. Assuming fully decomposed, the volume of CO_2 released at STP on heating 9.85 gm of BaCO_3 will be: (Atomic mass of Ba = 137 gm)
(A) 0.84 L (B) 2.24 L (C) 4.06 L (D) 1.12 L
39. Number of atoms in the following sample of substance is largest in
(A) 4 gm of hydrogen (B) 71 gm of chlorine (C) 127 gm of Iodine (D) 48 gm of magnesium

Rough Work

40. 1.5 moles of oxygen atoms are present in
(A) 0.5 moles of BaCO_3 (B) 1 mole of BaCO_3
(C) 2 moles of BaCO_3 (D) 0.25 moles of BaCO_3
41. The density of water is 1 gm/mL. What is the volume occupied by 1 molecule of water?
(A) 1.44×10^{-23} mL (B) 1 mL (C) 18 mL (D) 2.988×10^{-23} mL
42. A compound was found to contain 5.37% nitrogen. What is the minimum molecular weight of compound?
(A) 26.07 (B) 2.607 (C) 260.7 (D) 280.5
43. The number of electrons present in 9.5 gm of PO_4^{3-} is (Atomic number of P = 15)
(A) $6N_A$ (B) $5 N_A$ (C) $0.1 N_A$ (D) $4.7 N_A$
44. 20 gm of an ideal gas containing only atoms of 'S' and 'O' occupies 5.6 L at 1 atm and 273 K. What is the molecular weight of gas?
(A) 64 (B) 80 (C) 96 (D) 85
45. The total number of AlF_3 molecules in a sample of AlF_3 containing 3.01×10^{23} ions of F^- is
(A) 9×10^{24} (B) 3×10^{24} (C) 7.5×10^{23} (D) 10^{23}
46. Rearrange the following (I to IV) in the increasing order of masses:
(I) 0.5 mole of O_3 (II) 0.5 gm atom of Oxygen
(III) 3.011×10^{23} molecules of O_2 (IV) 5.6 litre of CO_2 at STP
(A) $\text{II} < \text{IV} < \text{III} < \text{I}$ (B) $\text{II} < \text{I} < \text{IV} < \text{III}$ (C) $\text{IV} < \text{II} < \text{III} < \text{I}$ (D) $\text{I} < \text{II} < \text{III} < \text{IV}$
47. Amount of oxygen in 32.2 g of $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ is
(A) 20.8 gm (B) 27.71 gm (C) 2.24 gm (D) 22.4 gm
48. Weight of one atom of an element is 6.644×10^{-23} gm. The g-atoms of element in 40 kg is
(A) 10^2 (B) 10^3 (C) 10^4 (D) 10^5
49. One litre of CO_2 is passed over hot coke. The volume becomes 1.4 L. The percent composition of product is
$$\text{CO}_2(g) + \text{C}(s) \rightarrow 2\text{CO}(g)$$

(A) 0.6 litre CO (B) 0.8 litre CO_2
(C) 0.6 litre CO_2 & 0.8 litre CO (D) 0.5 L CO_2 & 0.5 L CO
50. Equal masses of O_2 , H_2 & CH_4 are taken in a container. The respective mole ratio of these gases in container is
(A) 1 : 16 : 2 (B) 16 : 1 : 2 (C) 1 : 2 : 16 (D) 16 : 2 : 1

Rough Work

SECTION-II**(Numerical Value Answer Type)**

This section contains 10 questions. The answer to each question is a Numerical value. If the Answer is in the decimals, **Mark the nearest Integer only. Have to Answer any 5 only out of 10 questions** and question will be evaluated according to the following marking scheme:

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

51. An organic compound gives 0.220 g of CO_2 and 0.126 g of H_2O on complete combustion. If the % of carbon is 24 then the % of hydrogen is $x \times 10^{-1}$. Find the value of x .
52. The mass (in gm) of Na_2SO_4 would contain the same number of total ions as present in 930 gm of $\text{Ca}_3(\text{PO}_4)_2$ is $x \times 10^2$ gm. Find $\frac{(x-2)}{5}$.
53. The number of significant figure in result for given calculation of 0.00340×2305 is.
54. When 0.01 mol of an organic compound containing 60% carbon was burnt completely 4.4 gm of CO_2 was produced. The mass of compound is _____ gm/mole.
55. 16 gm of SO_x occupies 5.6 litre at STP assuming ideal gas nature. The value of x is _____.
56. When 45 mL of HCl is dissolved in 455 ml of H_2O . Find its v/v%.
57. The total number of atoms in 11.2 litre of CH_4 gas at NTP is $3x \times 10^{23}$. Find the value of x .
($N_A = 6.02 \times 10^{23}$)
58. The number of moles of electrons present in 1 kg of electron sample is $x \times 10^5$. Find the value of x . (Given: $N_A = 6 \times 10^{23}$, mass of electron = 9.1×10^{-31} kg).
59. 200 gm of CaCO_3 sample with 40% impurity is heated to give $1.344 x \times 10^y$ ml of $\text{CO}_2(\text{g})$ at NTP. Find $x + y$.
60. An element A is tetravalent & another B is divalent. The formula of compound is A_xB_y . Find the value of $(4x - y)$.

Rough Work

MATHEMATICS**Max.Marks:100****SECTION – I****(SINGLE CORRECT ANSWER TYPE)**

This section contains 20 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which **ONLY ONE** option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

61. Let the sum of n , $2n$, $3n$ terms of an AP be S_1 , S_2 and S_3 respectively, then $3(S_2 - S_1)$ is
(A) $2S_3$ (B) S_3 (C) $4S_3$ (D) None of these
62. If the sum and product of the first three terms in an A.P. are 33 and 1155, respectively, then a value of its 11^{th} term is:
(A) -35 (B) 25 (C) -36 (D) -25
63. An infinite GP has the first term 'x' and sum 5, then x belongs to
(A) $x < -10$ (B) $-10 < x < 0$ (C) $0 < x < 10$ (D) $x > 10$
64. If $x = 1 + a + a^2 + \dots \infty$ and $y = 1 + b + b^2 + \dots \infty$ where a and b are proper fractions, then $1 + ab + a^2b^2 + \dots \infty$ equals
(A) $\frac{xy}{y+x-1}$ (B) $\frac{x+y}{x-y}$ (C) $\frac{x^2+y^2}{x-y}$ (D) None of these
65. If a, b and c be three distinct real numbers in GP and $a + b + c = xb$, then x cannot be
(A) 4 (B) 2 (C) -2 (D) -3
66. pth term of an A.P. is q and qth term is p, its $(p + q)^{\text{th}}$ term is $(p \neq q)$
(A) 0 (B) $p + q$ (C) $-(p + q)$ (D) None of these
67. If the sum of first n terms of an A.P. is $Pn + Qn^2$ where P and Q are constants, then common difference of A.P. will be
(A) $P + Q$ (B) $P - Q$ (C) $2P$ (D) $2Q$
68. The nth term of a sequence whose sum of n terms is $5n^2 + 2n$, is
(A) $10n + 3$ (B) $10n - 5$ (C) $10n - 3$ (D) $5n + 6$
69. If the arithmetic mean of a and b is $\frac{a^n + b^n}{a^{n-1} + b^{n-1}}$ then the value of n is
(A) -1 (B) 0 (C) 1 (D) -2
70. If b_1 , b_2 and b_3 ($b_1 > 0$) are three successive terms of a G.P. with common ratio r, then complete set of value of r for which the inequality $b_3 > 4b_2 - 3b_1$ holds, is
(A) $(-\infty, 1)$ (B) $(-\infty, 1) \cup (3, \infty)$ (C) $(3, \infty)$ (D) $(1, 3)$
71. The product of three geometric means between 4 and $\frac{1}{4}$ is
(A) 1 (B) 2 (C) 3 (D) 4

Rough Work

72. If $a_1, a_2, \dots, a_n$ are in AP with common difference $d \neq 0$ then $(\sin d)$
 $[\sec a_1 \sec a_2 + \sec a_2 \sec a_3 + \dots + \sec a_{n-1} \sec a_n]$ is equal to
 (A) $\cot a_n - \cot a_1$ (B) $\cot a_1 - \cot a_n$ (C) $\tan a_n - \tan a_1$ (D) $\tan a_n - \tan a_{n-1}$
73. If a, b, c are in A. P. as well as in G. P., then
 (A) $a = b = c$ (B) $a \neq b = c$ (C) $a = b \neq c$ (D) $a \neq b \neq c$
74. If $(1 - p)(1 + 3x + 9x^2 + 27x^3 + 81x^4 + 243x^5) = 1 - p^6$, $p \neq 1$, then value of $\frac{p}{x}$ is
 (A) $\frac{1}{3}$ (B) 3 (C) $\frac{1}{2}$ (D) 2
75. Let $a_1, a_2, a_3, \dots$ be an A.P. if $\frac{a_1 + a_2 + \dots + a_{10}}{a_1 + a_2 + \dots + a_p} = \frac{100}{p^2}$, $p \neq 10$, then $\frac{a_{11}}{a_{10}}$ is equal to
 (A) $\frac{19}{21}$ (B) $\frac{100}{121}$ (C) $\frac{21}{19}$ (D) $\frac{121}{100}$
76. The sum of the infinite series $1 + \frac{2}{3} + \frac{7}{3^2} + \frac{12}{3^3} + \frac{17}{3^4} + \frac{22}{3^5} + \dots$ is equal to
 (A) $\frac{13}{4}$ (B) $\frac{9}{4}$ (C) $\frac{15}{4}$ (D) $\frac{11}{4}$
77. In a G.P., if the $(p + q)^{\text{th}}$ term is m and $(p - q)^{\text{th}}$ term is n , then its p^{th} term is
 (A) 0 (B) mn (C) $\sqrt{mn}$ (D) $\frac{1}{2}(m+n)$
78. If the 2^{nd} , 5^{th} and 9^{th} terms of a non-constant A.P. are in G.P., then the common ratio of this G.P. is:
 (A) $\frac{8}{5}$ (B) $\frac{4}{3}$ (C) $\frac{3}{4}$ (D) $\frac{7}{4}$
79. If $S = \frac{7}{5} + \frac{9}{5^2} + \frac{13}{5^3} + \frac{19}{5^4} + \dots$, then $160 S$ is equal to
 (A) 305 (B) 310 (C) 315 (D) 320
80. If a, b, c are in G.P; a, x, b are in A.P. and also b, y, c are in A.P., then $\frac{a}{x} + \frac{c}{y}$ is equal to
 (A) 1 (B) 2 (C) $\frac{1}{2}$ (D) $\frac{3}{2}$

Rough Work

SECTION-II**(Numerical Value Answer Type)**

This section contains 10 questions. The answer to each question is a Numerical value. If the Answer is in the decimals, **Mark the nearest Integer only. Have to Answer any 5 only out of 10 questions** and question will be evaluated according to the following marking scheme:

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

81. If $a_1, a_2, a_3, \dots, a_{24}$ are in arithmetic progression and $a_1 + a_5 + a_{10} + a_{15} + a_{20} + a_{24} = 225$, then $a_1 + a_2 + a_3 + \dots + a_{23} + a_{24}$ is equal to
82. If the sum of first $2n$ terms of A.P. 2, 5, 8,... is equal to the sum of the first n terms of the A.P. 57, 59, 61,, then n equals
83. If the sum of n terms of an AP is given by $S_n = 3n + 2n^2$, then the common difference of the AP is
84. The value of $9^{1/3} \times 9^{1/9} \times 9^{1/27} \times \dots \infty$ is equal to
85. In an increasing geometric series, the sum of the second and the sixth term is $\frac{25}{2}$ and the product of the third and fifth term is 25. Then, the sum of 4^{th} , 6^{th} and 8^{th} term is equal to:
86. The sum of first four terms of a geometric progression (G.P.) is $\frac{65}{12}$ and the sum of their respective reciprocals is $\frac{65}{18}$. If the product of first three terms of the G.P. is 1, and the third term is α , then 2α is
87. If A_1 be the A.M. and G_1, G_2 be two G.M.'s between two positive numbers then $\frac{G_1^3 + G_2^3}{G_1 G_2 A_1}$ is equal to:
88. If the sum of first n natural numbers is $1/5$ times the sum of their squares, then the value of n is
89. Let T_r be the r^{th} term of an A.P. whose first term is a and common difference is d . If for some positive integers $m, n, m \neq n, T_m = \frac{1}{n}$ and $T_n = \frac{1}{m}$, then $a - d =$
90. The product of three consecutive terms of a G.P, is 512. If 4 is added to each of the first and the second of these terms, the three terms now form an A.P. Then the sum of the original three terms of the given G.P, is:

Subject	Physics	Chemistry	Mathematics
Setter Branch	DEL-FAR		
SME Name	MR. RAHUL KUMAWAT	MR. KUNDAN KUMAR	MR. R.K. SEHGAL
Phone Number			

Rough Work