Experiment 7

:. Random Sampling and Sampling Distribution

Code

```
import numpy as np
import matplotlib.pyplot as plt
# Step 1: Generate a population (e.g., normal distribution)
population_mean = 50
population_std = 10
population_size = 100000
population = np.random.normal(population_mean, population_std, population_size)
# Step 2: Random sampling
sample_sizes = [30, 50, 100] # different sample sizes to consider
num_samples = 1000 # number of samples for each sample size 26
sample_means = {}
for size in sample_sizes:
sample_means[size] = []
for _ in range(num_samples):
sample = np.random.choice(population, size=size, replace=False)
sample_means[size].append(np.mean(sample))
# Step 3: Plotting sampling distributions
plt.figure(figsize=(12, 8))
for i, size in enumerate(sample_sizes):
plt.subplot(len(sample_sizes), 1, i+1)
plt.hist(sample_means[size], bins=30, alpha=0.7, label=f'Sample Size {size}')
plt.axvline(np.mean(population), color='red', linestyle='dashed', linewidth=1.5, label='Population
Mean')
plt.title(f'Sampling Distribution (Sample Size {size})')
plt.xlabel('Sample Mean')
plt.ylabel('Frequency')
plt.legend()
```

```
plt.tight_layout()
plt.show()
```

OUTPUT:

