

One task provoked symptoms, and convergence was outside the normal range

PROVOCATION IS A RISE OF 2+ POINTS OVER YOUR PRE-TEST RATING

Smooth pursuit	NO RISE
Horizontal saccades	NO RISE
Vertical saccades	NO RISE
Near point of convergence	PROVOKED +3 8.4 CM (≥ 5)
Horizontal VOR	NO RISE
Vertical VOR	NO RISE
Visual motion sensitivity	NO RISE

Your pre-test ratings were headache 4, dizziness 3, nausea 1, foggiess 2. Everything above is measured against those, which is why the screening asks for them first.

None of this waits. Go to an emergency department now if you have any of these:

- Neck pain or tenderness
- Double vision
- Weakness, numbness or tingling in the arms or legs
- Severe or increasing headache
- Seizure or convulsion
- Loss of consciousness
- Worsening confusion, agitation or unusual behaviour
- Repeated vomiting

WHAT TO TRY NEXT

Activity — stage 1: Symptom-limited activity

Daily activities that do not provoke symptoms more than mildly.

School or work — stage 2: School work at home

Homework in 20-30 minute blocks with real breaks between them.

One task provoked symptoms, and convergence was outside the normal range. Start at symptom-limited activity and school work at home. This is a starting floor, not a diagnosis.

See a clinician about this screening.

- One of the seven tasks provoked symptoms by 2 points or more.
- Convergence measured at 8.4 cm, at or beyond the 5 cm cut point.
- Your symptom inventory scored in the moderate band, from 4 of 22 items.

Baseline is a screening aid. It does not diagnose a concussion and it cannot clear you to return to contact — only a clinician who examines you can do that.