



PATUAKHALI SCIENCE AND TECHNOLOGY UNIVERSITY

**Group
Assignment of MAT-**

211

Group Assignment of MAT-211

Submitted By:

Name	ID
Aishwariya Sarder	2102012
Mehedi Hasan Nazmul	2102013
Noushin Rahman	2102014
Md Riadul Islam Mahi	2102015
Md. Mehedi Hasan	2102016
Durjoy Das	2102017
Jannatul Mim	2102018
Hridoy Chandra Sarker	2102019
Md. Sadman Kabir Bhuiyan	2102020

Submitted to:

MUHAMMAD MASUDUR RAHMAN

Associate Professor of Mathematics Department

Faculty of Computer Science And Engineering

Equation Reducible to Homogenous

An equation that is reducible to a homogeneous form is one that, through some transformations, can be converted into a form where the function can be expressed as a ratio of variables, often leading to a separable equation.

Homogeneous Differential Equations

A first-order ordinary differential equation (ODE) of the form:

$$M(x,y) dx + N(x,y) dy = 0$$

is homogeneous if both $M(x,y)$ and $N(x,y)$ are homogeneous functions of the same degree. A function $f(x,y)$ is homogeneous of degree n if:

$$f(tx,ty) = t^n f(x,y)$$

for any $t \in \mathbb{R}$.

Transformations to Homogeneous Form

Sometimes, an equation that is not explicitly in homogeneous form can be transformed into one that is. One common transformation is the substitution $y=vx$ where $v=y/x$.

Example: Reduction to Homogeneous Form

Consider the differential equation:

$$\frac{dy}{dx} = \frac{ax+by+c}{dx+ey+f}$$

This is not homogeneous as is. However, we can make a substitution to reduce it to a homogeneous form.

1. **Substitution:** $y=vx$

$$\frac{dy}{dx} = v + x \frac{dv}{dx}$$

2. **Rewrite the Original Equation:** Substitute $y=vx$ into the original equation:

$$v + x \frac{dv}{dx} = \frac{ax + b(vx) + c}{dx + e(vx) + f}$$

$$v + x \frac{dv}{dx} = \frac{ax + bvx + c}{dx + evx + f}$$

3. **Simplify:**

$$4. \quad v + x \frac{dv}{dx} = \frac{x(a + bv) + c}{x(d + ev) + f}$$

$$v + x \frac{dv}{dx} = \frac{a + bv + \frac{c}{x}}{d + ev + \frac{f}{x}}$$

5. **Analyze the Terms:**

If c and f are zero, the equation becomes:

$$v + x \frac{dv}{dx} = \frac{a + bv}{d + ev}$$

This is a homogeneous differential equation in terms of v and x .

Solving the Homogeneous Form

1. **Separate Variables:**

$$x \frac{dv}{dx} = \frac{a + bv}{d + ev} - v$$

$$x \frac{dv}{dx} = \frac{a + bv - vd - v^2 e}{d + ev}$$

$$x \frac{dv}{dx} = \frac{a-vd}{d+ev}$$

2. Integrate:

$$\int \frac{d+ev}{a-vd} dv = \int \frac{1}{x} dx$$

Example-1: Solve the DE $dy/dx = (x+2y-1)/(x+2y+1)$.

Solution:

Note that h, k do not exist in this case which can reduce this DE to homogeneous form. Thus, we use the substitution

$$x+2y=v$$

$$\Rightarrow 1+2 \, dy/dx$$

Thus, our DE becomes

$$1/2(dv/dx-1) = (v-1)/(v+1)$$

$$\Rightarrow dv/dx = (2v-2)/(v+1) + 1$$

$$= (3v-1)/(v+1)$$

$$\Rightarrow (v+1)/(3v-1) dv = dx$$

$$\Rightarrow 1/3(1+4/(3v-1)) dv = dx$$

Integrating, we have

$$1/3(v+4/3 \ln(3v-1)) = x + C_1$$

Substituting $v=x+2y$, we have

$$x+2y+4/3 \ln(3x+6y-1) = 3x + C_2$$

$$\Rightarrow y-x+2/3 \ln(3x+6y-1) = C$$

Example 2: Reducible to homogeneous differential equation.

Solve the DE $\frac{dy}{dx} = \frac{2y - x - 4}{y - 3x + 3}$

Solution: We substitute $x \rightarrow X+h$ and $y \rightarrow Y+k$ where h, k need to be determined :

$$\frac{dy}{dx} = \frac{dY}{dX} = \frac{\{(2Y - X) + (2K - h - 4)\}}{\{(Y - 3X) + (k - 3h + 3)\}}$$

h and k must be chosen so that

$$2k - h - 4 = 0$$

$$k - 3h + 3 = 0$$

This gives $h=2$ and $k=3$. Thus,

$$x = X + 2$$

$$y = Y + 3$$

Our DE now reduces to

$$\frac{dY}{dX} = \frac{2Y - X}{Y - 3X}$$

Using the substitution $Y=vX$, and simplifying, we have (verify),

$$\frac{v - 3}{v^2 - 5v + 1} dv = \frac{-dX}{X}$$

We now integrate this DE which is VS; the left-hand side can be integrated by the techniques described in the unit on Indefinite Integration.

Finally, we substitute $v = \frac{Y}{X}$ and

$$X = x - 2$$

$$Y = y - 3$$

to obtain the general solution.

Suppose our DE is of the form

$$\frac{dy}{dx} = f\left(\frac{ax + by + c}{dx + ey + f}\right)$$

We try to find h, k so that

$$ah + bk + c = 0$$

$$dh + ek + f = 0$$

What if this system does not yield a solution? Recall that this will happen if $\frac{a}{b} = \frac{d}{e}$. How do we reduce the DE to a homogeneous one in such a case?

let $\frac{a}{d} = \frac{b}{e} = \lambda$ (say). Thus,

$$\frac{ax + by + c}{dx + ey + f} = \frac{\lambda(dx + ey) + c}{dx + ey + f}$$

This suggests the substitution $dx + ey = v$, which'll give

$$\begin{aligned} d + e \frac{dy}{dx} &= \frac{dv}{dx} \\ \Rightarrow \frac{dy}{dx} &= \frac{1}{e} \left(\frac{dv}{dx} - d \right) \end{aligned}$$

Thus, our DE reduces to

$$\begin{aligned}\frac{1}{e} \left(\frac{dv}{dx} - d \right) &= \frac{\lambda v + c}{v + f} \\ \Rightarrow \frac{dv}{dx} &= \frac{\lambda e v + e c}{v + f} + d \\ &= \frac{(\lambda e + d)v + (e c + d f)}{v + f} \\ &\Rightarrow \frac{v + f}{(\lambda e + d)v + (e c + d f)}\end{aligned}$$

which is in VS form and hence can be solved.

Example 3. Solve $(3x-7y-3) \frac{dy}{dx} = 3y - 7x + 7$

Solution: $\frac{dy}{dx} = \frac{3y-7x+7}{3x-7y-3}$

Put $x = X+h$, $y = Y+k$, where h, k are some constants. Then $\frac{dy}{dx} = \frac{dY}{dX}$

And the given eq. becomes, $\frac{dY}{dX} = \frac{3Y-7X+(3K-7h+7)}{3X-7Y+(3h-7k-3)}$

Choose h, k such that $3h-7k-3=0$ and $3k-7h+7=0$, which give $h=1, k=0$.

$$\therefore \frac{dY}{dX} = \frac{3Y-7X}{3X-7Y}$$

put $Y = vX$, $\frac{dY}{dX} = v + X \frac{dv}{dX}$

$$\therefore v + X \frac{dv}{dX} = \frac{3v-7X}{3X-7vX} = \frac{3v-7}{3-7v}$$

$$\text{Or, } X \frac{dv}{dX} = \frac{3v-7}{3X-7vX} - v = \frac{7(v^2-1)}{3-7v}$$

$$\text{Or, } \frac{7dX}{X} = \frac{3-7v}{v^2-1} dv = - \left(\frac{2}{v-1} + \frac{5}{v+1} \right) dv$$

Integrating, $7 \log X = -2 \log(v-1) - 5 \log(v+1) + \log C$

$$\text{Or, } X^7(v-1)^2(v+1)^5 = C$$

$$\text{Or, } X^7 \left(\frac{Y}{X} - 1 \right)^2 \left(\frac{Y}{X} + 1 \right)^5 = C \text{ as } Y = vX$$

$$\text{Or, } (Y - X)^2(Y + X)^5 = C$$

$$\text{Or, } (y - x + 1)^2(y + x - 1)^5 = C \text{ as } x = X+1, y = Y+0$$

Example 4: Solve the DE $\frac{dy}{dx} = \frac{x+2y-1}{x+2y+1}$

Solution: Note that h, k do not exist in this case which can reduce this DE to homogeneous form. Thus, we use the substitution

$$x+2y=v$$

$$\Rightarrow 1+2y \frac{dy}{dx} = \frac{dv}{dx}$$

Thus, our DE becomes

$$\frac{1}{2} \left(\frac{dv}{dx} - 1 \right) = \frac{v-1}{v+1}$$

$$\Rightarrow \frac{dv}{dx} = \frac{2v-2}{v+1} + 1$$

$$= \frac{3v-1}{v+1}$$

$$\Rightarrow \frac{v+1}{3v-1} dv = dx$$

$$\Rightarrow \frac{1}{3} \left(1 + \frac{4}{3v-1} \right) dv = dx$$

Integrating, we have

$$\frac{1}{3} \left(v + \frac{4}{3} \ln(3v-1) \right) = x + C_1$$

Substituting $v=x+2y$, we have

$$x + 2y + \frac{4}{3} (\ln 3x + 6y - 1) = 3x + C_2$$

$$\Rightarrow y-x + \frac{2}{3} (\ln 3x + 6y - 1) = C$$

Example 5: Find the solution of the differential equation $\frac{dy}{dx} = \frac{x+y+3}{2x+2y+1}$

Solution: Here, $\frac{a}{a_1} = \frac{b}{b_1} = \frac{1}{2}$ i.e., the coefficients of x and y in the Nr and Dr of the expression for $\frac{dy}{dx}$ are proportional. Proper substitution in this case, therefore, will be to put v for x + y. Let x + y = v. Then, $1 + \frac{dy}{dx} = \frac{dv}{dx}$ with these substitutions the given equation reduces to

$$\frac{dv}{dx} - 1 = \frac{v+3}{2v+1}$$

$$\text{or } \frac{dv}{dx} = \frac{v+3}{2v+1} + 1 = \frac{3v+4}{2v+1}$$

$$\text{Or } dx = \frac{2v+1}{3v+4} dv = \left[\frac{2}{3} - \frac{\frac{5}{3}}{3v+4} \right] dv$$

$$\therefore \text{ On integrating, } x + C = \frac{2}{3}v - \frac{5}{3} \cdot \frac{1}{3} \log(3v + 4)$$

$$\Rightarrow \Rightarrow x + C = \frac{2}{3}v - \frac{5}{9} \log(3v + 4)$$

$$\text{Or } x + C = \frac{2}{3}(x + y) - \left(\frac{5}{9}\right) \log(3x + 3y + 4), \text{ here } v = x + y$$

Which is the required solution.

Example 6. Solve $(2x+y+3) \frac{dy}{dx} = x + 2y + 3$

Solution: $\frac{dy}{dx} = \frac{x+2y+3}{2x+y+3}$

Put $x = X+h$, $y = Y+k$, where h, k are some constants. Then $\frac{dy}{dx} = \frac{dY}{dX}$

And the given eq. becomes, $\frac{dY}{dX} = \frac{X+2Y+(h+2k+3)}{2X+Y+(2h+k+3)}$

Choose h, k such that $(2h + k + 3) = 0$ and $(h + 2k + 3) = 0$, which give $h = -1, k = -1$.

$$\therefore \frac{dY}{dX} = \frac{X+2Y}{2X+Y}$$

put $Y = vX$, $\frac{dY}{dX} = v + X \frac{dv}{dX}$

$$\therefore v + X \frac{dv}{dX} = \frac{X+2vX}{2X+vX}$$

Or, $X \frac{dv}{dX} = \frac{1+2v}{2+v} - v$

$$\text{Or, } \frac{dX}{X} = \frac{2+v}{1-v^2} dv = \left(\frac{3/2}{1-v} + \frac{1/2}{v+1} \right) dv$$

Integrating, $2 \log X = -3 \log(1-v) + \log(v+1) + \log C$

Or, $X^2(1-v)^3(v+1) = C$

Or, $X^2 \frac{(1-Y/X)^3}{(1+Y/X)} = C$

Or, $(X-Y)^3 = C(Y+X)$; where $x = X-1$, $y = Y-1$

Or, $(X-Y)^3 = C(Y+X-2)$ is the solution.

Example 7: Solve $(3x - 3y + 4) \frac{dy}{dx} = 2x - 2y + 4$

Solution: The equation is $(3x - 3y + 6) \frac{dy}{dx} = 2x - 2y + 4$
 $\Rightarrow \frac{dy}{dx} = \frac{2x-2y+4}{(3x-3y+6)}$

Put $x - y = v$, so that $1 - \frac{dy}{dx} = \frac{dv}{dx}$

$$\text{Or } \frac{dy}{dx} = 1 - \frac{dv}{dx}.$$

\therefore The equation becomes

$$\begin{aligned} 1 - \frac{dv}{dx} &= \frac{2v+4}{(3v+6)} \\ \text{or, } \frac{dv}{dx} &= 1 - \frac{2v+4}{(3v+6)} \\ &= \frac{v+2}{3v+6} \\ \text{Or, } dx &= \frac{3v+6}{v+2} dv \end{aligned}$$

\therefore Integrating, $x = 3v + \log(v + 2) + C$

$$\Rightarrow x = 3(x - y) + \log(x - y + 2) + C ; \text{ here } v = x - y$$

$$\Rightarrow 3y - 2x = \log(x - y + 2) + C,$$

Which is the required solution.

Reducible to homogeneous differential equation

Example 1: Solve the DE $\frac{dy}{dx} = \frac{2y-x-4}{y-3x+3}$

Solution: We substitute $x \rightarrow X+h$ and $y \rightarrow Y+k$ where h, k need to be determined

$$\frac{Dy}{dx} = \frac{dY}{dY} = \frac{(2Y-X)+(2k-h-4)}{(Y-3X)+(k-3h+3)}$$

h and k must be chosen so that

$$2k-h-4=0$$

$$k-3h+3=0$$

This gives $h=2$ and $k=3$. Thus,

$$x=X+2$$

$$y=Y+3$$

Our DE now reduces to

$$\frac{dy}{dx} = \frac{2Y-X}{Y-3X}$$

Using the substitution $Y=vX$ and simplifying, we have (verify),

$$\frac{v-3}{v^2-5v+1} dv = \frac{-dX}{X}$$

We now integrate this DE which is VS; the left-hand side can be integrated by the techniques described in the unit on Indefinite Integration.

Finally, we substitute $v=YX$ and

$$X=x-2$$

$$Y=y-3$$

to obtain the general solution.

Suppose our DE is of the form

$$\frac{dY}{dX} = f\left(\frac{ax+by+c}{dx+ey+f}\right)$$

We try to find h, k so that

$$ah+bk+c=0$$

$$dh+ek+f=0$$

Let $ad=be=\lambda$ (say). Thus,

$$\frac{ax + by + c}{dx + ey + f} = \frac{\lambda(dx + ey) + c}{dx + ey + f}$$

This suggests the substitution $dx+ey=v$ which'll give

$$d+e\frac{dy}{dx} = \frac{dv}{dx}$$

$$\frac{dy}{dx} = \frac{1}{e}\left(\frac{dv}{dx} - d\right)$$

Thus, our DE reduces to

$$\begin{aligned} \frac{1}{e}\left(\frac{dv}{dx} - d\right) &= \frac{\lambda v + c}{v + f} \\ \Rightarrow \frac{dv}{dx} &= \frac{\lambda ev + ec}{v + f} + d \\ &= \frac{(\lambda e + d)v + (ec + df)}{v + f} \\ \Rightarrow \frac{(v + f)}{(\lambda e + d)v + ec + df} dv &= dx \end{aligned}$$

which is in VS form and hence can be solved.