



A Note on Prediction with Misspecified Models

Khreshna Syuhada

Statistics Research Division, Institut Teknologi Bandung
Jalan Ganesa 10 Bandung 40132 Indonesia
Email: khreshna@math.itb.ac.id

Abstract. Suppose that a time series model is fitted. It is likely that the fitted model is not the true model. In other words, the model has been misspecified. In this paper, we consider the prediction interval problem in the case of a misspecified first-order autoregressive or AR(1) model. We have calculated the coverage probability of an upper one-step-ahead prediction interval for both properly specified and misspecified models through Monte Carlo simulation. It was found that dealing with prediction interval for misspecified model is complicated: the distribution of a future observation conditional on the last observation and the parameter estimator is not identical to the distribution of this future observation conditional on the last observation alone.

Keywords: *autoregressive; coverage probability; model misspecification; time series prediction.*

1 Introduction

A very informative way of specifying the accuracy of a time series prediction is to use a prediction interval. To do this, some authors, e.g. [1-5], have computed the prediction interval or limit for the autoregressive (AR) process and the autoregressive conditional heteroscedastic (ARCH) process. For these models, they have assumed that the fitted model is the true model. In other words, the authors have employed properly specified models.

In practice, however, time series models are likely to be misspecified. For instance, an AR(1) model is fitted but the true model is an AR(2) or MA(1) model. This paper deals with the prediction interval in the case of a misspecified AR(1) model. In the context of the prediction problem, there have been a few papers, such as [6] and [7], that provide derivations of expressions for mean-squared-error of prediction in the case of a misspecified Gaussian AR(p) model. An investigation of the coverage probability, conditional on the last observations, of a prediction interval based on a misspecified Gaussian AR(p) model was presented first by [8].

In the case of a misspecified Gaussian AR(p) model, the derivation of the asymptotic expansion of the coverage probability, conditional on the last

observations, of a prediction interval requires a great deal of care. To make clear the issues involved in this derivation, we consider the simple scenario described in Section 2. Namely, a misspecified zero-mean Gaussian AR(1) process fitted to a stationary zero-mean Gaussian process. We also consider a one-step-ahead prediction interval with target coverage probability. Similarly to [8], we aim to find the asymptotic expansion for the coverage probability of the prediction interval under consideration, conditional on the last observation.

In Section 3, we compute the conditional coverage probabilities for both properly and misspecified first order autoregressive models. This is aimed to examine the effect of misspecification on the coverage probability of the prediction interval. Furthermore, the technical argument for finding the conditional coverage probability of a prediction interval as proposed by de Luna (which differs from the estimative prediction interval) is reviewed in detail in Section 4. A numerical example to illustrate a gap in his argument is given in Section 5.

2 Description of the Time Series and Fitted Models

Consider a discrete time stationary Gaussian zero-mean process $\{Y_t\}$. Let $\gamma_k = E(Y_t Y_{t-k})$ denote the autocovariance function of $\{Y_t\}$ evaluated at lag k . Suppose that $\sum_{j=1}^{\infty} |\gamma_j| < \infty$. We do not require that $\{Y_t\}$ is an AR(1) process. The available data are Y_1, Y_2, \dots, Y_n . We will be careful to distinguish between random variables, written in upper case, and their observed values, written in lower case.

We consider one-step-ahead prediction using a stationary Gaussian zero-mean AR(1) model. This model may be misspecified. Let $\rho_k = \gamma_k / \gamma_0$. Define θ to be the value of a that minimizes $E(Y_{n+1} - aY_n)^2$. Thus $\theta = \gamma_1 / \gamma_0$. Define $\psi^2 = \text{var}(Y_{n+1} | Y_n = y_n)$. Note that

$$\psi^2 = \gamma_0 - \frac{\gamma_1^2}{\gamma_0}$$

and $E(Y_{n+1} - \theta Y_n)^2 = \psi^2$. Let $\widehat{\Gamma}_k$ denote a consistent estimator of γ_k such as $n^{-1} \sum_{j=k+1}^n Y_j Y_{j-k}$. Let $\widehat{\Theta} = \widehat{\Gamma}_1 / \widehat{\Gamma}_0$ and note that $\widehat{\Theta}$ is a consistent

estimator of θ . We use the notation $[a \pm b]$ for the interval $[a - b, a + b]$ ($b > 0$).

3 The Coverage Properties of Misspecified AR(1) Model

In this section, we examine the effect of misspecification on the coverage probability of the prediction interval. The true model $\{Y_t\}$ is a zero-mean Gaussian AR(2) process, satisfying

$$Y_t = \theta_1 Y_{t-1} + \theta_2 Y_{t-2} + \eta_t \quad (1)$$

for all integer t , where η_t are independent and identically $N(0, \sigma^2)$ distributed. The roots of $1 - \theta_1 m - \theta_2 m^2$ are outside the unit circle to ensure stationarity of the process.

We consider an upper one-step-ahead prediction interval with conditional mean square prediction error equal to V^2 , i.e.

$$I_u(Y_n; \hat{\Theta}, \hat{V}^2) = \hat{\Theta} Y_n + \Phi^{-1}(1 - \alpha) \hat{V}, \quad (2)$$

where the estimators $\hat{\Theta}$ and \hat{V}^2 are obtained by maximizing the loglikelihood function conditional on $Y_1 = y_1$, that are

$$\hat{\Theta} = \frac{\sum_{t=1}^{n-1} Y_t Y_{t+1}}{\sum_{t=1}^{n-1} Y_{t+1}^2}$$

and

$$\hat{V}^2 = \frac{1}{n-1} \sum_{t=1}^{n-1} (Y_t - \hat{\Theta} Y_{t+1})^2.$$

The coverage probability of $I_u(Y_n; \hat{\Theta}, \hat{V}^2)$, conditional on $Y_n = y_n$ is calculated as follows. First, we observe that

$$\begin{aligned}
& P(Y_{n+1} \leq \widehat{\Theta} Y_n + \Phi^{-1}(1-\alpha) \widehat{V} | Y_n = y_n, Y_{n-1} = y_{n-1}) \\
&= P(\theta_1 Y_n + \theta_2 Y_{n-1} + \eta_{n+1} \leq \widehat{\Theta} Y_n + \Phi^{-1}(1-\alpha) \widehat{V} | Y_n = y_n, Y_{n-1} = y_{n-1}) \\
&= P(\eta_{n+1} \leq (\widehat{\Theta} - \theta_1) y_n - \theta_2 y_{n-1} + \Phi^{-1}(1-\alpha) \widehat{V} | Y_n = y_n, Y_{n-1} = y_{n-1}) \\
&= E \left(\Phi \left((\widehat{\Theta} - \theta_1) \frac{y_n}{\sigma} - \theta_2 \frac{y_{n-1}}{\sigma} + \Phi^{-1}(1-\alpha) \frac{\widehat{V}}{\sigma} \right) | Y_n = y_n, Y_{n-1} = y_{n-1} \right) \quad (3)
\end{aligned}$$

and we estimate this conditional expectation by simulation. We use the backward representation:

$$Y_t = \theta_1 Y_{t+1} + \theta_2 Y_{t+2} + v_t \quad (4)$$

for all integer t , where v_t are independent and identically $N(0, \sigma^2)$ distributed. It is noted that v_t and Y_{t+1}, Y_{t+2}, \dots are independent. We begin the simulation run by setting $(Y_n, Y_{n-1}) = (y_n, y_{n-1})$ and then use (4) to run the process backwards in time.

Table 1 Estimated coverage probabilities, conditional on $(Y_n, Y_{n-1}) = (0, 1)$, of the upper 0.95 one-step-ahead forecasting intervals for true and misspecified models. Standard errors are in brackets.

a_1, a_2, σ^2	Model		
$a_1 = 1, a_2 = 0.25, \sigma^2 = 1$	True	0.9401 (0.00079)	0.9451 (0.00055)
	Misspecified	0.9999 (0.00001)	1 (0.000002)
$a_1 = 0.1, a_2 = -0.4, \sigma^2 = 1$	True	0.9327 (0.00091)	0.9423 (0.00056)
	Misspecified	0.9311 (0.0011)	0.9316 (0.0009)
$a_1 = -0.3, a_2 = -0.3, \sigma^2 = 1$	True	0.9342 (0.00088)	0.9419 (0.00057)
	Misspecified	0.9656 (0.0010)	0.9728 (0.0007)
$a_1 = -0.75, a_2 = 0.5, \sigma^2 = 1$	True	0.9427 (0.00074)	0.9468 (0.00049)
	Misspecified	0.9988 (0.00006)	0.9993 (0.00002)

Table 1 presents the estimate of conditional coverage probability (3). Standard errors of these estimates are given in brackets. We also provide the estimate of conditional coverage probability of when the predictor is also a zero-mean Gaussian AR(2) model (properly specified model) as in [4].

4 The Conditional Distribution for Misspecified Model

de Luna (2000) considers the data generating mechanism $\{Y_t\}$, as described earlier, and h -step-ahead prediction using a zero-mean Gaussian AR(p) model. This model may be misspecified. Consider the case that $h=1$ and $p=1$. His $1-\alpha$ prediction interval for Y_{n+1} is

$$J = [\hat{\Theta}Y_n - \sqrt{\psi^2 + \omega^2(Y_n)} z_{1-\alpha/2}, \hat{\Theta}Y_n + \sqrt{\psi^2 + \omega^2(Y_n)} z_{1-\alpha/2}],$$

where $\omega^2(Y_n) = n^{-1} w Y_n^2$, $w = \sum_{k=1}^{\infty} (\rho_{k+1} + \rho_{k-1} - 2\rho_1\rho_k)^2$ and $z_{1-\alpha/2}$ is the $(1-\alpha/2)$ quantile of the $N(0,1)$ distribution. In practice, ψ^2 and w are unknown since they depend on an unknown autocovariance function of the process $\{Y_t\}$. However, de Luna assumes that both ψ^2 and w are known.

de Luna claims that $P(Y_{n+1} \in J | Y_n = y_n) = 1 - \alpha + O(n^{-1})$. His justification for this claim, with intermediate steps added, is the following.

$$\begin{aligned} & P(Y_{n+1} \in J | Y_n = y_n) \\ &= P(Y_{n+1} \in [\hat{\Theta}y_n \pm \sqrt{\psi^2 + \omega^2(y_n)} z_{1-\alpha/2}] | Y_n = y_n) \\ &\quad \text{(by the substitution theorem for conditional expectations)} \\ &= E(P(Y_{n+1} \in [\hat{\Theta}y_n \pm \sqrt{\psi^2 + \omega^2(y_n)} z_{1-\alpha/2}] | Y_n = y_n, \hat{\Theta}) | Y_n = y_n), \end{aligned}$$

by the double expectation theorem. To evaluate this conditional expectation, de Luna proceeds as follows. Define $F(\cdot)$ and $f(\cdot)$ to be the cdf and pdf of Y_{n+1} conditional on $Y_n = y_n$, respectively. He then assumes that

$$P(Y_{n+1} \in [\hat{\Theta}y_n \pm \sqrt{\psi^2 + \omega^2(y_n)} z_{1-\alpha/2}] | Y_n = y_n, \hat{\Theta})$$

is equal to

$$F(\widehat{\Theta}y_n + \sqrt{\psi^2 + \omega^2(y_n)}z_{1-\alpha/2}) - F(\widehat{\Theta}y_n - \sqrt{\psi^2 + \omega^2(y_n)}z_{1-\alpha/2}), \quad (5)$$

since the Taylor expansions are applied to F and not some other function. Then, by Taylor expansion of F around $\theta y_n + \psi z_{1-\alpha/2}$,

$$\begin{aligned} & F(\widehat{\Theta}y_n + \sqrt{\psi^2 + \omega^2(y_n)}z_{1-\alpha/2}) \\ &= F(\theta y_n + \psi z_{1-\alpha/2}) \\ &+ f(\theta y_n + \psi z_{1-\alpha/2})(y_n(\widehat{\Theta} - \theta) + (\sqrt{\psi^2 + \omega^2(y_n)} - \psi)z_{1-\alpha/2}) + \dots \end{aligned}$$

Now, $F(\theta y_n + \psi z_{1-\alpha/2}) = 1 - \alpha/2$. Thus

$$E(F(\widehat{\Theta}y_n + \sqrt{\psi^2 + \omega^2(y_n)}z_{1-\alpha/2}) | Y_n = y_n)$$

is equal to

$$1 - \frac{\alpha}{2} + f(\theta y_n + \psi z_{1-\alpha/2})(y_n E(\widehat{\Theta} - \theta | Y_n = y_n) + (\sqrt{\psi^2 + \omega^2(y_n)} - \psi)z_{1-\alpha/2}) + \dots$$

Similarly,

$$\begin{aligned} & E(F(\widehat{\Theta}y_n - \sqrt{\psi^2 + \omega^2(y_n)}z_{1-\alpha/2}) | Y_n = y_n) \\ &= \frac{\alpha}{2} + f(\theta y_n - \psi z_{1-\alpha/2})(y_n E(\widehat{\Theta} - \theta | Y_n = y_n) - (\sqrt{\psi^2 + \omega^2(y_n)} - \psi)z_{1-\alpha/2}) + \dots \end{aligned}$$

If it is assumed that $E(\widehat{\Theta} - \theta | Y_n = y_n) = O(n^{-1})$ then $P(X_{n+1} \in J | Y_n = y_n) = 1 - \alpha + O(n^{-1})$.

The assumption (5) is crucial. It is true if $\{Y_t\}$ is also an AR(1) process (i.e. if the model is properly specified), but it is not necessarily true if the model is misspecified. Let $F^\dagger(\cdot; \hat{\theta})$ denote the cdf of Y_{n+1} conditional on $Y_n = y_n$ and

$\widehat{\Theta} = \hat{\theta}$. Now

$$\begin{aligned}
P(Y_{n+1} \in [\hat{\Theta} y_n \pm \sqrt{\psi^2 + \omega^2(y_n)} z_{1-\alpha/2}] | Y_n = y_n, \hat{\Theta}) \\
= F^\dagger(\hat{\Theta} y_n + \sqrt{\psi^2 + \omega^2(y_n)} z_{1-\alpha/2}; \hat{\Theta}) - F^\dagger(\hat{\Theta} y_n - \sqrt{\psi^2 + \omega^2(y_n)} z_{1-\alpha/2}; \hat{\Theta}).
\end{aligned}$$

If $\{Y_t\}$ is also an AR(1) process then, by the Markov property, $F^\dagger(\cdot; \hat{\theta}) = F(\cdot)$ and so (5) is satisfied. If, however, the model is misspecified then $F^\dagger(\cdot; \hat{\theta})$ is not necessarily equal to $F(\cdot)$.

5 Computational Example to Illustrate the Error in the Calculating Conditional Distribution

In this section, we provide a numerical example to support the following argument. For a misspecified AR(1) model, the distribution of Y_{n+1} conditional on $Y_n = y_n$ is not identical to the distribution of Y_{n+1} conditional on $Y_n = y_n$ and $\hat{\Theta} = \hat{\theta}$. In doing so, we seek an estimator $\hat{\Theta}$, an observed value $\hat{\theta}$ and a misspecified AR(1) model such that the condition $\hat{\Theta} = \hat{\theta}$, when added to the condition $Y_n = y_n$, tells us a great deal more about the value of Y_{n+1} than the condition $Y_n = y_n$ alone.

Consider the following estimator:

$$\hat{\Theta} = \frac{\left(\sum_{t=2}^T Y_t Y_{t-1} \right)^2}{\sum_{t=2}^T Y_t^2 \sum_{t=2}^T Y_{t-1}^2} \quad (6)$$

and let $(y_1, \dots, y_n) = (Y_1, \dots, Y_n)$ and $(x_1, \dots, x_n) = (Y_1, \dots, Y_{n-1})$. Thus, we write the estimator (6) as

$$\hat{\Theta} = \frac{\left(\sum_{i=1}^n y_i x_i \right)^2}{\sum_{i=1}^n y_i^2 \sum_{j=1}^n x_j^2}$$

which is less than or equal to 1, according to the Cauchy-Schwarz inequality.

Equality occurs if $x_i = a^* y_i$, i.e. $Y_t = a^* Y_{t+1}$, where

$$a^* = \frac{\sum_{i=1}^n y_i x_i}{\sum_{i=1}^n y_i^2} = \frac{\sum_{t=2}^T Y_t Y_{t-1}}{\sum_{t=2}^T Y_t^2}$$

Thus,

$$\widehat{\Theta} = 1 \Rightarrow Y_t = a^* Y_{t+1}$$

where $a^* \approx \widehat{\Theta} = 1$. Meanwhile,

$$\widehat{\Theta} \approx 1 \text{ and } Y_n = y_n \Rightarrow Y_{n-1} \approx y_n.$$

Now, consider a stationary zero-mean Gaussian AR(1) process $\{Y_t\}$ satisfying

$$Y_t = \beta Y_{t-2} + \varepsilon_t$$

where ε_t are independent and identically $N(0, \sigma^2)$ distributed, β is close to 1 but satisfies $\beta < 1$. We find the autocovariance properties of $\{Y_t\}$ as follows:

$$\gamma_0 = \frac{\sigma^2}{1 - \beta^2}$$

$$\gamma_{2k} = \frac{\beta^k \sigma^2}{1 - \beta^2}, k = 1, 2, \dots$$

whilst $\gamma_l = 0$, for $l = 1, 3, \dots$. By the properties above, $\theta = \gamma_1 / \gamma_0 = 0$ for the above process. Thus the distribution of Y_{n+1} conditional on $Y_n = y_n$ is $N(0, \gamma_0)$, i.e.

$$(Y_{n+1} | Y_n = y_n) \sim N(0, \gamma_0) = N\left(0, \frac{\sigma^2}{1 - \beta^2}\right). \quad (7)$$

If we choose $\sigma^2 = 1$, $\beta = 0.95$ and y_n fairly large,

$$y_n = 2\sqrt{\frac{1}{1-0.95^2}} = 6.405, \text{ say,}$$

then

$$(Y_{n+1} | Y_n = 6.405) \sim N(0, 1/(1-0.95^2)) = N(0, 10.256).$$

Now, if we condition on $|\widehat{\Theta} - 1| < \delta$, where $\delta = 0.01$, say, and n is not too large then we expect that, to a good approximation,

$$(Y_{n+1} | Y_n = y_n, |\widehat{\Theta} - 1| < \delta) \sim N(\beta y_n, \sigma^2) = N(0.95 y_n, 1). \quad (8)$$

which is quite different from (7).

As for illustration, a simulation is carried out as follows. Initialize the counter: set $l = 0$. The k th simulation run is

- simulate an observation of (Y_1, \dots, Y_{n-1}) conditional on $Y_n = y_n$.
- calculate $\widehat{\Theta}$
- if $|\widehat{\Theta} - 1| < \delta$, then $l = l + 1$ and store $x_l = y_{n-1}$

After M simulation runs, suppose that $l = N$. The probability density function of Y_{n+1} conditional on $Y_n = y_n$ and $|\widehat{\Theta} - 1| < \delta$ is approximated by

$$\frac{1}{N} \sum_{l=1}^N f(y; 0.95 x_l)$$

where $f(y; 0.95 x_l)$ denotes the $N(0.95 x_l, 1)$ probability density function evaluated at y . Note that to simulate an observation of (Y_1, \dots, Y_{n-1}) conditional on $Y_n = y_n$




Figure 1 The pdf of Y_{n+1} conditional on $Y_n = y_n$ (top) and the pdf of Y_{n+1} conditional on $Y_n = y_n$ and $|\hat{\Theta} - 1| < \delta$ (bottom), $M = 10.000$.

we shall know that the vector $Y = (Y_1, \dots, Y_n)^T$ follows a multivariate normal distribution with mean vector μ and the covariance matrix Σ i.e.

$$\begin{pmatrix} Y_1 \\ \vdots \\ Y_{n-1} \\ Y_n \end{pmatrix} \sim N \left(\begin{bmatrix} 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \gamma_0 & \gamma_1 & \cdots & \gamma_{n-1} \\ \gamma_1 & \gamma_0 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \gamma_1 \\ \gamma_{n-1} & \cdots & \gamma_1 & \gamma_0 \end{bmatrix} \right)$$

Now if we partition the mean vector μ and the covariance matrix Σ such that

$$\mu = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}$$

with sizes for μ_1 and μ_2 are $(n-1) \times 1$ and 1×1 , respectively, and

$$\Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix}$$

with sizes

$$\begin{pmatrix} (n-1) \times (n-1) & (n-1) \times 1 \\ 1 \times (n-1) & 1 \times 1 \end{pmatrix}$$




Figure 2 The pdf of Y_{n+1} conditional on $Y_n = y_n$ (top) and the pdf of Y_{n+1} conditional on $Y_n = y_n$ and $|\hat{\Theta} - 1| < \delta$ (bottom), $M=50.000$.

Then the distribution of (Y_1, \dots, Y_{n-1}) conditional on $Y_n = y_n$ is multivariate normal with mean vector $\widetilde{\mu}$ and covariance matrix $\widetilde{\Sigma}$ ([9]), i.e.

$$(Y_1, \dots, Y_{n-1} | Y_n = y_n) \sim N(\widetilde{\mu}, \widetilde{\Sigma})$$

where the mean vector and the covariance matrix

$$\widetilde{\mu} = \mu_1 + \Sigma_{12}\Sigma_{22}^{-1}(y_n - \mu_2)$$

$$\widetilde{\Sigma} = \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21}$$

Figures 1-3 show the pdf of Y_{n+1} conditional only on $Y_n = y_n$ (top figure) and the pdf of Y_{n+1} conditional on $Y_n = y_n$ and $|\widehat{\Theta} - 1| < \delta$ (bottom figure). We carry out $n = 300$ and $M = 10,000, 50,000, \text{ and } 100,000$ simulation runs.

6 Discussion

The derivation of the asymptotic conditional coverage probability for a misspecified model requires a great deal more care than the derivation of the asymptotic conditional coverage probability for a properly specified model. Results similar to those of de Luna can be derived by defining the random variable $\varepsilon_{n+1} = X_{n+1} - E(X_{n+1}|X_n, X_{n-1}, \dots)$ and noting that ε_{n+1} and (X_n, X_{n-1}, \dots) are independent random vectors.




Figure 3 The pdf of Y_{n+1} conditional on $Y_n = y_n$ (top) and the pdf of Y_{n+1} conditional on $Y_n = y_n$ and $|\widehat{\Theta} - 1| < \delta$ (bottom), $M=100,000$.

$X_{n+1} - E(X_{n+1}|X_n, X_{n-1}, \dots)$ and noting that ε_{n+1} and (X_n, X_{n-1}, \dots) are independent random vectors.

Acknowledgement

I am grateful to Paul Kabaila for his valuable comments and suggestions to clarify many aspects on this paper.

References

- [1] Barndorff-Nielsen, O.E. & Cox, D.R., *Prediction and Asymptotics*, Bernoulli, **2**, pp. 319-340, 1996.
- [2] Kabaila, P., *The Relevance Property for Prediction Intervals*, Journal of Time Series Analysis, **20**(6), pp. 655-662, 1999.
- [3] Kabaila, P. & Syuhada, K., *The Relative Efficiency of Prediction Intervals*, Communication in Statistics: Theory and Methods, **36**(15), pp. 2673-2686, 2007.
- [4] Kabaila, P. & Syuhada, K., *Improved Prediction Limits for AR(p) and ARCH(p) Processes*, Journal of Time Series Analysis, **29**, pp. 213-223, 2008.
- [5] Vidoni, P., *Improved Prediction Intervals for Stochastic Process Models*, Journal of Time Series Analysis, **25**, pp. 137-154, 2004.
- [6] Davies, N. & Newbold, P., *Forecasting with Misspecified Models*, Applied Statistics, **29**, pp. 87-92, 1980.
- [7] Kunitomo, N. & Yamamoto, T., *Properties of Predictors in Misspecified Autoregressive Time Series Models*, Journal of the American Statistical Association, **80**, pp. 941-950, 1985.
- [8] de Luna, X., *Prediction Intervals Based on Autoregression Forecasts*, The Statistician, **49**, pp. 87-93, 2000.
- [9] Graybill, F.A., *Theory and Application of The Linear Model*, North Scituate, Mass: Duxbury Press, 1976.