

ON COVERS OF DIHEDRAL 2-GROUPS BY POWERFUL SUBGROUPS

RISTO ATANASOV, ADAM GREGORY, LUKE GUATELLI, AND ANDREW PENLAND

ABSTRACT. A finite p -group G is called *powerful* if either p is odd and $[G, G] \subseteq G^p$ or $p = 2$ and $[G, G] \subseteq G^4$. A *cover* for a group is a collection of subgroups whose union is equal to the entire group. We will discuss covers of p -groups by powerful subgroups. The size of the smallest cover of a p -group by powerful subgroups is called the *powerful covering number*. Our focus in this paper is to determine the powerful subgroup covering number of the Dihedral 2-groups.

1. INTRODUCTION

If G is a group, a *cover* of G is a collection of proper subgroups whose union is equal to G , and the *covering number* of G is the minimal number of proper subgroups that occur in any cover of G . We write $\sigma(G)$ for the covering number of G .

If G is a p -group, we define a *powerful cover* of G to be a covering of G by powerful subgroups. We define the *powerful covering number* of G to be the minimal number of subgroups in any powerful covering of G . Our main result in this paper is to establish the powerful covering number of the dihedral 2-groups.

Interest in coverings and covering numbers goes back to G.A. Miller, who considered covers by subgroups with pairwise trivial intersection []. Such a cover is known as a *partition*. reference and culminating in a theorem which completely classifies all finite groups with a partition. todo: references see https://projecteuclid.org/download/pdf_1/euclid.ijm/1258488174 todo: Partition Numbers by Foguel

Scorza todo:reference, who first noted that there is no group with $\sigma(G) = 2$. There has been a lot of interest in calculating the covering number for todo: more references for $\sigma(G)$ – Cohn, Tomkinson, etc.

Additionally, one may consider coverings in which all subgroups share some certain property. Bargava (todo: reference) classified the groups which have coverings by normal subgroups. todo: look also at <https://mathoverflow.net/questions/266368/normal-covering-number-and-maximal-cyclic-subgroups>

Foguel and Ragland (todo: reference) investigated coverings by abelian groups, all of which are pairwise isomorphic. Atanasov, Foguel, and Penland considered coverings by subgroups that all have the same order and have mutually isomorphic pairwise intersections.

2. BACKGROUND

In the interest of making this paper self-contained and accessible to non-experts, we will make very few assumptions about the reader's knowledge of group theory. This section reviews all definitions and well-known facts necessary to establish our results. Most of this material is standard and can be found in a standard text such as [1].

Henceforth in this paper, all groups are assumed to be finite.

2.1. Group Theory. Let G be a group. If S is a subset of G , the *group generated by S* is the smallest subgroup of G that contains S . If T is a set, we write $|T|$ for the cardinality of T . A subgroup generated by a single element $g \in G$ is denoted $\langle g \rangle$ in an abuse of notation. Such a subgroup is called *cyclic*. For $g \in G$, the *order* of g is $|\langle g \rangle|$. For g, h in G , the *commutator of g and h* is defined as $g^{-1}h^{-1}gh$. The *commutator subgroup of G* , denoted $[G, G]$, is the group generated by all commutators of all pairs of elements in G . Let p be a fixed prime number. We say that G is a p -group if every element in G has order equal to a power of p . The subgroup G^p is defined as

$$G^p = \langle \{g^p \mid g \in G\} \rangle.$$

Date: November 2018.

Key words and phrases.

A subgroup K of G is *maximal* if whenever there exists a subgroup $H \subseteq G$ such that K is a proper subgroup of H , it follows that $H = G$. If H is any subgroup of G , the *index* of H in G is equal to $\frac{|G|}{|H|}$. For a p -group, it is known that all maximal subgroups have index p . If T is a group, a homomorphism is a map $\alpha : G \rightarrow T$ such that $\alpha(gh) = \alpha(g)\alpha(h)$ for all g, h in G . If T is a group and $\alpha : G \rightarrow T$ is a homomorphism, then the index of $\ker \alpha$ is equal to the size of the image of α .

If G_1 and H are groups, then the *direct product* $G_1 \times G_2$ is a group with the set $\{(g, h) \mid g \in G, h \in H\}$ and group operation given componentwise.

2.2. Powerful Groups. A finite p -group G is called *powerful* if either p is odd and $[G, G] \subseteq G^p$ or $p = 2$ and $[G, G] \subseteq G^4$. Powerful groups were first introduced by todo: whoever introduced them.

todo: references for places where powerful groups have been used to obtain important results (such as classification of groups by finite coclass) todo:

2.3. Group Covers. If G is a group, the *covering number* of G is defined as the minimal number of subgroups needed to cover the group. We define the *powerful covering number* of G as the minimal number of proper powerful subgroups needed to cover the group. We define the *abelian covering number* of G as the minimal number of abelian powerful subgroups needed to cover the G . We write $\sigma(G)$ for the covering number, $\sigma_P(G)$ for the powerful covering number, and $\sigma_A(G)$ for the abelian covering number.

Lemma 2.1. *Let G be a finite noncyclic p -group. Then G has a powerful covering.*

Proof. G has a covering by the set of all cyclic subgroups, and cyclic groups are powerful. □

Lemma 2.2. *If G is a finite p -group, the relationship*

$$\sigma_A(G) \leq \sigma_P(G) \leq \sigma(G)$$

always holds.

Proof. This follows immediately from the fact that abelian groups are powerful. □

Theorem 2.3. *Let G be a finite p -group. If K is a finite homomorphic image of G such that K is not cyclic, then $\sigma_P(K) \leq \sigma_P(G)$.*

Note (AP): I believe that the following claim is true, but it needs verification. Be careful! It is not true for covering numbers in general. Understanding why would be a good exercise.

Proof. Suppose $G = \bigcup_{i=1}^n H_i$, where each H_i is a proper powerful subgroup. Let $\alpha : G \rightarrow K$ be a surjective homomorphism. We see that each $\alpha(H_i)$ is also a powerful subgroup, and that

$$\bigcup_{i=1}^n \alpha(H_i) = K.$$

□

Theorem 2.4. *Let G be a finite p -group and let K be a powerful finite p -group. Then $\sigma_P(G \times K) = \sigma_P(G)$.*

Proof. □

todo: add reference to Cohn or Tomkinson for the theorem below.

Theorem 2.5. *Let G be a p -group. Then $\sigma(G) = p + 1$.*

This immediately leads to the following result for abelian p -groups.

Corollary 2.6. *Let G be an abelian p -group. Then $\sigma_A(G) = \sigma_P(G) = \sigma(G) = p + 1$.*

The covering number σ is well-behaved with respect to subgroup inclusion: if H is a subgroup of G , then $\sigma(H) \leq \sigma(G)$. However, for powerful covers, this may not be true, since not all subgroups of a powerful group are powerful.

todo: example of a subgroup where the powerful covering number is larger than the powerful covering number of a group?

2.4. Cyclic and Dihedral Groups. We define the *cyclic group of order t* to be the group generated by a single element of order t . We write C_t for the cyclic group of order t . We let σ represent the generator of C_2 .

The following facts about

For $n \geq 2$, we define $D(2^n)$ to be the set of isometries of a regular 2^n -gon. The group $D(2^n)$ has 2^{n+1} elements.

We say a group is a *dihedral 2-group* if it is equal to $D(2^n)$ for some $n \geq 3$, or if it is isomorphic to C_2 , $C_2 \times C_2$, or the trivial group.

3. ELEMENTS AND SUBGROUPS OF DIHEDRAL GROUPS

todo: We need to define the exact generators a and b are two elements of order two so that ab is the rotation by $\frac{2\pi}{2^n}$ radians.

Lemma 3.1. Suppose $g \in D(2^n)$.

- (i.) The element g can be written uniquely as $g = (ab)^j a^k$ for some integers j and k with $0 \leq j < 2^n$ and $0 \leq k \leq 1$.
- (ii.) If $g = (ab)^j a$ for some j with $0 \leq j < 2^n$, then $|g| = 2$.
- (iii.) If $g = (ab)^j$ for some j with $0 \leq j < 2^n$, then $|g| = \frac{2^{n-1}}{\gcd(j, 2^{n-1})}$.

Proof. Proof of (i.) This follows from well-known facts about cosets in a group. (todo: Provide lemma above.) (Proof of (ii.)) Suppose $g = (ab)^j a$ for some j with $0 \leq j \leq 2^{n-1}$. Then we calculate

$$\begin{aligned} ((ab)^j a)^2 &= (ab)^j a (ab)^j a \\ &= a (ba)^j (ab)^j a \end{aligned}$$

Since $(ab)^j$ is the inverse of $(ba)^j$ and $a^2 = e$, $a(ba)^j (ab)^j a$ is trivial, and the result follows. (Proof of (iii.)) This follows from the fact that (ab) has order 2^{n-1} and Lemma ??.

Lemma 3.2. Let $n \geq 2$. For the element $ab \in D(2^n)$, the following properties hold.

- (i.) ab has order 2^n .
- (ii.) For any r , $(ab)^r a = a(ab)^{-r}$.
- (iii.) $(ab)^{2^{n-1}} = (ba)^{2^{n-1}}$.
- (iv.) For any $g \in D(2^n)$, $g(ab)^{2^{n-1}} = (ab)^{2^{n-1}} g$.

Proof. The order of ab follows directly from the fact that ab represents a rotation of $\frac{2\pi}{2^n}$ radians. For (ii.), notice that that $(ab)(ba) = e$, know that $ba = (ab)^{-1}$ and hence represents a rotation of $-\frac{2\pi}{2^n}$ radians. Since $(ab)^r a$ represents a reflection, it has order two, and thus $(ab)^r a$ is equal to its own inverse, which is equivalent to $a(ab)^{-r}$. Since a rotation of $2^{n-1} \frac{2\pi}{2^n} = \pi$ radians is the same as a rotation of $2^{n-1} \cdot -\frac{2\pi}{2^n} = -\pi$ radians, we have that $(ab)^{2^{n-1}} = (ba)^{2^{n-1}}$. To see (iv.), let $g \in D(2^n)$ and write $g = (ab)^j a^k$ as in Lemma 3.1(i.). If $k = 0$, then the statement holds since powers of ab commute. If $k = 1$, then we have

$$\begin{aligned} (ab)^{2^{n-1}} g &= (ab)^{2^{n-1}} (ab)^j a \\ &= (ab)^{2^{n-1}+j} a \\ &= (ab)^j (ab)^{2^{n-1}} a \\ &= (ab)^j a (ba)^{2^{n-1}} \\ &= (ab)^j a (ab)^{2^{n-1}} \\ &= g(ab)^{2^{n-1}}. \end{aligned}$$

□

Lemma 3.3. *If G is a finite abelian group generated by two elements of order two, then G is isomorphic to $C_2 \times C_2$.*

Proof. □

Lemma 3.4. *If G is a finite group generated by two elements of order two, then G is isomorphic to a dihedral 2-group.*

Proof. todo □

Proposition 3.5. *For any $n \geq 1$, $D(2^n)$ has three maximal subgroups. Two of these maximal subgroups are isomorphic to $D(2^{n-1})$, and one is isomorphic to C_{2^n} .*

Proof. The maximal subgroups of a dihedral 2-group $D(2^n)$ correspond to the kernels of nontrivial homomorphisms from $D(2^n)$ to C_2 . Such a homomorphism is completely determined by the images of the generators a and b . Let α_1 be the homomorphism for which $\alpha_1(a)$ is trivial and $\alpha_1(b)$ is not. Let $H_1 = \ker \alpha_1$. It is not hard to see that this group consists of all elements with an even number of b 's, and therefore is generated by a and aba . Since a and aba both have order two, H_1 is isomorphic to a finite dihedral group by Lemma 3.4. We know the order of H_1 is 2^n , so H_1 is isomorphic to $D(2^{n-1})$. Let α_2 be the homomorphism for which $\alpha_2(b)$ is trivial and $\alpha_2(a)$ is not. Letting $H_2 = \ker \alpha_2$, an identical argument to that of α_1 establishes that $H_2 \cong D(2^{n-1})$. Finally, let α_3 be the homomorphism such that neither $\alpha_3(a)$ nor $\alpha_3(b)$ is trivial. Let $H_3 = \ker \alpha_3$. It is clear that H_3 contains ab , so H_3 contains the cyclic subgroup $\langle ab \rangle$, which has order 2^n by Lemma 3.2. Since H_3 is maximal, it has order 2^n , so $H_3 = \langle ab \rangle \cong C_{2^n}$. □

Proposition 3.6. *Every subgroup of a dihedral 2-group $D(2^n)$ is either dihedral or cyclic.*

Proof. We will proceed by induction. For the base case $n = 2$, this fact is easy to see for $D(4)$. Now assume the statement is true for some $k \geq 2$, and consider $D(2^{k+1})$. By Proposition 3.5, the maximal subgroups of this group are either dihedral or cyclic. Then, by the induction hypothesis and the fact that every proper subgroup is contained in some maximal subgroup, the desired result follows. □

Corollary 3.7. *If a subgroup H of $D(2^n)$ is powerful, then either $H \cong C_2 \times C_2$ or $H \cong C_k$ where k is a divisor of 2^{n-1} . If $H \cong C_k$ for $k > 2$, then H is a subgroup of $\langle ab \rangle$.*

Proof. The first statement follows from Proposition 3.6. The second statement follows from Lemma 3.1. □

Note that the form for elements given in Lemma 3.1 may not be the shortest or most intuitive form for writing the element. For instance, the generator $b \in D_4$ would be written as $(ab)^3a$ under the convention we've established. Lemma 3.1 also establishes the following characterization of elements of order two in $D(2^n)$.

Corollary 3.8. *Let $G = D_{2^n}$. If $g \in G$, then g has order two if and only if $g = (ab)^j a$ for some $0 \leq k \leq 2^n$, or $g = (ab)^{2^n}$.*

Proposition 3.9. *Let $\mathcal{C} = \{H_1, \dots, H_q\}$ be any subgroup cover of $D(2^n)$. Then there is some i with $1 \leq i \leq q$ such that $H_i = \langle ab \rangle$.*

Proof. By definition of subgroup cover, there must be some $i \in \{1, \dots, q\}$ such that $ab \in H_i$. Since H_i contains ab , it follows that H_i contains the subgroup $\langle ab \rangle$. Since $\langle ab \rangle$ is maximal and H_i must be a proper subgroup, it follows that $H_i = \langle ab \rangle$. □

Note to Risto: I was able to do the proof without what we were calling “Lemma 2” in your office.

Lemma 3.10. *Let H be a subgroup of a dihedral 2-group. Then H is isomorphic to $C_2 \times C_2$ if and only if $H = \langle (ab)^s a, (ab)^t a \rangle$, where $t \neq s$ and $t + s = 2^{n-1}$.*

Proof. Let t and s be distinct positive integers $s + t = 2^{n-1}$. We let $x = (ab)^s a$ and $y = (ab)^t a$, and we let H be a group generated by x and y . First, notice that x and y each have order two, by Lemma ???. We

calculate that

$$\begin{aligned}
xy &= (ab)^s a(ab)^t a \\
&= (ab)^s a(ab)^{2^{n-1}-s} a \\
&= (ab)^s aa(ba)^{2^{n-1}-s} \\
&= (ab)^s aa(ab)^{2^{n-1}-s} \\
&= (ab)^s (ab)^{2^{n-1}-s} \\
&= (ab)^{2^{n-1}}.
\end{aligned}$$

A similar calculation establishes that $yx = (ba)^{2^{n-1}}$, and since $(ab)^{2^{n-1}} = (ba)^{2^{n-1}}$ by Lemma 3.2, it follows that $xy = yx$. Hence $\langle x, y \rangle \cong C_2 \times C_2$ by Lemma 3.3.

For the other direction, we suppose that H is a subgroup of $D(2^n)$ such that H is isomorphic to $C_2 \times C_2$. Note that H must contain an element of the form $(ab)^s a$, otherwise H would be contained in the cyclic group $\langle ab \rangle$. H must also contain another element g of order two. We consider two cases for g , following Corollary 3.8.

Case 1: $g = (ab)^{2^{n-1}}$. From Lemma 3.2, we know that $(ab)^s a = a(ab)^{-s}$ and $(ab)^s a$ commutes with $(ab)^{2^{n-1}}$, giving us

$$\begin{aligned}
(ab)^{2^{n-1}} (ab)^s a &= (ab)^{2^{n-1}} a(ab)^{-s} \\
&= (ab)^{2^{n-1}} (ab)^{-s} a \\
&= (ab)^{2^{n-1}-s} a.
\end{aligned}$$

Letting $t = 2^{n-1} - s$, we have shown that in this case, $H = \langle (ab)^s a, (ab)^t a \rangle$, where $s \neq t$ and $s + t = 2^{n-1}$.

Case 2: $g = (ab)^r a$ for some r . Without loss of generality, assume $s > r$. Then H also contains $(ab)^s a g$, which is equal to $(ab)^s a (ab)^r a$, which simplifies to $(ab)^{s-r}$ using the fact that $(ab)^m = (ba)^{2^n-m}$. By Lemma 3.1, if $s-r \neq 2^n$, then $(ab)^{s-r}$ generates a cyclic subgroup of order larger than two, contradicting our assumption that $H \cong C_2 \times C_2$. Thus it must be the case that $s-r = 2^n$, and the result follows. \square

Corollary 3.11. *Let $g \in D(2^n)$ such that $g = (ab)^j a$ for some $0 \leq j \leq 2^{n-1}$. If H is a powerful subgroup of $D(2^n)$ that contains g , then H is either trivial, isomorphic to C_2 , or isomorphic to $C_2 \times C_2$.*

Remark. From Lemma 3.10, it follows that there are exactly 2^{n-1} subgroups of $D(2^n)$ that are isomorphic to $C_2 \times C_2$ – one for each pair of numbers s, t with $0 \leq s, t \leq 2^{n-1}$ with $s \neq t$ such that $s + t = 2^n$. It also follows from this Lemma that if H_i and H_j are distinct subgroups isomorphic to $C_2 \times C_2$, then $H_i \cap H_j = \langle (ab)^{2^{n-1}} \rangle$, which is the center of $D(2^n)$.

4. MAIN RESULTS

Proposition 4.1. *There exists a powerful cover of $D(2^n)$ with $2^{n-1} + 1$ subgroups.*

Proof. Let $H_1 = \langle ab \rangle$, and for $1 \leq r \leq 2^{n-1}$, let $H_{r+1} = \langle (ab)^r a, (ab)^{2^n-r} a \rangle$. There are $2^{n-1} + 1$ subgroups, each H_i is abelian, and every element of $D(2^n)$ is contained in some H_i . \square

Theorem 4.2. *The powerful subgroup covering number of $D(2^n)$ is $2^{n-1} + 1$.*

Proof. By Proposition 4.1, we know that the powerful covering number of $D(2^n)$ is at most $2^{n-1} + 1$. Now we show that $D(2^n)$ can not be covered by fewer than $2^{n-1} + 1$ powerful subgroups. Let $\mathcal{C} = \{H_1, \dots, H_q\}$ be a powerful cover of $D(2^n)$. Appealing to Proposition 3.9 and re-indexing if necessary, we may assume that $H_1 = \langle ab \rangle$, so $|H_1| = 2^n$. Now we claim that for each i with $2 \leq i \leq q$, the subgroup H_i is isomorphic to either C_2 or $C_2 \times C_2$. This follows from Proposition 3.7, and from the fact that if $H_i \cong C_k$ for some $k > 2$, we would have $H_i \subseteq \langle ab \rangle$. This would make the covering redundant, and hence no minimal. From Remark 3 each H_i that is isomorphic to $C_2 \times C_2$ contains e and $(ab)^{2^{n-1}}$, which are already in H_1 , so each

H_i that is isomorphic to $C_2 \times C_2$ contributes two new elements, while each H_i that is isomorphic to C_2 contributes one new element. This means that

$$\left| \bigcup_{i=1}^q H_i \right| \leq |H_1| + 2(q-1) = 2^n + 2(q-1).$$

In other words, the subgroups H_2, \dots, H_q can contain at most $2(q-1)$ elements not contained in H_1 . If $q < 2^{n-1}$, we would then have

$$\left| \bigcup_{i=1}^q H_i \right| < 2^n + 2(2^{n-1}) = 2^{n+1},$$

meaning that this collection of subgroups could not be a cover for the 2^{n+1} elements of $D(2^n)$. Thus, any cover of $D(2^n)$ by powerful subgroups must contain at least $2^{n-1} + 1$ powerful subgroups. This completes the proof. \square

Corollary 4.3. *For $n \geq 2$, $\sigma_P(D(2^n)) = \sigma_A(D(2^n))$.*

Proof. This follows from Theorem 4.2 and Corollary 3.7. \square

5. CONCLUSION

There are three 2-groups of coclass equal to 1. We have explicitly calculated the powerful covering number for one of them.

REFERENCES

- [1] *Abstract Algebra*. John Wiley & Sons, 2004. 1

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, STILLWELL 426, WESTERN CAROLINA UNIVERSITY, CULLOWHEE, NC 28723 USA

E-mail address: ratanasov@email.wcu.edu

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, STILLWELL 426, WESTERN CAROLINA UNIVERSITY, CULLOWHEE, NC 28723 USA

E-mail address: adgregory1@catamount.wcu.edu

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, STILLWELL 426, WESTERN CAROLINA UNIVERSITY, CULLOWHEE, NC 28723 USA

E-mail address: lrguatelli1@catamount.wcu.edu

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, STILLWELL 426, WESTERN CAROLINA UNIVERSITY, CULLOWHEE, NC 28723 USA

E-mail address: adpenland@email.wcu.edu