



COMMUNITY PHARMACY BLOOD PRESSURE ANALYSIS

BY
DR. RITA AIWUYO

Total Encounters

381

Unique Patients

140


Returning Patients

63


Hypertensive Encounter Rate

43.6%


BP Classification by Sex



BP Status at Latest Follow-up




Unique Patients by Age Group



KEY INSIGHTS

- 43.6% of recorded BP encounters were within the hypertensive range (Stage 1 or Stage 2).
- 45.0% of patients returned for at least one follow-up BP assessment (63 of 140 patients).
- Adults aged 50–59 years formed the largest patient group, accounting for 47 of 140 patients (33.6%).
- Prehypertension was the most frequently recorded BP category (134 encounters), followed by Stage 1 hypertension (117 encounters).

Total Encounters by BP Classification





FOLLOW-UP & BLOOD PRESSURE CHANGE

Longitudinal analysis of patients with repeat BP encounters

BY
DR. RITA AIWUYO

Returning Patients

63

Improved Patients

18


Unchanged Patients

30


Worsened Patients

15

Patient BP Status at Latest Follow-up



Average BP: Baseline vs Latest Follow-up



Follow-up BP Status by Sex

sex	Improved Patients	Unchanged Patients	Worsened Patients
Female	9	17	6
Male	9	13	9
Total	18	30	15

KEY INSIGHTS

- 18 of 63 returning patients (28.6%) moved to a lower BP category by their latest recorded follow-up.
- Average BP decreased from 131.4/83.2 mmHg at baseline to 126.7/81.1 mmHg at the latest recorded follow-up.
- 15 of 63 returning patients (23.8%) moved to a higher BP category, highlighting a subgroup that may warrant closer clinical review.
- Nearly half of returning patients (47.6%) remained in the same BP category at their latest follow-up.