

Assignment 1

S. Rithvik Reddy - cs20btech11049

Download all python codes from

<https://github.com/rithvikreddy6300/Assignment-1/blob/main/Assignment-1.py>

and latex-tikz codes from

<https://github.com/rithvikreddy6300/Assignment-1/blob/main/Assignment-1.tex>

The expectation value of X represented by E(X) is given by

$$E(x) = \sum_{X=1}^6 Pr(X = n).X$$

$$\Rightarrow E(X) = \sum_{X=1}^6 \frac{(2X - 1)}{36}.X \quad (0.0.2)$$

$$\Rightarrow E(X) = \sum_{X=1}^6 \frac{(2X^2 - X)}{36} \quad (0.0.3)$$

$$\Rightarrow E(X) = \frac{2}{36} \cdot \sum_{X=1}^6 X^2 - \frac{1}{36} \sum_{X=1}^6 X \quad (0.0.4)$$

$$\Rightarrow E(X) = \frac{2}{36}.91 - \frac{1}{36}.21 \quad (0.0.5)$$

$$\Rightarrow E(X) = \mathbf{4.4722} \quad (0.0.6)$$

QUESTION

Two numbers are selected at random (without replacement) from the first six positive integers. Let X denote the larger of the two numbers obtained. Find E(X)?

SOLUTION

Let X_1, X_2 be the 1st, 2nd numbers drawn randomly from 1 to 6 and $X = \max(X_1, X_2)$
let $\max(X_1, X_2) = n$, $X_i \in \{1, 2, 3, 4, 5, 6\}, i = 1, 2$ so $X \in \{1, 2, 3, 4, 5, 6\}$, The probability mass function is

$$p_{X_i}(n) = Pr(X_i = n) = \begin{cases} \frac{1}{6}, & \text{if } 1 \leq n \leq 6 \\ 0, & \text{otherwise} \end{cases}$$

$$p_X(n) = Pr(\max(X_1, X_2) = n)$$

$$= Pr(X_1 = n \text{ and } X_2 < n) + Pr(X_2 = n \text{ and } X_1 < n) + Pr(X_1 = X_2 = n) \quad (0.0.1)$$

Since choosing of X_1, X_2 are independent events we can write

$$Pr(X_1 \text{ and } X_2) = Pr(X_1).Pr(X_2)$$

Substituting this in (0.0.1) gives us

$$p_X(n) = Pr(X_1 = n).Pr(X_2 < n) + Pr(X_2 = n).Pr(X_1 < n) + Pr(X_1 = n).Pr(X_2 = n)$$

$$\Rightarrow p_X(n) = Pr(X = n) = \frac{1}{6} \cdot \frac{(n-1)}{6} + \frac{1}{6} \cdot \frac{(n-1)}{6} + \frac{1}{6} \cdot \frac{1}{6}$$

$$\Rightarrow p_X(n) = Pr(X = n) = \frac{(2n-1)}{36}$$

Therefore the expectation value of X, E(X) = **4.4722**.