Assignment-1

S. RITHVIK REDDY- cs20btech11049

Download all python codes from here

https://github.com/rithvikreddy6300/ Assignment-1/blob/main/Assignment-1.py

and latex-tikz codes from

https://github.com/rithvikreddy6300/ Assignment-1/blob/main/Assignment-1.tex

QUESTION-4.10

Two numbers are selected at random (without replacement) from the first six positive integers. Let X denote the larger of the two numbers obtained. Find E(X)?

SOLUTION

Let X_1, X_2 be the $1^{st}, 2^{nd}$ numbers drawn randomly from 1 to 6 and X = max (X_1, X_2)

let max (X_1,X_2) =n , $X_i \in \{1,2,3,4,5,6\}, i=1,2$ so $X \in \{1,2,3,4,5,6\}$, The probability mass function is

$$p_{X_i}(n) = Pr(X_i = n) = \begin{cases} \frac{1}{6}, & \text{if } 1 \le n \le 6\\ 0, & \text{otherwise} \end{cases}$$
$$p_X(n) = Pr(\max(X_1, X_2) = n)$$

$$= Pr(X_1 = n \text{ and } X_2 < n) + Pr(X_2 = n \text{ and } X_1 < n) + Pr(X_1 = X_2 = n)$$
 (1)

Since choosing of X_1, X_2 are independent events we can write

$$Pr(X_1 \text{ and } X_2) = Pr(X_1).Pr(X_2)$$

Substituting this in (1) gives us

$$p_X(n) = Pr(X_1 = n).Pr(X_2 < n) + Pr(X_2 = n)$$

$$.Pr(X_1 < n) + Pr(X_1 = n).Pr(X_2 = n)$$
(2)

$$\implies p_X(n) = Pr(X = n) = \frac{1}{6} \cdot \frac{(n-1)}{6} + \frac{1}{6} \cdot \frac{(n-1)}{6} + \frac{1}{6} \cdot \frac{1}{6}$$

$$\implies p_X(n) = Pr(X = n) = \frac{(2n-1)}{36}$$

The expectation value of X represented by $\mathrm{E}(\mathrm{X})$ is given by

$$E(x) = \sum_{X=1}^{6} Pr(X = n).X$$

$$\Longrightarrow E(X) = \sum_{X=1}^{6} \frac{(2X-1)}{36} . X \tag{3}$$

$$\Longrightarrow E(X) = \sum_{X=1}^{6} \frac{(2X^2 - X)}{36} \tag{4}$$

$$\Longrightarrow E(X) = \frac{2}{36} \cdot \sum_{X=1}^{6} X^2 - \frac{1}{36} \sum_{X=1}^{6} X \tag{5}$$

$$\implies E(X) = \frac{2}{36}.91 - \frac{1}{36}.21$$
 (6)

$$\Longrightarrow E(X) = 4.4722 \tag{7}$$

Therefore the expectation value of X, E(X) = 4.4722.