


Conserved domains on [lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10]

Local query sequence

Graphical summary ☐ Zoom to residue level show extra options »



List of domain hits

Name	Accession	Description
Lipase	pfam00151	Lipase;
Lipase;		:

Pssm-ID: 395099 Cd Length: 336 Bit Score: 604.44 E-value: 0e+00

```

      10      20      30      40      50      60      70      80
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 271 EVCYERLGCFSDDSPWAGIV-ERPLKILPWSPKVIDTRFLLYTNENPNQCLITGDPETIRNSNFNTRSRKTRFIHGFID 349
Cdd:pfam00151 2 EVCYGQLGCGFDKIPWAGNTLVPRVKSLPWSPKDIDTRFLLYTNENPNQCLITGDPETIRNSNFNTRSRKTRFIHGFID 81

      90     100     110     120     130     140     150     160
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 350 KG-EESWLSTMCQNMFKVESVNCICVDWKSGRSTAYSQASQNVRIVGAEVAYLVGLQSSFDYSPSNVHIIGHSLGSHAA 428
Cdd:pfam00151 82 KGyEESWLSDMCKALQVEDVNVICVDWKSGRSTHYTQAVQNIIRVGAEVANLLQWLSNELNYSPSNVHLIGHSLGAHVA 161

      170     180     190     200     210     220     230     240
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 429 GEAGRRRTGAVGRITGLDPAEPCFQGTPELVRLDPDQAQFVDVIHTDIAPIPNLGFQSGTQAGHLDFPNGGKEMPGQC 508
Cdd:pfam00151 162 GEAGRRRTGKLGRTGLDPAGPYFQGTPEELVRLDPGDAQFVDAIHTDTRP-IPGLGFQISQPVGHVDFPNGGSEQPGQC 240

      250     260     270     280     290     300     310     320
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 509 KNLVLSQIVDIDGIWQGRdFAACNHLRSYKYTDSILNPDGFAFGSCASYSDFTANKCFPCSSGECQMGMHYADRFPGR 588
Cdd:pfam00151 241 KNLISQIIDIDGIWEGTQ-FVACNHLRSYHYIDSLNPRGFGPGYPCSSYDAFSQNKCLCPKGGCPQMGMHYADKFPGKT 319

      330
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 589 KGVGQLFYLTNGDASNF 605
Cdd:pfam00151 320 SKLEQTFYLTNGSSSPF 336
```

Smac_DIABLO pfam09057 Second Mitochondria-derived Activator of Caspases; Second Mitochondria-derived Activator of ...
Second Mitochondria-derived Activator of Caspases promotes apoptosis by activating caspases in the cytochrome c/Apaf-1/caspase-9 pathway, and activity of inhibitor of apoptosis proteins (XIAP-BIR3). The protein assumes an elongated three-helix bundle structure, and forms a dimer in solution.

Pssm-ID: 462664 Cd Length: 231 Bit Score: 387.66 E-value: 9.97e-129

```

      10      20      30      40      50      60      70      80
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 39 SRSVTSFFRYRQ-CLCVPVVFANfkrccfseliRPWHKVTITIGFVTLCAVPFAQKSEPHSLSEALMRAVSLVTDST 117
Cdd:pfam09057 1 RRSASVLLRSRghLQSFGLAAN-----RPWSKAVSVGVGATLCAVPFTQKSEP-SLSHEALIRRAVSLVTDSTANT 70

      90     100     110     120     130     140     150     160
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 118 FLSQTTALIEAITETKAVYTLTSLYRQYTSLLGKMNSEEDVWQVIGARAEMTSKHQEYLKLETTWMTAVGLSEMA 197
Cdd:pfam09057 71 FLSQTTLALIDALTEYTKAVYTLISLQKQYANLLGKMNPEEDAIWQVIGARVEMTDQEQEYLRFSNMWMTAVNLSMA 150

      170     180     190     200     210     220     230     240
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 198 AEAAYQTGADQASITARNHIQLVKLVQVEVHQLSRKAETKLAEAQIEELRQKTQEEGEERAEE-----QEAYLRE 268
Cdd:pfam09057 151 AEAAYQSGADQASVTARTHIQVAQSQVEEVRLSQEAETKLAEVQTEIIRRTQGERGASAPQGVelseediPEAYLRE 230

      269 D 269
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 269 D 269
Cdd:pfam09057 231 D 231
```

Tryp_SPc cd00190 Trypsin-like serine protease; Many of these are synthesized as inactive precursor zymogens ...
Trypsin-like serine protease; Many of these are synthesized as inactive precursor zymogens that are cleaved during limited proteolysis to generate their active forms. Alignment contains also inactive enzyme substitutions of the catalytic triad residues.

Pssm-ID: 238113 [Multi-domain] Cd Length: 232 Bit Score: 298.42 E-value: 1.03e-94

```

      10      20      30      40      50      60      70      80
lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10 719 VVGGEDARPHSWPQISLQYlknDTRHRCGGTLIASNFVLTAACHISNT--RTYRVAVGKNNLEVEDDEESLFGVVDTI 790
Cdd:cd00190 1 IVGGEAKIGSFWPQVSLQY---TGGRHFCGGSLSIPRWVLTAACHVYSSApSNYTVRLGSHDLSN-EGGGQVIKVKV 76
```



90100110120130140150160

.....|.....|.....|.....|.....|.....|.....|

lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10797HVHKRWNALLLRNDIALIKLAEHVELSDTIQVACLPEKDSLPPKDYPCYVTGWGRLWTNGPIADKLQOGLQPVVDHATCS876

Cdd:cd0019077IVHPNYNPSTYDNDIALLLKLRPVTLSDNVRPICLPSSGYNLPAGTTCTVSGWGRTSEGGPLPDVLQEVNVPIVSNAECK156

170180190200210220230

.....|.....|.....|.....|.....|.....|.....|

lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10877RIDWGWFRVKKTMVCAGG-DGVISACNGDSGGPLNCQLeNGSWEVFGIVSFGSrrGCNTRKKPVVYTRVSAYIDWINEK954

Cdd:cd00190157RAYSYGGTITDNMLCAGGLEGGKDACQDGSGGPLVCND-NGRGVLVGIVSWGSGCARNPYPGVYTRVSSYLDWIQKT232

PLAT_PL cd01759 PLAT/LH2 domain of pancreatic triglyceride lipase. Lipases hydrolyze phospholipids and ...
PLAT/LH2 domain of pancreatic triglyceride lipase. Lipases hydrolyze phospholipids and triglycerides to generate fatty acids for energy production or for storage and to release inositol phosphates that ac
The central role of triglyceride lipases is in energy production. The proposed function of PLAT/LH2 domains is to mediate interaction with lipids or membrane bound proteins.

:

Pssm-ID: 238857 Cd Length: 113 Bit Score: 208.76 E-value: 3.05e-63

1020304050607080

.....|.....|.....|.....|.....|.....|.....|

lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10608WRYRVVDVTLSGK-KVTGHLVSLFGNKGNSRQYEIFQGLKPDNTYSNEFSDVEVGDLEKVKFIWYNNVINLTLPKVGA686

Cdd:cd017591WRYKVSVDLTSGLKkKVTGTILVSLYGNKGNTROQYEIFKGTLPKNTYSAFIDVDVDVGLPLTKVKFIWNNVINLTLPKVGA80

90100110

.....|.....|.....|

lcl|seqsig_MLRAA_2bd2e7bbc0dc13c100c25f6e28d32f10687SKITVERN-DGSVFNFCSEETVREDVLLTLTAC718

Cdd:cd0175981EKITVQSGkDGKVFNFCSSEETVRENVLQTLTPC113

References:

 Wang J et al. (2023), "The conserved domain database in 2023", **Nucleic Acids Res.**51(D)384-8.

 Lu S et al. (2020), "The conserved domain database in 2020", **Nucleic Acids Res.**48(D)265-8.

 Marchler-Bauer A et al. (2017), "CDD/SPARCLE: functional classification of proteins via subfamily domain architectures.", **Nucleic Acids Res.**45(D)200-3.