

Partial Differential Equations (MA20103)

Assignment – 3

Second order PDE

Q1. Solve the following second order homogenous PDE with constant coefficients (symbols

have usual meanings, say, $r = \frac{\partial z}{\partial x}$; $s = \frac{\partial^2 z}{\partial x \partial y}$; $t = \frac{\partial z}{\partial y}$)

(i) $25r - 40s + 16t = 0$

(ii) $r + (a + b)s + abt = xy$

(iii) $r - t = x - y$

(iv) $r + t + 2s = xy$

(v) $2r - 3s - 2t = 0$

(vi) $r - 4s + 4t = 0$

(vii) $r + 3s + 2t = 2x + 3y$

(viii) $r - s - 2t = (y - 1)e^x$

(ix) $r - 5s + 4t = \sin(4x + y)$

(x) $r + t = \cos mx \cos ny$

Q2. Classify and reduce the following equations in to canonical form

(i) $\frac{\partial^2 z}{\partial x^2} - x^2 \frac{\partial^2 z}{\partial y^2} = 0$ (also find general solution)

(ii) $\frac{\partial^2 z}{\partial x^2} + x^2 \frac{\partial^2 z}{\partial y^2} = 0$ (also find general solution)

(iii) $16 \frac{\partial^2 z}{\partial x^2} - y^{10} \frac{\partial^2 z}{\partial y^2} = 5y^9 \frac{\partial z}{\partial y}$

(iv) $y^2 \frac{\partial^2 z}{\partial x^2} - 2xy \frac{\partial^2 z}{\partial x \partial y} + x^2 \frac{\partial^2 z}{\partial y^2} = \frac{y^2}{x} \frac{\partial z}{\partial x} + \frac{x^2}{y} \frac{\partial z}{\partial y}$

(v) $(y - 1) \frac{\partial^2 z}{\partial x^2} - (y^2 - 1) \frac{\partial^2 z}{\partial x \partial y} + y(y - 1) \frac{\partial^2 z}{\partial y^2} + \frac{\partial z}{\partial x} - \frac{\partial z}{\partial y} = 2ye^{2x}(1 - y)^3$

Q3. Show that the solution of the equation

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$$

satisfying the conditions

(i) $u \rightarrow 0$ as $t \rightarrow \infty, \forall x$

(ii) $u = 0$ for $x = 0$ and $x = a \forall t > 0$

(iii) $u = x$ when $t = 0$ and $0 < x < a$ is

$$u(x, t) = \frac{2a}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \sin\left(\frac{n\pi x}{a}\right) \exp\left[-\left(\frac{n\pi}{a}\right)^2 t\right]$$

Q4. Solve $\nabla^2 u = 0$

subject to $u(x, 0) \neq 0, u(x, a) = 0, u(x, y) \rightarrow 0$ as $x \rightarrow \infty$ where $x \geq 0$ and $0 \leq y \leq a$

Q5. Solve the 2 dimensional Laplace equation in polar co-ordinates r and θ in the region $0 \leq r \leq a, 0 \leq \theta \leq 2\pi$ subject to

(i) u remains finite as $r \rightarrow 0$

(ii) $u = \sum_n c_n \cos(n\theta)$ on $r = a$

Q6. A tightly stretched string with fixed end point $x = 0$ and $x = l$ is initially in a position given by $u = u_0 \sin^3 \frac{\pi x}{l}$. If it is released from rest from this position, show that the displacement is given by

$$u(x, t) = \frac{u_0}{4} \left(3 \cos \frac{\pi ct}{l} \sin \frac{\pi x}{l} - \cos \frac{3\pi ct}{l} \sin \frac{3\pi x}{l} \right)$$