



< Previous



Next >

## Distance Exercises

Bookmark this page

## Distance Exercises #1

1/1 point (graded)

If you have not done so already, install the data package **tissuesGeneExpression**.

```
library(devtools)
install_github("genomicsclass/tissuesGeneExpression")
```

The data represents RNA expression levels for seven tissues, each with several *biological replicates*. We call samples that we consider to be from the same population, such as liver tissue from different individuals, *biological replicates*:

```
library(tissuesGeneExpression)
data(tissuesGeneExpression)
head(e)
head(tissue)
```

How many biological replicates are there for hippocampus?

✓ Answer: 31

31

Explanation

```
sum(tissue=="hippocampus")
##to answer this question for all tissues look at
table(tissue)
```

Submit

You have used 1 of 5 attempts

❗ Answers are displayed within the problem

## Distance Exercises #2

1/1 point (graded)

What is the distance between samples 3 and 45?

✓ Answer: 152.5662

152.5662

Explanation

```
sqrt( crossprod(e[,3]-e[,45]) )
## or
sqrt( sum((e[,3]-e[,45])^2 ) )
```

Submit

You have used 1 of 5 attempts

**i** Answers are displayed within the problem

### Distance Exercises #3

1/1 point (graded)  
What is the distance between gene `210486_at` and `200805_at` ?

41.01153

✓ Answer: 41.01153

41.01153

Explanation

```
sqrt( crossprod(e["210486_at"],-e["200805_at"],)) )  
## or  
sqrt( sum((e["210486_at"],-e["200805_at"],)^2 ))
```

Submit

You have used 1 of 5 attempts

**i** Answers are displayed within the problem

### Distance Exercises #4

1/1 point (graded)  
If I run the command (**don't run it!**):

```
d = as.matrix(dist(e))
```

How many cells (number of rows times number of columns) would this matrix have?

493506225

✓ Answer: 493506225

493506225

Explanation

```
##every pair of rows has an entry:  
nrow(e)^2
```

Submit

You have used 1 of 5 attempts

**i** Answers are displayed within the problem

## Distance Exercises #5

1/1 point (graded)

Compute the distance between all pairs of samples:

```
d = dist(t(e))
```

Read the help file for `dist()`.

How many distances are stored in `d`? (Hint: What is the length of `d`)?

✓ Answer: 17766

17766

Explanation

```
length(d)
```

Submit

You have used 2 of 5 attempts

❗ Answers are displayed within the problem

## Distance Exercises #6

1/1 point (graded)

Why is the answer above not `ncol(e)^2`?

☐ R made a mistake there

☐ Distances of 0 are left out

☒ Because R takes advantage of symmetry: only the lower triangular matrix is stored, thus there are only `ncol(e)*(ncol(e)-1)/2` values.

☐ Because it is equal to `nrow(e)^2`



Explanation

Note that the distance between samples *i* and *j* is the same as distance between samples *j* and *i*. Also the distance between a sample and itself is 0. The object returned by `dist()` avoids storing all those values.

Submit

You have used 1 of 2 attempts

❗ Answers are displayed within the problem



# edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

# Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

# Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)

