

# Dynamics of non-household contacts during the COVID-19 pandemic in 2020 and 2021 in the Netherlands

Jantien A. Backer<sup>1,\*</sup>, Laurens Bogaardt<sup>1</sup>, more authors<sup>?</sup>, and Jacco Wallinga<sup>1,2</sup>

<sup>1</sup>National Institute for Public Health and the Environment, Bilthoven, the Netherlands

<sup>2</sup>Leiden University Medical Center, Leiden, the Netherlands

\*jantien.backer@rivm.nl

## ABSTRACT

The COVID-19 pandemic in 2020 and 2021 was for a large part mitigated by reducing contacts in the general population. To monitor how these contacts changed over the course of the pandemic in the Netherlands, a survey was conducted where participants reported every two weeks on their at-risk contacts, as part of the European CoMix survey. The survey included 1659 participants from April to August 2020 and 2514 participants from December 2020 to September 2021.

We categorized the number of unique contacted persons excluding household members, reported per participant per day into six activity levels, viz 0, 1, 2, 3-4, 5-9 and 10 or more reported contacts. After correcting for age, vaccination status, risk status for severe outcome of infection, and duration of participation, activity levels increased over time, coinciding with relaxation of COVID-19 control measures.

## Introduction

From early 2020 onwards, COVID-19 grew into a global pandemic within a few months affecting nearly all countries worldwide. SARS-CoV-2 virus that causes COVID-19 is transmitted between persons during contact events with physical proximity. Reducing transmission with non-pharmaceutical measures can be achieved by lowering the transmission probability per contact, e.g. by maintaining a safe distance from each other or using protective equipment such as a face mask, and by reducing the number of contacts per person, e.g. by closing schools, theaters and restaurants, teleworking or inviting fewer people at home.

In the Netherlands, the first nation-wide lockdown was imposed on March 15th, 2020 involving all of the above measures. It was lifted stepwise on May 11th (partly opening of schools), June 2nd (further opening of schools, museums and terraces open a.o.) and July 6th (bars and restaurants open a.o.). The rise of the Alpha variant in winter 2020/2021 called for an even stricter lockdown including an evening curfew, that was relaxed from April 28th, 2021 onwards while vaccination coverages were increasing. It is not known how the change in control measures affected the number of at-risk contacts in the Dutch general population.

To learn about the effect of control measures on contact behaviour, many countries set up contact surveys<sup>1-8</sup>. The POLYMOD study<sup>9</sup> served as a blueprint for these as it did for many pre-COVID-19 studies<sup>10</sup>. But while these pre-COVID-19 studies aimed to measure 'normal' contact behaviour, the unprecedented measures that were taken during the pandemic gave a unique opportunity to determine contact behaviour under restricted conditions. As part of a European project, the CoMix survey was set up in April 2020 in Belgium, the UK, and the Netherlands, later followed by 19 other European countries<sup>11</sup>. In this contact study, participants were asked to regularly report their contact behaviour, i.e. the number of unique persons they had contact with in certain locations per day, as well as some characteristics of the contacted persons such as age and sex. Participants were asked to report only at-risk contacts that could have led to an infection, specifically face-to-face conversational contacts and/or physical contacts. What sets the CoMix survey apart from other contact surveys is that it uses a standardised questionnaire in multiple countries and is repeated at regular intervals.

At the same time the CoMix survey started, a large-scale survey on SARS-CoV-2 serology was initiated in the Netherlands<sup>12</sup>. This Pienter Corona (PiCo) study was repeated every 4 to 6 months in a representative sample of the Dutch population involving a few thousand participants per survey round, and is still ongoing. It also included a contact question where the participants were asked to report the number of unique persons they contacted the day before in specific age groups. An early analysis of the first two survey rounds in 2020 showed that the first lockdown decreased the number of community contacts per participant on average by 76% compared to pre-COVID behaviour<sup>4</sup>. The PiCo survey rounds are however too far apart to closely monitor the contact behaviour in time.

Our aim is to describe how control measures affected the contact behaviour of the Dutch general population, using the

CoMix data that was collected in the Netherlands during the survey periods in 2020 and 2021. We will focus on the number of contacts per participant outside the household because contacts with household members are not subject to control measures. As a validation, the results are compared to four survey rounds of the PiCo survey that coincided with the CoMix study periods.

## Methods

### CoMix participant recruitment and survey design in the Netherlands

This contact survey was part of the CoMix study that was conducted in several EU countries and the UK<sup>11</sup>. Participants were recruited by the market research company Ipsos-MORI, and filled out the contact survey online. The study was carried out in two survey periods: 8 survey rounds from 16 April to 5 August 2020, referred to as the 2020 series, and 20 survey rounds from 23 December 2020 to 22 September 2021, referred to as the 2021 series. At the start of the 2020 series, 1500 participants of 18 years and older were recruited, reflecting the age distribution of the Dutch population. An overall drop-out rate of 10% per round was allowed, but these rates differed per age group leading to very few participants in the youngest age group of 18-24 in the last rounds of the 2020 series (Fig 1). For this reason, a preferential sampling scheme was adopted at the start of the 2021 series, oversampling the younger age groups. In total 1200 participants of 18 years and older included and additionally 300 children between 0 and 17 years of age were included by asking adult participants to fill out the contact survey for one of their children. After 10 rounds, the age cohorts were supplemented to include 1500 participants in total again in May 2021.

distributed over different age groups according to the Dutch population over 18 years of age in the 2020 series, and according to a preferential sampling scheme in the 2021 series

The contact survey was repeated every two weeks and a survey round lasted at most 7 days, except for the survey rounds where many new participants were recruited that were allowed more time to fill out the questionnaire. In each survey round participants were asked about characteristics that can change over time, such as risk perception, risk status and vaccination status against COVID-19. High risk participants were people with chronic respiratory disease, chronic heart disorder, chronic kidney disease, diabetes, reduced resistance to infection (due to illness or medication), morbid obesity (BMI over 40), and pregnant women, i.e. reflecting the medical indication for influenza vaccination. Missing data in the risk status were completed with the most common risk status reported over the survey rounds by the participant. The risk status of participants can change by survey round. The fraction of high risk participants is calculated per age group using only the participants with unambiguous risk status, and compared to the population fraction with a medical indication for influenza vaccination<sup>13</sup>.

The 2021 series coincided with the roll-out of the COVID-19 vaccination campaign. Health care workers, persons with a high medical risk and care home residents were vaccinated with priority, while the non-risk groups were vaccinated from elderly to young. Most persons were vaccinated with either BionTech/Pfizer, Moderna or Janssen vaccine by the Municipal Health Services, except for 60-64 year olds who were vaccinated at an early stage with AstraZeneca by their general practitioner. Participants are considered to be vaccinated after they have reported to have had at least one vaccination dose. The fraction of vaccinated participants is compared to the vaccination coverage of one vaccine dose in the general population over time<sup>14</sup>.

In the main part of the survey, participants reported their contact behaviour of the previous day. For each unique person they had conversational or physical contact with between 5am on the previous day until 5am of the survey day, participants filled out the characteristics of the contacted person, such as age group and gender, as well as duration and location of the contact event. Multiple contact events with the same person were to be considered as one contact, and only contact events with a close physical proximity were to be reported, i.e. excluding online or phone contacts. Besides these individual contacts, it was also possible to report group contacts from the third survey round onwards in late May 2020. Group contacts were reported as the total number of contacted persons in broad age groups (0-17, 18-64, 65+) at work, school or another location. More details on the CoMix study and questionnaire can be found in previous publications<sup>11,15,16</sup>.

### Analysis of number of contacts

The participants are stratified into three age groups (0-17, 18-64, 65+) and two series (2020, 2021). Only contacts with non-household members are included, as these are most likely to change over the course of the study period. Twelve occasions were excluded where participants reported more than 1000 contacts in a single survey round, and 58 participants were excluded that participated in four or more rounds but did not report any contact.

The distribution of the number of contacts per participant per round had a heavy tail, which can result in unstable estimates of the variance<sup>17</sup>. Instead of using a parametric distribution, we define activity levels of 0, 1, 2, 3-4, 5-9,  $\geq 10$  contacts per participant per round and analyse these using ordinal regression. The activity levels are used as variables of a generalised additive model with fixed effects for participant vaccination status, participant risk status, weekends and (school)holidays, cubic splines for calendar time, participant age, participant round (i.e. the  $n^{\text{th}}$  time a participant participated), and a random intercept for each participant. The statistic model can be described as:

$$Pr(\text{activity} \leq l) = \Phi[\theta_l - (\beta_{id} + \beta_0 + \beta_1 * s(\text{date}) + \beta_2 * s(\text{part\_round}) + \beta_3 * s(\text{part\_age}) + \beta_4 * \text{part\_vacc} + \beta_5 * \text{part\_risk} + \beta_6 * \text{weekend} + \beta_7 * \text{holiday})],$$

where  $\Phi$  is the logistic function,  $\theta_l$  are the thresholds for the activity levels  $l$ ,  $\beta_{id}$  are the random intercepts and the  $s()$  function indicates that splines are used. The fits are assessed by explained deviance and by comparing the fitted activity levels to the observations.

Next, the fitted results are used in a synthetic population that has the same size as the study population, but reflects the general population, with respect to age<sup>18</sup>, risk status<sup>13</sup> and vaccination status<sup>14</sup> at each day of the study periods. Participant id's are sampled with replacement from the study population, and participant round is set to 1 for each day during the study period. 200 parameter sets are sampled from the estimated model parameters, and combined with 200 synthetic populations, to capture both the uncertainty of the model parameters and the uncertainty caused by the sample size of the study population. As a validation, the predicted activity levels of the synthetic population are compared to the activity levels measured in PiCo survey rounds 1 (Apr 2020), 2 (Jun 2020), 4 (Feb 2021) and 5 (Jun 2021).

All analyses are performed in R version 4.1.3<sup>19</sup> using the *mgcv* package<sup>20</sup> with family *ocat* for ordered categorical variables. Code is available on Github<sup>21</sup>, and data is published on Zenodo<sup>22</sup> in the standardised format of *socialcontactdata.org*.

## Results

### Study population

In total, 1659 participants were included in the 2020 series, and 2514 participants in the 2021 series. Participants were equally distributed over men and women in 2020, while female participants were slightly overrepresented in 2021. Household sizes of 2 persons were most common (Tab. 1). Each series started with a target population of 1500 participants, and the number of participants declined in each survey round (Tab S1, Fig S1). In 2020 the drop-out rate was on average 4% per round in the oldest age group and 14% in the youngest age group; in 2021 drop out rates were more comparable for all age groups (Fig. 1A).

The fraction of high risk adult participants is higher than in the general population, especially in the 2020 series (Fig. 1B). The vaccination coverage of the study population increases over time largely in agreement with the general population (Fig. 1C). The participants in the age groups of 18-34 years achieve a higher vaccination coverage at an earlier stage than the general population. The participants in the age group of 65 years and older lag behind the general population.

### Contact behaviour

The distribution of the number of contacts reveals that the adult and elderly age groups in both series exhibit heavy tails (Fig. 2). Exponents for these four groups range from 1.1 to 1.5, indicating that the estimates of the sample mean are stable but the estimates of the sample variance are not. The numbers of contacts per participant per round are categorized in activity levels of 0, 1, 2, 3-4, 5-9,  $\geq 10$  contacts per participant per round.

Separate analyses are carried out for the different series and age groups. Of the fixed effects only the weekends significantly reduces the activity levels in all analyses (Tab. S2). Activity levels increase over time during the two series, indicating that the underlying number of non-household contacts per participant increases (Fig. S2), corresponding to the lifting of COVID-19 control measures. In all analyses the activity levels decrease with participant round. This means that participants tend to report fewer contacts the more often they participate. This is especially the case for the first few rounds, after which the effect stabilises.

The deviance explained by this model ranges from 14% to 21% (Tab. S2). Using these model results the fitted activity levels in the study population are compared to the data (Fig. 3). With the exception of some rounds, notably the first round in 2020 and the fourth round in 2021, the fitted activity levels agree well with the observed activity levels.

Next, the model results are extrapolated to the general population over the full course of the study periods (Fig. 4). The saw-tooth pattern is caused by the weekend effect, which is largest for the 0-17 age group. Over time activity levels increase in all age groups in both series, indicated by the decrease of the predicted lines in Fig. 4. As a comparison, the results of the PiCo study<sup>4</sup> are plotted, of which rounds 1, 2, 4 and 5 fell within the CoMix study periods. Participants of the PiCo study reported higher activity levels than CoMix participants. The discrepancy is largest for the 0-17 age group, whereas the 65+ age group all but agrees with the PiCo study results.

## Discussion

In this paper we showed how the number of at-risk contacts per person outside the household increased as the COVID-19 control measures were lifted in the Netherlands, based on the Dutch data of the European CoMix survey.

The independent contact surveys of the PiCo studies show a similar increasing trend in activity levels in the general population as our results, but the activity levels reported in the PiCo data are higher than in the CoMix data. The two studies differed in how the participants report their contacts. In the CoMix study contacts were reported individually with a lot of detail on location, duration, distance from contact, protection measures, etc. Also for the group contacts at work, school, or other places additional details were requested. In the PiCo study a participant reported the total number of contacts in specific age classes, which made reporting many contacts easier. This may partly explain the higher activity levels found in the PiCo study.

Participants tend to report fewer contacts when they participate in more rounds. This fatigue effect is also seen in the CoMix data of other countries, but the fatigue effect in the Netherlands is found to be extreme compared to these<sup>23</sup>. A complicating factor in interpreting the fatigue effect is that the participant rounds are not equally distributed over the study period. Most participants have their first round in survey round 1 in 2020, and survey rounds 1 and 11 in 2021, due to the study design. In an alternative design as was used in the UK, the study population is supplemented to reach the target size in every survey round, which leads to a better distribution of the participant rounds over the survey rounds<sup>16</sup>. Another possible adjustment to the study design is to decrease the frequency of the survey rounds, which may lead to a smaller fatigue effect.

In order to say anything about the general population, the fatigue effect needed to be corrected for, and differences between the study population and the general population had to be taken into account. Particularly the age distribution of the study participants varied over the survey rounds, because drop-out rates differed between age groups. Also vaccination and risk status of the participants did not always reflect the vaccination and risk status in the general population. The participants in the age groups of 18-34 years achieved a higher vaccination coverage at an earlier stage than the general population. This could indicate that these groups contained relatively many health care workers, who were vaccinated with priority and may have a higher willingness to get vaccinated. The participants in the age group of 65 years and older lagged behind the general population, because the study population probably did not contain many care home residents who were vaccinated with priority. The fraction of high risk adult participants was higher than in the general population. The risk status was reported by participants themselves and compared to the fraction of the population that is invited for the annual influenza vaccination. Although the high risk definition is the same for both populations, participants may have assessed their risk status differently than their GP would. This is supported by the finding that the high risk participants are mainly overrepresented in 2020 (Fig. 1B, Tab. 1) when risks may have been perceived to be higher than in 2021. This would explain the discrepancy, but it can not be excluded that they were really high risk participants.

Our findings suggest that the contact data collected in the CoMix survey describe trends in contact behaviour in the general population. As such it can be useful in monitoring the effect of control measures that are aimed at contact reduction<sup>16,24,25</sup>, describe determinants for contact behaviour<sup>26,27</sup>, or serve as input for infectious disease models<sup>28-30</sup>. These results can help guide policy in future waves of COVID-19 or other emerging respiratory diseases and emphasize that contact surveys such as the CoMix study are indispensable for providing a quantitative basis to public health policy.

## References

1. Zhang, J. *et al.* Changes in contact patterns shape the dynamics of the COVID-19 outbreak in China. *Science* **368**, 1481–1486, DOI: <http://dx.doi.org/10.1126/science.abb8001> (2020).
2. Latsuzbaia, A., Herold, M., Bertemes, J. P. & Mossong, J. Evolving social contact patterns during the COVID-19 crisis in Luxembourg. *PLoS One* **15**, e0237128, DOI: <http://dx.doi.org/10.1371/journal.pone.0237128> (2020).
3. Quaife, M. *et al.* The impact of COVID-19 control measures on social contacts and transmission in Kenyan informal settlements. *BMC Med* **18**, 316, DOI: <http://dx.doi.org/10.1186/s12916-020-01779-4> (2020).
4. Backer, J. A. *et al.* Impact of physical distancing measures against COVID-19 on contacts and mixing patterns: repeated cross-sectional surveys, the Netherlands, 2016–17, April 2020 and June 2020. *Euro Surveill* **26**, DOI: <http://dx.doi.org/10.2807/1560-7917.ES.2021.26.8.2000994> (2021).
5. Bosetti, P. *et al.* Lockdown impact on age-specific contact patterns and behaviours, France, April 2020. *Euro Surveill* **26**, DOI: <http://dx.doi.org/10.2807/1560-7917.ES.2021.26.48.2001636> (2021).
6. Feehan, D. M. & Mahmud, A. S. Quantifying population contact patterns in the United States during the COVID-19 pandemic. *Nat Commun* **12**, 893, DOI: <https://dx.doi.org/10.1038/s41467-021-20990-2> (2021).
7. Tomori, D. V. *et al.* Individual social contact data and population mobility data as early markers of SARS-CoV-2 transmission dynamics during the first wave in Germany-an analysis based on the COVIMOD study. *BMC Med* **19**, 271, DOI: <http://dx.doi.org/10.3201/doi:10.1186/s12916-021-02139-6> (2021).
8. Drolet, M. *et al.* Time trends in social contacts before and during the COVID-19 pandemic: the CONNECT study. *BMC Public Heal.* **22**, 1032, DOI: <https://dx.doi.org/10.1186/s12889-022-13402-7> (2022).

9. Mossong, J. *et al.* Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med.* **5**, e74, DOI: <http://dx.doi.org/10.1371/journal.pmed.0050074> (2008).
10. Hoang, T. *et al.* A Systematic Review of Social Contact Surveys to Inform Transmission Models of Close-contact Infections. *Epidemiology* **30**, 723–736, DOI: <http://dx.doi.org/10.1097/EDE.0000000000001047> (2019).
11. Verelst, F. *et al.* SOCRATES-CoMix: a platform for timely and open-source contact mixing data during and in between COVID-19 surges and interventions in over 20 European countries. *BMC Med* **19**, 254, DOI: <http://dx.doi.org/10.1186/s12916-021-02133-y> (2021).
12. Vos, E. R. A. *et al.* Associations Between Measures of Social Distancing and Severe Acute Respiratory Syndrome Coronavirus 2 Seropositivity: A Nationwide Population-based Study in the Netherlands. *Clin Infect Dis* **73**, 2318–2321, DOI: <http://dx.doi.org/10.1093/cid/ciab264> (2021).
13. Hooiveld, M., Heins, M., Hendriksen, J. & Korevaar, J. Aantal mensen met medische indicatie voor vaccinatie tegen COVID-19 (Number of people with medical indication for vaccination against COVID-19). <https://www.nivel.nl/sites/default/files/bestanden/1003980.pdf> (2021). [Online; accessed 8 August 2022].
14. RIVM. Cumulatieve opkomst tenminste één COVID-19 vaccinatie naar geboortjaar en week (Cumulative coverage of at least one COVID-19 vaccination per birth year and calendar week). <https://www.rivm.nl/covid-19-vaccinatie/cijfers-vaccinatieprogramma> (2022). [Online; accessed 14 April 2022].
15. Jarvis, C. I. *et al.* Quantifying the impact of physical distance measures on the transmission of COVID-19 in the UK. *BMC Med* **18**, 124, DOI: <http://dx.doi.org/10.1186/s12916-020-01597-8> (2020).
16. Gimma, A. *et al.* Changes in social contacts in England during the COVID-19 pandemic between March 2020 and March 2021 as measured by the CoMix survey: A repeated cross-sectional study. *PLoS Med* **19**, e1003907, DOI: <https://doi.org/10.1371/journal.pmed.1003907> (2022).
17. Tagore, S. *Epidemic Models: Their Spread, Analysis and Invasions in Scale-Free Networks*, 1–25 (Springer International Publishing, Cham, 2015).
18. Statistics Netherlands. Bevolking; geslacht, leeftijd en burgerlijke staat, 1 januari 2021. <https://opendata.cbs.nl/statline/#/CBS/nl/dataset/7461bev/table> (2021). [Online; accessed 1 July 2021].
19. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria (2022).
20. Wood, S. *Generalized Additive Models: An Introduction with R* (Chapman and Hall/CRC, 2017), 2 edn.
21. RIVM. CoMixNL. <https://github.com/rivm-syso/CoMixNL> (2022).
22. Backer, J. A., Bogaardt, L. & Wallinga, J. CoMix social contact data (Netherlands). <http://zenodo.org/record/4790347#.YwTvc3ZByUm/>, DOI: <https://dx.doi.org/10.5281/zenodo.4790347> (2022). [Online; accessed 19 July 2022].
23. Wong, K. L. M. *et al.* Social contact patterns during the COVID-19 pandemic in 21 European countries – evidence from a two-year study. *medRxiv* DOI: <http://dx.doi.org/10.1101/2022.07.25.22277998> (2022).
24. Coletti, P. *et al.* CoMix: comparing mixing patterns in the Belgian population during and after lockdown. *Sci Rep* **10**, 21885, DOI: <https://dx.doi.org/10.1038/s41598-020-78540-7> (2020).
25. Jarvis, C. I. *et al.* The impact of local and national restrictions in response to COVID-19 on social contacts in England: a longitudinal natural experiment. *BMC Med* **19**, 52, DOI: <http://dx.doi.org/10.1186/s12916-021-01924-7> (2021).
26. Wambua, J. *et al.* The influence of risk perceptions on close contact frequency during the SARS-CoV-2 pandemic. *Sci Rep* **12**, 5192, DOI: <http://dx.doi.org/10.1038/s41598-022-09037-8> (2022).
27. Wong, K. L. M. *et al.* Pregnancy during COVID-19: social contact patterns and vaccine coverage of pregnant women from CoMix in 19 European countries. *medRxiv* DOI: <http://dx.doi.org/10.1101/2022.06.01.22275775> (2022).
28. Coletti, P. *et al.* A data-driven metapopulation model for the Belgian COVID-19 epidemic: assessing the impact of lockdown and exit strategies. *BMC Infect Dis* **21**, 503, DOI: <https://dx.doi.org/10.1186/s12879-021-06092-w> (2021).
29. Munday, J. D. *et al.* Estimating the impact of reopening schools on the reproduction number of SARS-CoV-2 in England, using weekly contact survey data. *BMC Med* **19**, 233, DOI: <http://dx.doi.org/10.1186/s12916-021-02107-0> (2021).
30. Franco, N. *et al.* Inferring age-specific differences in susceptibility to and infectiousness upon SARS-CoV-2 infection based on Belgian social contact data. *PLoS Comput. Biol* **18**, e1009965, DOI: <https://doi.org/10.1371/journal.pcbi.1009965> (2022).



## Acknowledgements

The authors would like to thank all participants of the CoMix survey. This project has received funding from the European Union's Horizon 2020 research and innovation programme - project EpiPose (grant agreement number 101003688). The study was financed by the Netherlands Ministry of Health, Welfare and Sport.

## Author contributions statement

J.B. managed the data collection, L.B. curated the data, J.B. and L.B. analysed the data, [more authors?], J.W. set up the study. All authors reviewed the manuscript.

## Additional information

**Competing interests** The authors declare no competing interests.

**Ethics statement** This study involves human participants. The research protocol was shared with the Medical Ethics Review Committee Utrecht, and an official waiver for ethical approval was obtained, which stated that ethical assessment was unnecessary given the non-invasive nature of data collection. Participants gave informed consent to participate in the study before taking part.


2020 series		n	%	reference <sup>a</sup> (%)
total		1659		
age group	18-24	84	5	11
	25-34	236	14	16
	35-44	257	15	15
	45-54	337	20	18
	55-64	312	19	17
	65+	433	26	24
gender	Female	814	49	51
	Male	845	51	49
high risk	mixed	359	<sup>b</sup>	
	no	744	58	78
	yes	540	42	22
	NA	16	<sup>b</sup>	
household size	1	409	25	17 <sup>c</sup>
	2	716	43	31
	3	241	15	17
	4	203	12	23
	5+	90	5	12
2021 series		n	%	reference (%)
total		2514		
age group	0-11	240	10	12
	12-17	290	12	7
	18-24	511	20	9
	25-34	378	15	13
	35-44	254	10	12
	45-54	271	11	14
	55-64	230	9	14
	65+	340	14	20
gender	Female	1383	55	50
	Male	1131	45	50
high risk	mixed	385	<sup>b</sup>	
	no	1720	81	81
	yes	400	19	19
	NA	9	<sup>b</sup>	
household size	1	573	23	17 <sup>c</sup>
	2	766	30	31
	3	444	18	17
	4	496	20	23
	5+	235	9	12

**Table 1.** Participants in the CoMix survey in the Netherlands in 2020 and 2021.

<sup>a</sup> Characteristics for 2020 general population based on 18+ population to match study population.


<sup>b</sup> Percentage of risk status in study population only calculated for participants with unambiguous risk status to allow comparison with general population.

<sup>c</sup> Expected household size distribution per Dutch citizen, based on Dutch households consisting of one (38%), two (33%), three (12%), four (12%) and five or more (5%) persons<sup>4</sup>




**Figure 1.** Characteristics of study population, compared to general population. **A** Number of participants included per survey round in the 2020 and 2021 series, by age group. After 10 survey rounds in May 2021, the study population was supplemented to meet the target numbers of the first survey round again. **B** The fraction of high risk participants of the 2020 and 2021 series (points with 95% confidence interval), and of the general population<sup>13</sup> (bars). Only participants with unambiguous risk status are included. **C** Vaccination coverage in 2021 per age group of the study population (points with 95% confidence interval) and of the general population<sup>14</sup> (line). The dashed line denotes the average date when non-risk groups were invited for vaccination. The dashed line for the 55-64 age group is in fact the average for 55-59 year olds, as 60-64 year olds were invited by their general practitioner.






**Figure 2.** Distribution of number of contacts per participant per round excluding household members, by series (line type) and age group (color), with complementary cumulative distribution function on y-axis. The inset shows the same plot on log-log scale.



**Figure 3.** Fitted and observed activity levels over time by series (columns) and age group (rows). Activity levels are shown as the fraction of participants that report more than a certain number of non-household contacts per day. With five limits ( $>0$ ,  $>1$ ,  $>2$ ,  $>4$  and  $>9$ ) six activity levels are defined, e.g. the fraction between the limits of  $>2$  and  $>4$  is the activity level that represents 3 or 4 contacts per participant. The model fits per round are shown by the median (lines) and 95% interval (shaded), from the first to last participation date of that survey round. The observed activity levels per round (open circles) are placed at the mean participation date of that survey round. Holidays and school holiday periods are shaded in grey.




**Figure 4.** Predicted activity levels over time by series (columns) and age group (rows) for the general population. Activity levels are shown as the fraction of the population that has more than a certain number of non-household contacts per day. With five limits ( $>0$ ,  $>1$ ,  $>2$ ,  $>4$  and  $>9$ ) six activity levels are defined, e.g. the fraction between the limits of  $>2$  and  $>4$  is the activity level that represents 3 or 4 contacts per person. The model predictions are shown by the median (lines) and 95% interval (shaded). The activity levels observed in the contact survey of the independent PiCo study are shown for comparison per round at the mean participation date (points), with bootstrapped 95% confidence intervals. Holidays and school holiday periods are shaded in grey.

## Supplementary information

year	round	0-11	12-17	18-24	25-34	35-44	45-54	55-64	65+	total
2020	target	0	0	162	235	225	277	249	352	1500
2020	1	0	0	88	227	236	308	277	386	1522
2020	2	0	0	43	176	211	277	255	351	1313
2020	3	0	0	31	145	168	246	237	316	1143
2020	4	0	0	30	134	153	223	212	268	1020
2020	5	0	0	35	131	174	247	231	306	1124
2020	6	0	0	21	137	174	233	228	318	1111
2020	7	0	0	20	103	140	215	195	279	952
2020	8	0	0	22	104	142	214	203	284	969
2021	target	150	150	248	206	163	184	161	238	1500
2021	1	137	145	239	202	158	180	158	232	1451
2021	2	126	126	139	175	136	177	176	257	1312
2021	3	107	115	174	175	136	149	128	187	1171
2021	4	99	104	112	157	110	125	142	213	1062
2021	5	81	95	94	127	89	132	127	197	942
2021	6	87	75	110	151	126	118	95	92	854
2021	7	73	67	64	126	85	93	115	143	766
2021	8	68	76	80	133	102	123	78	103	763
2021	9	49	67	78	117	94	86	61	64	616
2021	10	40	65	65	104	89	96	71	31	561
2021	11	116	154	195	205	160	198	167	260	1455
2021	12	107	133	178	194	155	160	142	208	1277
2021	13	96	115	124	163	121	154	152	233	1158
2021	14	84	98	85	143	117	134	140	228	1029
2021	15	77	86	84	135	117	123	113	188	923
2021	16	71	74	50	117	98	117	112	192	831
2021	17	59	82	48	115	103	130	119	106	762
2021	18	55	69	41	90	65	72	105	181	678
2021	19	64	56	43	90	79	73	70	201	676
2021	20	47	81	41	86	84	100	87	160	686

**Table S1.** Number of participants per survey round of study series in 2020 and 2021 in eight age groups. The target number of participants was to be achieved in round 1 in 2020, and in rounds 1 and 11 in 2021.




**Figure S1.** Number of participants by survey round and participant round (i.e. the  $n^{\text{th}}$  time a participant participated). Series 2020 (left) consisted of 8 rounds, series 2021 (right) consisted of 20 rounds. Most participants participated for the first time the start of the series, but some also started during the study period. The panel was supplemented to the target population in round 11 of series 2021.

series	age_group	Variable	Estimate	Std. Error	t value	Pr(> t )
2020	18-64	(Intercept)	-0.24	0.06	-4.15	0.00
2020	18-64	part_vaccTRUE	0.00	0.00		
2020	18-64	part_elevated_riskTRUE	-0.03	0.08	-0.38	0.70
2020	18-64	weekendTRUE	-0.21	0.06	-3.65	0.00
2020	18-64	holidayTRUE	0.07	0.10	0.69	0.49
2020	18-64	Deviance explained	0.16			
2020	65+	(Intercept)	0.03	0.13	0.27	0.79
2020	65+	part_vaccTRUE	0.00	0.00		
2020	65+	part_elevated_riskTRUE	0.25	0.12	2.05	0.04
2020	65+	weekendTRUE	-0.32	0.08	-3.89	0.00
2020	65+	holidayTRUE	-0.08	0.11	-0.70	0.48
2020	65+	Deviance explained	0.14			
2021	0-17	(Intercept)	-0.82	0.08	-10.02	0.00
2021	0-17	part_vaccTRUE	0.12	0.19	0.63	0.53
2021	0-17	part_elevated_riskTRUE	-0.14	0.28	-0.52	0.60
2021	0-17	weekendTRUE	-0.67	0.08	-8.35	0.00
2021	0-17	holidayTRUE	-0.20	0.10	-2.02	0.04
2021	0-17	Deviance explained	0.17			
2021	18-64	(Intercept)	-1.04	0.06	-17.68	0.00
2021	18-64	part_vaccTRUE	0.09	0.07	1.24	0.22
2021	18-64	part_elevated_riskTRUE	-0.09	0.08	-1.20	0.23
2021	18-64	weekendTRUE	-0.27	0.05	-5.95	0.00
2021	18-64	holidayTRUE	0.06	0.06	0.94	0.35
2021	18-64	Deviance explained	0.21			
2021	65+	(Intercept)	-0.86	0.15	-5.74	0.00
2021	65+	part_vaccTRUE	0.07	0.19	0.38	0.70
2021	65+	part_elevated_riskTRUE	-0.09	0.11	-0.81	0.42
2021	65+	weekendTRUE	-0.45	0.08	-5.69	0.00
2021	65+	holidayTRUE	0.04	0.10	0.40	0.69
2021	65+	Deviance explained	0.16			

**Table S2.** Results for fixed effects of generalised additive model, by series (2020, 2021) and age group (0-17, 16-64, 65+). Included fixed effects are participant vaccination status (part\_vacc), participant risk status (part\_elevated\_risk), weekend and holiday. Holidays include general holidays and school holidays. The last row of each section refers to the explained deviance.





**Figure S2.** Results for non-stationary fixed effects of generalised additive model, by series (2020, 2021) and age group (0-17, 16-64, 65+). Calendar time (expressed as days since first survey), participant age, and participant round (i.e. the  $n^{\text{th}}$  time a participant participated) are included as cubic splines. Shown are the fitted splines (solid lines)  $\pm$  the standard error (dashed lines).