



1500 series

CS & IT ENGINEERING

Discrete Mathematics



Lecture No.- 01

By- Satish Yadav Sir

Topics to be Covered



Topic

**Inclusion - Exclusion
Question Discussion**



Inclusion-Exclusion :

↓
Venn diagram
11-18


→ Derangement

→ no int soltn.


$$x_1 + x_2 + x_3 = 20$$

→ $(I + E) \rightarrow$ permutatn with
identical objects.

→ Graph Theory.



In a survey of students at State University the following information was obtained: 260 were taking a statistics course, 208 were taking a mathematics course, 160 were taking a computer programming course, 76 were taking statistics and mathematics, 48 were taking statistics and computer programming, 62 were taking mathematics courses and computer programming, 30 were taking all 3 kinds of courses, and 150 were taking none of the 3 courses.



Total Students.

$$\begin{aligned}
 S \cap M &= 76 \\
 (S \cap M \cap \bar{P}) \\
 &= (S \cap M) - (S \cap M \cap P) \\
 &= 76 - 30 = 46
 \end{aligned}$$

- a) How many students were surveyed? $U = (\text{sum up}) +$
 (b) How many students were taking a statistics and
 a mathematics course but not a computer sum up
 programming course? (snpm)
 (c) How many were taking a statistics and a
 computer course but not a mathematics course?

c)
 a) 46
 ✓ b) 18
 c) 32
 d) not a.

b)
 a) 46 ✓
 b) 18
 c) 32
 d) not

A) a) 600
 b) 622 ✓
 c) 390
 d) 310

(d) How many were taking a computer programming and a mathematics course but not a statistics course?

(e) How many were taking a statistics course but not taking a course in mathematics or in computer programming?

(f) How many were taking a mathematics course but not taking a statistics course or a computer programming course?

(g) How many were taking a computer programming course but not taking a course in mathematics or in statistics

g) 80 ✓
90
100
110

f) 100 ✓
200
300
400

e) 160
166 ✓
120
130

d) 18
b) 36
c) 32 ✓
d) not a.

In how many ways can the letters {5. a, 4.b 3.c} be arranged so that all the letters of the same kind are not in a single block?

Total possibility = $\frac{12!}{5! \cdot 4! \cdot 3!} - (A_1 \cup A_2 \cup A_3)$ { $\overset{5}{\boxed{aaaaa}} \overset{4}{\boxed{bbbb}} \overset{3}{\boxed{ccc}}$ }

A₁ → all a's are in same block.

$\overset{1}{\boxed{aaaaa}} \overset{4}{\boxed{bbbb}} \overset{3}{\boxed{ccc}}$
 $\frac{8!}{4! \cdot 3!}$

A₂ → all b's will be in same block.

$\overset{5}{\boxed{aaaaa}} \overset{1}{\boxed{bbbb}} \overset{3}{\boxed{ccc}}$
 $\frac{9!}{5! \cdot 3!}$

Total

-(at least 1 block) $\frac{10!}{5! \cdot 4!}$

A₃

→ all c will be in same block

$\boxed{aaaaa} \boxed{bbbb} \boxed{ccc}$

$$\text{Ans} = \text{Total} - (A_1 \cup A_2 \cup A_3)$$

aaaaa bbbb ccc.

$$A_1 \cup A_2 \cup A_3 = A_1 + A_2 + A_3 - A_1 \cap A_2 - A_2 \cap A_3 - A_1 \cap A_3 + A_1 \cap A_2 \cap A_3.$$

$$A_1 \rightarrow \overset{1}{\boxed{\text{aaaaa}}} \overset{4}{\text{bbbb}} \overset{3}{\text{ccc}} \quad \frac{8!}{4! \cdot 3!}$$

$$A_1 \cap A_2 \rightarrow \overset{1}{\boxed{\text{aaaa}}} \overset{1}{\boxed{\text{bbbb}}} \overset{3}{\text{ccc}} \quad \frac{5!}{3!}$$

$$A_2 \rightarrow \overset{5}{\text{aaaaa}} \overset{1}{\boxed{\text{bbbb}}} \overset{3}{\text{ccc}} \quad \frac{9!}{5! \cdot 3!}$$

$$A_1 \cap A_3 \rightarrow \overset{1}{\boxed{5.a}} \overset{4}{\text{bbbb}} \overset{1}{\boxed{3.c}} \quad \frac{6!}{4!}$$

$$A_3 \rightarrow \overset{5}{\text{aaaaa}} \overset{4}{\text{bbbb}} \overset{1}{\boxed{\text{ccc}}} \quad \frac{7!}{5!}$$

$$A_2 \cap A_3 \rightarrow \overset{5}{\text{aaaaa}} \overset{1}{\boxed{4b}} \overset{1}{\boxed{3c}} \quad \frac{7!}{5!}$$

$$A_1 \cap A_2 \cap A_3 \rightarrow \overset{1}{\boxed{3a}} \overset{1}{\boxed{4b}} \overset{1}{\boxed{3c}} \quad 3!$$

$$\frac{10!}{5! \cdot 4!}$$

how many ways $\{4.a, 3.b, 2.c\}$ be arranged
so that all letters of the same kind should
not be in single
block

A) 870

B) 871

C) 872

d) 873

$$x_1 + x_2 + x_3 + x_4 = 18 \quad x_i \leq 7$$

Count the number of integral solutions _____

$$x_1 + x_2 + x_3 + x_4 = 18 \quad x_i \leq 7$$


$$x_1 + x_2 + x_3 + x_4 = 20 \quad x_i \leq 7$$

at least: $x_i \geq 2$ (Combi with Reptn)

at least/at most \rightarrow G.F

at most \rightarrow I.F

$$= \text{Total} - (A_1 \cup A_2 \cup A_3 \cup A_4)$$



$$\left\{ x_1 + x_2 + x_3 + x_4 = 20 \mid \underline{x_i \geq 8} \right\}$$

$$x_1 + x_2 + x_3 = 20$$

$$\begin{aligned} 2 &\leq x_1 \leq 5 \\ 4 &\leq x_2 \leq 7 \\ -2 &\leq x_3 \leq 9 \end{aligned}$$

Total:

$$x_1 + x_2 + x_3 = 20$$

$$\begin{aligned} \downarrow & \quad \downarrow & \quad \downarrow \\ x_1 &\geq 2, & x_2 &\geq 4, & x_3 &\geq -2 \text{ (Initial)} \end{aligned}$$

$$\text{C.B.R.} \quad 20 - 2 - 4 + 2 + (3-1)$$

$$\text{line: } (3-1)$$

$$3-1$$

Required

$$\text{ans} = \text{Total} - (A_1 \cup A_2 \cup A_3)$$

$$A_1 \rightarrow \{ x_1 + x_2 + x_3 = 20 \mid \underline{x_1 \geq 6}, x_2 \geq 4, x_3 \geq -2 \}$$

$$20 - 6 - 4 + 2 + (3-1) \binom{14}{3-1} C_2$$

$$A_2 \rightarrow \{ x_1 + x_2 + x_3 = 20 \mid x_2 \geq 8, x_1 \geq 2, x_3 \geq -2 \}$$

$$A_3 \rightarrow \{ x_1 + x_2 + x_3 = 20 \mid x_3 \geq 10, x_1 \geq 2, x_2 \geq 4 \}$$

$$A_1 \cap A_2 = \begin{cases} x_1 + x_2 + x_3 = 20 \\ x_1 \geq 6 \\ x_2 \geq 8 \\ x_3 \geq -2 \end{cases}$$

Required
Ans:



2	4	-2
3	5	-1
4	6	0
5	7	1
		2
		3
		4
		5
		6
		7
		8
		9
		10

A₁

A₂

A₃

Total



$$x_1 + x_2 = 20$$

1	2
20	20

There are 5 villages in country side , an engineer is to devise a system of two- way roads so that after the system is completed , no village will be isolated , in how many ways can he do this ?

$$\text{Total} - (A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_5)$$

$$= 2^{10} - \left(5C_1 \times 2^{\frac{4 \times 3}{2}} \right.$$



Total paths.
= Total Graphs.
 $= 2^{\frac{5 \times 4}{2}} = 2^{10}$

A) 450

B) 340

C) 768 ✓

D) 560

$$- 5C_2 \times 2^{\frac{3 \times 2}{2}} + 5C_3 \times 2^2 - 5C_4 \times 2^{\frac{1 \times 0}{2}} + 5C_5 \times 2^0$$

✓
A — B



A •



X X

Find the number of derangements of the integers from 1 to 10 inclusive, satisfying the condition that the set of elements in the first 5 places is

(a) 1, 2, 3, 4, 5, in some order, $ANS : D_5 : D_5 = 1936$.

(b) 6, 7, 8, 9, 10, in some order. $(5!)^2 = 14,400$

4. An advertising agency has 1,000 clients. Suppose that T is the set of clients that use television advertising, R is the set of clients that use radio advertising, and N is the set of clients who use newspaper advertising. Suppose that $|T| = 415$, $|R| = 350$, $|N| = 280$, 100

clients use all 3 types of advertising, 175 use television and radio, 180 use radio and newspapers, and $|T \cap N| = 165$.

(a) Find $|T \cap R \cap N|$.

(b) How many clients use radio and newspaper advertising but not television?

(c) How many use television but do not use newspaper advertising and do not use radio advertising?

(d) Find $|\overline{T} \cap \overline{R} \cap \overline{N}|$.

4. (a) 75.

(b) $|R \cap N \cap \overline{T}| = 80$

(c) $|T \cap \overline{N} \cap \overline{R}| = 175$.

(d) $|\overline{T} \cap \overline{R} \cap \overline{N}| = |\overline{T \cup R \cup N}| = 1,000 - 625 = 375$.

10. Find the number of permutations of the integers 1 to 10 inclusive
- (a) such that exactly 4 of the integers are in their natural positions (that is, exactly 6 of the integers are deranged)
 - (b) such that 6 or more of the integers are deranged.
 - (c) that do not have 1 in the first place, nor 4 in the fourth place, nor 7 in the seventh place.
 - (d) such that no odd integer will be in the natural position.
 - (e) that do not begin with a 1 and do not end with 10.

10. (a) $C(10,6)D_6$.

(b) $\binom{10}{6}D_6 + \binom{10}{7}D_7 + \binom{10}{8}D_8 + \binom{10}{9}D_9 + \binom{10}{10}D_{10}$

(c) $10! - (3)9! + (3)8! - 7!$.

(d) $10! - \binom{5}{1}9! + \binom{5}{2}8! - \binom{5}{3}7! + \binom{5}{4}6! - \binom{5}{5}5!$.

(e) $10! - (2)9! + 8!$.

17. At a theater 10 men check their hats. In how many ways can their hats be returned so that

- (a) no man receives his own hat?
- (b) at least 1 of the men receives his own hat?
- (c) at least 2 of the men receive their own hats?

17. (a) D_{10} .

(b) $10! - D_{10}$.

(c) $10! - D_{10} - 10D_9$.

25. The squares of a chessboard are painted 8 different colors. The squares of each row are painted all 8 colors and no 2 consecutive squares in one column can be painted the same color. In how many ways can this be done?

25. The first row can be painted $8!$ ways. Each row after the first can be painted D_8 ways. Hence the number of ways is $8!(D_8)^7$.

5. Determine the number of positive integers n , $1 \leq n \leq 2000$, that are

- a) not divisible by 2, 3, or 5
- b) not divisible by 2, 3, 5, or 7
- c) not divisible by 2, 3, or 5, but are divisible by 7

- (a) c_1 : number n is divisible by 2
 c_2 : number n is divisible by 3
 c_3 : number n is divisible by 5
 $N(c_1) = \lfloor 2000/2 \rfloor = 1000$, $N(c_2) = \lfloor 2000/3 \rfloor = 666$,
 $N(c_3) = \lfloor 2000/5 \rfloor = 400$, $N(c_1 c_2) = \lfloor 2000/(2)(3) \rfloor = 333$,
 $N(c_2 c_3) = \lfloor 2000/(3)(5) \rfloor = 133$, $N(c_1 c_3) = \lfloor 2000/(2)(5) \rfloor = 200$,
 $N(c_1 c_2 c_3) = \lfloor 2000/(2)(3)(5) \rfloor = 66$.
 $N(\bar{c}_1 \bar{c}_2 \bar{c}_3) = 2000 - (1000 + 666 + 400) + (333 + 200 + 133) - 66 = 534$
- (b) Let c_1, c_2, c_3 be as in part (a). Let c_4 denote the number n is divisible by 7. Then
 $N(c_4) = 285$, $N(c_1 c_4) = 142$, $N(c_2 c_4) = 95$, $N(c_3 c_4) = 57$, $N(c_1 c_2 c_4) = 47$, $N(c_1 c_3 c_4) = 28$, $N(c_2 c_3 c_4) = 19$, $N(c_1 c_2 c_3 c_4) = 9$. $N(\bar{c}_1 \bar{c}_2 \bar{c}_3 \bar{c}_4) = 2000 - (1000 + 666 + 400 + 285) + (333 + 200 + 133 + 142 + 95 + 57) - (66 + 47 + 28 + 19) + 9 = 458$
- (c) $534 - 458 = 76$.
2. a) List all the derangements of 1, 2, 3, 4, 5 where the first three numbers are 1, 2, and 3, in some order.
b) List all the derangements of 1, 2, 3, 4, 5, 6 where the first three numbers are 1, 2, and 3, in some order.
3. How many derangements are there for 1, 2, 3, 4, 5?
4. How many permutations of 1, 2, 3, 4, 5, 6, 7 are not derangements?
5. a) Let $A = \{1, 2, 3, \dots, 7\}$. A function $f: A \rightarrow A$ is said to have a *fixed point* if for some $x \in A$, $f(x) = x$. How many one-to-one functions $f: A \rightarrow A$ have at least one fixed point?
b) In how many ways can we devise a secret code by assigning to each letter of the alphabet a different letter to represent it?
2. (a) There are only two derangements with this property: 23154 and 31254.
(b) Here there are four such derangements:
(i) 231546 (ii) 231645 (iii) 312546 (iv) 312645
3. The number of derangements for 1,2,3,4,5 is $5![1 - 1 + (1/2!) - (1/3!) + (1/4!) - (1/5!)] = 5![(1/2!) - (1/3!) + (1/4!) - (1/5!)] = (5)(4)(3) - (5)(4) + 5 - 1 = 60 - 20 + 5 - 1 = 44$.
4. There are $7! = 5040$ permutations of 1,2,3,4,5,6,7. Among these there are $7![1 - 1 + (1/2!) - (1/3!) + (1/4!) - (1/5!) + (1/6!) - (1/7!)] = 1854$ derangements. Consequently, we have $5040 - 1854 = 3186$ permutations of 1,2,3,4,5,6,7 that are not derangements.
5. (a) $7! - d_7$ ($d_7 \doteq (7!)e^{-1}$); (b) $d_{26} \doteq (26!)e^{-1}$
6. How many derangements of 1, 2, 3, 4, 5, 6, 7, 8 start with
(a) 1, 2, 3, and 4, in some order? (b) 5, 6, 7, and 8, in some order?
7. For the positive integers 1, 2, 3, \dots , $n - 1$, n , there are 11,660 derangements where 1, 2, 3, 4, and 5 appear in the first five positions. What is the value of n ?

6. (a) There are $(d_4)^2 = 9^2 = 81$ such derangements.
 (b) In this case we get $(4!)^2 = 24^2 = 576$ derangements.
7. Let $n = 5 + m$. Then $11,660 = d_5 \cdot d_m = 44(d_m)$, and so $d_m = 265 = d_6$. Consequently, $n = 11$.

9. In how many ways can Mrs. Ford distribute ten distinct books to her ten children (one book to each child) and then collect and redistribute the books so that each child has the opportunity to peruse two different books?

$$9. \quad (10!)d_{10} \doteq (10!)^2(e^{-1})$$

12. Ms. Pezzulo teaches geometry and then biology to a class of 12 advanced students in a classroom that has only 12 desks. In how many ways can she assign the students to these desks so that (a) no student is seated at the same desk for both classes? (b) there are exactly six students each of whom occupies the same desk for both classes?

$$12. \quad (a) \quad (12!)d_{12} \qquad (b) \quad (12!)\binom{12}{6}d_6$$

1. Determine how many $n \in \mathbb{Z}^+$ satisfy $n \leq 500$ and are not divisible by 2, 3, 5, 6, 8, or 10.

1. We need only consider the divisors 2, 3, and 5. Let c_1 denote divisibility by 2, c_2 divisibility by 3, and c_3 divisibility by 5.

$$N = 500; \quad N(c_1) = \lfloor 500/2 \rfloor = 250; \quad N(c_2) = \lfloor 500/3 \rfloor = 166; \quad N(c_3) = \lfloor 500/5 \rfloor = 100; \\ N(c_1c_2) = \lfloor 500/6 \rfloor = 83; \quad N(c_1c_3) = \lfloor 500/10 \rfloor = 50; \quad N(c_2c_3) = \lfloor 500/15 \rfloor = 33; \\ N(c_1c_2c_3) = \lfloor 500/30 \rfloor = 16.$$

$$N(\bar{c}_1\bar{c}_2\bar{c}_3) = 500 - (250 + 166 + 100) + (83 + 50 + 33) - 16 = 134.$$

19. Caitlyn has 48 different books: 12 each in mathematics, chemistry, physics, and computer science. These books are ar-

ranged on four shelves in her office with all books on any one subject on its own shelf. When her office is cleaned, the 48 books are taken down and then replaced on the shelves — once again with all 12 books on any one subject on its own shelf. In how many ways can this be done so that (a) no subject is on its original shelf? (b) one subject is on its original shelf? (c) no subject is on its original shelf and no book is in its original position? [For example, the book originally in the third (from the left) position on the first shelf must not be replaced on the first shelf and must not be in the third (from the left) position on the shelf where it is placed.]

19. a) $d_4(12!)^4$
 b) $\binom{4}{1}d_3(12!)^4$
 c) $d_4(d_{12})^4$

4. Annually, the 65 members of the maintenance staff sponsor a "Christmas in July" picnic for the 400 summer employees at their company. For these 65 people, 21 bring hot dogs, 35 bring fried chicken, 28 bring salads, 32 bring desserts, 13 bring hot dogs and fried chicken, 10 bring hot dogs and salads, 9 bring hot dogs and desserts, 12 bring fried chicken and salads, 17 bring fried chicken and desserts, 14 bring salads and desserts, 4 bring hot dogs, fried chicken, and salads, 6 bring hot dogs, fried chicken, and desserts, 5 bring hot dogs, salads, and desserts, 7 bring fried chicken, salads, and desserts, and 2 bring all four food items. Those (of the 65) who do not bring any of these four food items are responsible for setting up and cleaning up for the picnic. How many of the 65 maintenance staff will (a) help to set up and clean up for the picnic? (b) bring only hot dogs? (c) bring exactly one food item?

c_1 : Staff member brings hot dogs
 c_2 : Staff member brings fried chicken
 c_3 : Staff member brings salads
 c_4 : Staff member brings desserts
 $N = 65$

$N(c_1) = 21$; $N(c_2) = 35$; $N(c_3) = 28$; $N(c_4) = 32$
 $N(c_1c_2) = 13$; $N(c_1c_3) = 10$; $N(c_1c_4) = 9$; $N(c_2c_3) = 12$; $N(c_2c_4) = 17$; $N(c_3c_4) = 14$
 $N(c_1c_2c_3) = 4$; $N(c_1c_2c_4) = 6$; $N(c_1c_3c_4) = 5$; $N(c_2c_3c_4) = 7$
 $N(c_1c_2c_3c_4) = 2$.

(a) $N(\bar{c}_1\bar{c}_2\bar{c}_3\bar{c}_4) = 65 - [21 + 35 + 28 + 32] + [13 + 10 + 9 + 12 + 17 + 14] - [4 + 6 + 5 + 7] + 2 = 65 - 116 + 75 - 22 + 2 = 4$.

(b) $N(\bar{c}_2\bar{c}_3\bar{c}_4) = N - [N(c_2) + N(c_3) + N(c_4)] + [N(c_2c_3) + N(c_2c_4) + N(c_3c_4)] - N(c_2c_3c_4)$, so
 $N(\bar{c}_2\bar{c}_3\bar{c}_4) = N(c_1) - [N(c_1c_2) + N(c_1c_3) + N(c_1c_4)] + [N(c_1c_2c_3) + N(c_1c_2c_4) + N(c_1c_3c_4)] - N(c_1c_2c_3c_4) = 21 - [13 + 10 + 9] + [4 + 6 + 5] - 2 = 21 - 32 + 15 - 2 = 2$.

(c) $N(\bar{c}_1c_2\bar{c}_3\bar{c}_4) = N(c_2) - [N(c_1c_2) + N(c_2c_3) + N(c_2c_4)] + [N(c_1c_2c_3) + N(c_1c_2c_4) + N(c_2c_3c_4)] - N(c_1c_2c_3c_4) = 35 - [13 + 12 + 17] + [4 + 6 + 7] - 2 = 35 - 42 + 17 - 2 = 8$

$N(\bar{c}_1\bar{c}_2c_3\bar{c}_4) = N(c_3) - [N(c_1c_3) + N(c_2c_3) + N(c_3c_4)] + [N(c_1c_2c_3) + N(c_1c_3c_4) + N(c_2c_3c_4)] - N(c_1c_2c_3c_4) = 28 - [10 + 12 + 14] + [4 + 5 + 7] - 2 = 28 - 36 + 16 - 2 = 6$.
 $N(\bar{c}_1\bar{c}_2\bar{c}_3c_4) = N(c_4) - [N(c_1c_4) + N(c_2c_4) + N(c_3c_4)] + [N(c_1c_2c_4) + N(c_1c_3c_4) + N(c_2c_3c_4)] - N(c_1c_2c_3c_4) = 32 - [9 + 17 + 14] + [6 + 5 + 7] - 2 = 32 - 40 + 18 - 2 = 8$.
 So the answer is $2 + 8 + 6 + 8 = 24$.

