

3. PAIR OF LINEAR EQUATION IN TWO VARIABLES

Marks: 20

MCQ's TEST – 1

Time: 25 min



1. The pair of equations $y = 0$ and $y = -7$ has
(a) one solution (b) two solutions (c) infinitely many solutions (d) no solution
 2. The pair of equations $x = a$ and $y = b$ graphically represents the lines which are
(a) parallel (b) intersecting at (a, b) (c) coincident (d) intersecting at (b, a)
 3. The value of c for which the pair of equations $cx - y = 2$ and $6x - 2y = 3$ will have infinitely many solutions is
(a) 3 (b) -3 (c) -12 (d) no value
 4. When lines l_1 and l_2 are coincident, then the graphical solution system of linear equation has
(a) infinite number of solutions (b) unique solution (c) no solution
(d) one solution
 5. When lines l_1 and l_2 are parallel, then the graphical solution system of linear equation has
(a) infinite number of solutions (b) unique solution (c) no solution
(d) one solution
 6. The coordinates of the vertices of triangle formed between the lines and y-axis from the graph is
(a) $(0, 5)$, $(0, 0)$ and $(6.5, 0)$
(b) $(4, 2)$, $(0, 0)$ and $(6.5, 0)$
(c) $(4, 2)$, $(0, 0)$ and $(0, 5)$
(d) none of these
-
7. Five years ago Nuri was thrice old as Sonu. Ten years later, Nuri will be twice as old as Sonu. The present age, in years, of Nuri and Sonu are respectively
(a) 50 and 20 (b) 60 and 30 (c) 70 and 40 (d) 40 and 10
 8. The pair of equations $5x - 15y = 8$ and $3x - 9y = 24/5$ has
(a) infinite number of solutions (b) unique solution (c) no solution
(d) one solution
 9. The pair of equations $x + 2y + 5 = 0$ and $-3x - 6y + 1 = 0$ have
(a) infinite number of solutions (b) unique solution (c) no solution
(d) one solution
 10. The sum of the digits of a two-digit number is 9. If 27 is added to it, the digits of the number get reversed. The number is
(a) 36 (b) 72 (c) 63 (d) 25

11. If a pair of equation is consistent, then the lines will be
 (a) parallel (b) always coincident (c) always intersecting
 (d) intersecting or coincident
12. The solution of the equations $x + y = 14$ and $x - y = 4$ is
 (a) $x = 9$ and $y = 5$ (b) $x = 5$ and $y = 9$ (c) $x = 7$ and $y = 7$ (d) $x = 10$ and $y = 4$
13. The sum of the numerator and denominator of a fraction is 12. If the denominator is increased by 3, the fraction becomes $\frac{1}{2}$, then the fraction is
 (a) $\frac{4}{7}$ (b) $\frac{5}{7}$ (c) $\frac{6}{7}$ (d) $\frac{3}{7}$
14. The value of k for which the system of equations $x - 2y = 3$ and $3x + ky = 1$ has a unique solution is
 (a) $k = -6$ (b) $k \neq -6$ (c) $k = 0$ (d) no value
15. If a pair of equation is inconsistent, then the lines will be
 (a) parallel (b) always coincident (c) always intersecting
 (d) intersecting or coincident
16. The value of k for which the system of equations $2x + 3y = 5$ and $4x + ky = 10$ has infinite many solutions is
 (a) $k = -3$ (b) $k \neq -3$ (c) $k = 0$ (d) none of these
17. The value of k for which the system of equations $kx - y = 2$ and $6x - 2y = 3$ has a unique solution is
 (a) $k = -3$ (b) $k \neq -3$ (c) $k = 0$ (d) $k \neq 0$
18. The solution of the equations $0.4x + 0.3y = 1.7$ and $0.7x - 0.2y = 0.8$ is
 (a) $x = 1$ and $y = 2$ (b) $x = 2$ and $y = 3$ (c) $x = 3$ and $y = 4$ (d) $x = 5$ and $y = 4$
19. The value of k for which the system of equations $x + 2y = 3$ and $5x + ky + 7 = 0$ has no solution is
 (a) 10 (b) 6 (c) 3 (d) 1
19. The value of k for which the system of equations $3x + 5y = 0$ and $kx + 10y = 0$ has a non-zero solution is
 (a) 0 (b) 2 (c) 6 (d) 8
20. The sum of the digits of a two – digit number is 12. The number obtained by interchanging its digits exceeds the given number is.....
 (a) 72 (b) 75 (c) 57 (d) None of these