

Every Algebraic Number is a Critical Value of an Integer Polynomial

by RJ Acuña

Abstract

For every irreducible $f \in \mathbb{Z}[x]$ of positive degree we construct $g \in \mathbb{Z}[x]$ and an irreducible $F \in \mathbb{Z}[x]$ with $F \mid g'$ and $F^2 \mid f \circ g$. Equivalently, every algebraic number is a critical value of an integer polynomial, so the set of critical values of $\mathbb{Z}[x]$ is exactly $\overline{\mathbb{Q}}$. The construction is explicit. The proof was assisted by AI, and has been formalized in Lean.

1 Introduction

A critical value of a nonconstant $g \in \mathbb{Z}[x]$ is a number α with $g(\beta) = \alpha$ and $g'(\beta) = 0$ for some β . Since $\overline{\mathbb{Q}}$ is closed under polynomial evaluation, every critical value is algebraic. This note proves the converse, and does so with an explicit construction.

Theorem 1. *Let f be an irreducible integral polynomial of positive degree. Then there exist F and g in $\mathbb{Z}[x]$, with F irreducible and of positive degree, such that*

$$F \mid g' \quad \text{and} \quad F^2 \mid f \circ g.$$

The corollary is the statement in the title.

Corollary 1. *The set of critical values is exactly $\overline{\mathbb{Q}}$.*

2 The construction

Let f be irreducible of degree $n \geq 1$, with root α ; write $f(x) = \sum_{k=0}^n a_k x^{n-k}$, so that $a_n = f(0)$. If $f = x$, then $g = x^2$ and $F = x$ work. Otherwise $f(0) \neq 0$, and since $\gcd(f, f') = 1$ in $\mathbb{Q}[x]$, the elimination ideal $(f, f') \cap \mathbb{Z}$ is nonzero; let $u, v \in \mathbb{Z}[x]$ and $\rho \in \mathbb{Z}$, $\rho \neq 0$, be such that

$$u f + v f' = \rho,$$

and set

$$M = \rho a_n, \quad F(x) = \frac{f(Mx)}{a_n} = \sum_{k=0}^n A_k x^{n-k}, \quad W(x) = \frac{v(0)F(x) - v(Mx)}{\rho}.$$

The polynomial claimed by Theorem 1 is

$$g(x) = Mx + f(Mx)W(x), \tag{1}$$

together with this F .

Theorem (restatement of Theorem 1). *F is irreducible, and $F \mid g'$ and $F^2 \mid f \circ g$.*

pf. From $A_k = a_k M^{n-k}/a_n$ we get $A_n = 1$ and $A_k = a_k \rho^{n-k} a_n^{n-k-1}$ for $k < n$, so $F \in \mathbb{Z}[x]$ with $F(0) = 1$, which gives $W(0) = 0$. And ρ divides M and every A_k with $k < n$, so it divides $v(0)F(x) - v(Mx)$ and $W \in \mathbb{Z}[x]$. Therefore g is an integral polynomial.

Let $\beta = \alpha/M$, a root of F . Then $W(\beta) = -v(\alpha)/\rho = -1/f'(\alpha)$, so

$$g(\beta) = \alpha, \quad g'(\beta) = M(1 + f'(\alpha)W(\beta)) = 0.$$

Hence β is a double root of $f \circ g$, and F is primitive irreducible, so $F \mid g'$ and $F^2 \mid f \circ g$ by Gauss. ■

The computation uses $f(\alpha) = 0$ and nothing else about α , so it runs at every root of f at once: all n conjugates are critical values of the same g , at the n points α_i/M . That is what makes F^2 , and not merely $(x - \beta)^2$, divide $f \circ g$.

Any ρ will do, but $M = \rho a_n$ multiplies into every coefficient of F , W and g , so the smallest gives the smallest answer: the generator of $(f, f') \cap \mathbb{Z}$, the reduced resultant, of which $\text{Res}(f, f')$ is a multiple. On $f = x^2 + 1$ the generator is 2 against a resultant of 4, and on Dedekind's cubic below 1006 against 2012.

3 Examples

Each line below is a closed identity, checkable by expansion.

Example 1 ($f = qx - p$, $p \neq 0$, $\gcd(p, q) = 1$). Here $f' = q$ and $p = qx - f$, so $(f, f') \cap \mathbb{Z} = \mathbb{Z}$ and $\rho = 1$. Choose $a, b \in \mathbb{Z}$ with $ap + bq = 1$ and take $u = -a$, $v = ax + b$; then $f(0) = -p$, $M = -p$, $F = qx + 1$, $W = x$, and

$$g = -pqx^2 - 2px, \quad g' = -2p(qx + 1), \quad f \circ g = -p(qx + 1)^2.$$

Example 2 ($f = x^2 + 1$). With $u = 2$, $v = -x$, $\rho = 2$, $f(0) = 1$, $M = 2$, $F = 4x^2 + 1$ and $W = x$,

$$g = 4x^3 + 3x, \quad g' = 3(4x^2 + 1), \quad f \circ g = (4x^2 + 1)^2(x^2 + 1).$$

Here $\deg g = 3 = n + 1$, the least degree any solution can have. The unit group $\mathcal{O}_K^\times = \{\pm 1, \pm i\}$ has rank 0, and it is $\mathbb{Z}[i/2] = \mathbb{Z}[i][1/2]$ that supplies the needed unit $1 + i$.

Example 3 ($f = x^3 - 2$). With $u = 3$, $v = -x$, $\rho = -6$, $f(0) = -2$, $M = 12$, $F = 1 - 864x^3$ and $W = -2x$,

$$g = -3456x^4 + 16x, \quad g' = 16(1 - 864x^3).$$

Example 4 ($f = x^3 - x - 1$). The construction gives $M = 23$ and a sextic. A hand-tuned quintic also witnesses the *statement*, and is a useful independent check of it:

$$g = -24334x^5 + 39675x^4 - 26450x^3 + 9085x^2 - 1610x + 118, \quad F = 23x^3 - 23x^2 + 8x - 1,$$

with $g' = -230(23x - 7)F$ and $F^2 \mid f \circ g$, the cofactor having degree 9.

Example 5 (Dedekind's cubic). $f = x^3 - x^2 - 2x - 8$, the standard example of a field whose ring of integers is not monogenic: 2 is a common index divisor [Ded78]. With $u = 21x - 125$, $v = -7x^2 + 44x - 3$, $\rho = 1006$, $f(0) = -8$, $M = -8048$,

$$F = 65158925824x^3 + 8096288x^2 - 2012x + 1, \quad W = -194310912x^3 + 426544x^2 + 358x,$$

$$g = 101288722414414331904x^6 - 209759614012620800x^5 \\ - 217370176548864x^4 - 14767629312x^3 + 2350016x^2 - 10912x.$$

4 Formalization

The development above has been formalized in Lean 4 [MU21] against Mathlib [The20], in about 1,200 lines. The main statement is

```
1 theorem main (f : Z[X]) (hf : 1 <= f.natDegree) :
2   exists g H : Z[X], Irreducible H /\ H \mid derivative g /\ H^2 \mid f.comp g
```

with a strengthened form carrying $2 \leq \deg g$ and $1 \leq \deg H$, and with Corollary 1 proved from it. Every result depends only on the three standard axioms `propext`, `Classical.choice` and `Quot.sound`.

References

- [Ded78] Richard Dedekind. “Über den Zusammenhang zwischen der Theorie der Ideale und der Theorie der höheren Kongruenzen”. In: *Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen* (1878).
- [The20] The mathlib Community. “The Lean Mathematical Library”. In: *Certified Programs and Proofs (CPP 2020)*. ACM, 2020.
- [MU21] Leonardo de Moura and Sebastian Ullrich. “The Lean 4 Theorem Prover and Programming Language”. In: *Automated Deduction – CADE 28*. Lecture Notes in Computer Science. Springer, 2021.