

Every Algebraic Number is a Critical Value of an Integer Polynomial

by RJ Acuña

Abstract

For every nonconstant $f \in \mathbb{Z}[X]$ we construct $g \in \mathbb{Z}[X]$ and an irreducible $H \in \mathbb{Z}[X]$ with $H \mid g'$ and $H^2 \mid f \circ g$. Equivalently, every algebraic number is a critical value of an integer polynomial, so the set of critical values of $\mathbb{Z}[X]$ is exactly $\overline{\mathbb{Q}}$. The construction is explicit. The proof was assisted by AI, and has been formalized in Lean.

1 Introduction

A critical value of a nonconstant $g \in \mathbb{Z}[X]$ is a number α with $g(\beta) = \alpha$ and $g'(\beta) = 0$ for some β . Since $\overline{\mathbb{Q}}$ is closed under polynomial evaluation, every critical value is algebraic. This note proves the converse, and does so with an explicit construction.

Theorem 1. *Let $f \in \mathbb{Z}[X]$ be nonconstant. Then there exist $g, H \in \mathbb{Z}[X]$ with H irreducible in $\mathbb{Z}[X]$ and $1 \leq \deg H$, such that*

$$H \mid g' \quad \text{and} \quad H^2 \mid f \circ g \quad \text{in } \mathbb{Z}[X].$$

The g produced satisfies $\deg g \geq 2$.

The corollary is the statement in the title.

Corollary 1. *The set of critical values is exactly $\overline{\mathbb{Q}}$.*

2 The construction

Let $h \in \mathbb{Z}[X]$ be a primitive irreducible factor of f , of degree n , with root θ and $h(0) \neq 0$. Put

$$\rho = \text{Res}(h, h') \in \mathbb{Z}, \quad v \in \mathbb{Z}[X], \quad \deg v < n, \quad v h' \equiv \rho \pmod{h},$$

which the extended Euclidean algorithm produces, and set

$$M = \rho h(0), \quad H(X) = \frac{h(MX)}{h(0)}, \quad W(X) = \frac{v(0)H(X) - v(MX)}{\rho}.$$

The polynomial claimed by Theorem 1 is

$$g(X) = MX + h(MX)W(X)$$

together with this H .

Theorem (restatement of Theorem 1). *H is irreducible in $\mathbb{Z}[X]$, and $H \mid g'$ and $H^2 \mid f \circ g$ in $\mathbb{Z}[X]$.*

Verification. Let $\beta = \theta/M$, a root of H . Since $H(0) = 1$, the constant terms in W cancel, and every other coefficient carries M^j with $\rho h(0) \mid M$, so $W \in \mathbb{Z}[X]$ and $W(0) = 0$; likewise $H, g \in \mathbb{Z}[X]$. Then $W(\beta) = -v(\theta)/\rho = -1/h'(\theta)$, so

$$g(\beta) = \theta, \quad g'(\beta) = M(1 + h'(\theta)W(\beta)) = 0.$$

Hence β is a double root of $f \circ g$, and H is primitive irreducible, so $H \mid g'$ and $H^2 \mid f(g)$ in $\mathbb{Z}[X]$ by Gauss.

The computation uses $h(\theta) = 0$ and nothing else about θ , so it runs at every root of h at once: all n conjugates are critical values of the same g , at the n points θ_i/M . That is what makes H^2 , and not merely $(X - \beta)^2$, divide $f \circ g$.

Only the congruence $vh' \equiv \rho \pmod{h}$ is used, so any nonzero ρ admitting one will serve; the resultant is simply the standard choice. Smaller choices give a smaller g , and the best available is the generator of $(h, h') \cap \mathbb{Z}$ — see Example 2, where $\rho = 2$ while $\text{Res}(h, h') = 4$.

3 Examples

Each line below is a closed identity, checkable by expansion.

Example 1 ($f = qX - p, p \neq 0$). With $u = 0, v = 1, \rho = q, h_0 = -p, M = -pq, H = q^2X + 1$ and $W = qX$,

$$g = -pq^3X^2 - 2pqX, \quad g' = -2pq(q^2X + 1), \quad f \circ g = -p(q^2X + 1)^2.$$

Example 2 ($f = X^2 + 1$). With $u = 2, v = -X, \rho = 2, h_0 = 1, M = 2, H = 4X^2 + 1$ and $W = X$,

$$g = 4X^3 + 3X, \quad g' = 3(4X^2 + 1), \quad f \circ g = (4X^2 + 1)^2(X^2 + 1).$$

Here $\deg g = 3 = n + 1$, the least degree any solution can have. Note also $\rho = 2$ while $\text{Res}(h, h') = 4$: the reduced resultant is the better choice. The unit group $\mathcal{O}_K^\times = \{\pm 1, \pm i\}$ has rank 0, and it is $\mathbb{Z}[i/2] = \mathbb{Z}[i][1/2]$ that supplies the needed unit $1 + i$.

Example 3 ($f = X^3 - 2$). With $u = 3, v = -X, \rho = -6, h_0 = -2, M = 12, H = 1 - 864X^3$ and $W = -2X$,

$$g = -3456X^4 + 16X, \quad g' = 16(1 - 864X^3).$$

Example 4 ($f = X^3 - X - 1$). The construction gives $M = 23$ and a sextic. A hand-tuned quintic also witnesses the *statement*, and is a useful independent check of it:

$$g = -24334X^5 + 39675X^4 - 26450X^3 + 9085X^2 - 1610X + 118, \quad H = 23X^3 - 23X^2 + 8X - 1,$$

with $g' = -230(23X - 7)H$ and $H^2 \mid f \circ g$, the cofactor having degree 9.

Example 5 (Dedekind's cubic). $f = X^3 - X^2 - 2X - 8$, the standard example of a field whose ring of integers is not monogenic: 2 is a common index divisor [Ded78]. With $u = 21X - 125, v = -7X^2 + 44X - 3, \rho = 1006, h_0 = -8, M = -8048$,

$$H = 65158925824X^3 + 8096288X^2 - 2012X + 1, \quad W = -194310912X^3 + 426544X^2 + 358X,$$

$$g = 101288722414414331904X^6 - 209759614012620800X^5 \\ - 217370176548864X^4 - 14767629312X^3 + 2350016X^2 - 10912X.$$

4 Formalization

The development above has been formalized in Lean 4 [MU21] against Mathlib [The20], in about 1,200 lines. The main statement is

```
1 theorem main (f : Z[X]) (hf : 1 <= f.natDegree) :
2   exists g H : Z[X], Irreducible H /\ H \mid derivative g /\ H^2 \mid f.comp g
```

with a strengthened form carrying $2 \leq \deg g$ and $1 \leq \deg H$, and with Corollary 1 proved from it. Every result depends only on the three standard axioms `propext`, `Classical.choice` and `Quot.sound`.

References

- [Ded78] Richard Dedekind. “Über den Zusammenhang zwischen der Theorie der Ideale und der Theorie der höheren Kongruenzen”. In: *Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen* (1878).
- [The20] The mathlib Community. “The Lean Mathematical Library”. In: *Certified Programs and Proofs (CPP 2020)*. ACM, 2020.
- [MU21] Leonardo de Moura and Sebastian Ullrich. “The Lean 4 Theorem Prover and Programming Language”. In: *Automated Deduction – CADE 28*. Lecture Notes in Computer Science. Springer, 2021.