

21-269  
Vector Analysis

Rohan Jain

# Contents

Chapter 1	Page 2
1.1 The Real Numbers	2
1.2 First Recitation, 1/18	3
1.3 Natural Numbers	4
1.4 Density of Rationals	6
1.5 1/23 - Recitation - Proving Irrationality of $\sqrt{2}$	7
1.6 Exponents	9
1.7 1/25 - Recitation - Sequences of Set	10
1.8 Vector Spaces	12
1.9 Inner Products, Norms, and Metric Spaces	14
1.10 Open Sets	16
1.11 2/1 - Rectitation	18
1.12 Closure	19
1.13 Bolzano-Weierstrass	20
1.14 2/6 - Recitation - Spaces	21
1.15 Compactness	21
1.16 2/8 - Recitation	24
1.17 Limits	24
1.18 Limits Continued	26
1.19 Squeeze Theorem	27

# Chapter 1

## 1.1 The Real Numbers

### Definition 1.1.1: Partial Order

Let  $X$  be a set with a binary relation  $\leq$ .  $\leq$  is a *partial order* if:

1.  $x \leq x$  for all  $x \in X$  (reflexivity)
2.  $x \leq y$  and  $y \leq z$  implies  $x \leq z$  for all  $x, y, z \in X$  (transitivity)
3.  $x \leq y$  and  $y \leq x$  implies  $x = y$  for all  $x, y \in X$  (antisymmetry)

### Definition 1.1.2: Partially Ordered Set (poset)

A set  $X$  with a partial order  $\leq$  is called a *partially ordered set* or *poset*. It is notated as  $(X, \leq)$ .

### Definition 1.1.3: Total Order

A partial order  $\leq$  is a *total order* if for all  $x, y \in X$ , we have  $x \leq y$  or  $y \leq x$ .

### Example 1.1.1 (poset)

Let  $Y$  be a set. Define  $X = \{\text{all subsets of } Y\} = \mathcal{P}(Y)$ . Let  $E, F \in Y$ , we say that  $E \leq F$  if  $E \subseteq F$ . Then  $(X, \leq)$  is a poset. This is not a total order.

### Definition 1.1.4: Upper Bound, Bounded Above, Supremum, Maximum

Let  $(X, \leq)$  be a poset. Let  $E \subseteq X$ .

1.  $y \in X$  is an *upper bound* of  $E$  if  $x \leq y$  for all  $x \in E$ .
2.  $E$  is *bounded above* if it has at least one upper bound.
3. If  $E$  is nonempty and bounded above, then the *supremum*, if it exists, of  $E$ , denoted  $\sup E$ , is the least upper bound of  $E$ .
4.  $E$  has a *maximum* if there is  $y \in E$  such that  $x \leq y$  for all  $x \in E$ .

Properties worth mentioning:

1. If  $E$  has a maximum, then  $\sup E$  exists and is equal to the maximum.

*Proof.* Let  $y$  be the maximum of  $E$ . If  $z \in X$ , is an upper bound of  $E$ , then  $z \geq y$  because  $y \in E$ . Since  $z$  was arbitrary, this is true for any upper bound. Thus,  $y$  is the least upper bound of  $E$ .  $\odot$

### Example 1.1.2

Let  $Y$  be a nonempty set,  $(\mathcal{P}(Y), \subseteq)$  poset.  
 Fix nonempty  $Z \subseteq Y$ .

$$E = \{W \subseteq Y : W \subset Z\}$$

Trivially,  $Z$  is an upper bound of  $E$ . Realize that any superset of  $Z$  is an upper bound as well. We can postulate that the supremum of  $E$  is  $Z$ . We will now prove it:

*Proof.* Need to show that if  $F$  is an upper bound of  $E$ , then  $F \supseteq Z$ . If  $x \in Z$ , then  $\{x\} \in E$  by definition of  $E$ , so  $F \supseteq x$  for all  $x \in Z$ . Thus,  $F \supseteq Z$ .  $\odot$

Note that there is no maximum of  $E$ .

### Definition 1.1.5: Lower Bound, Bounded Below, Infimum, Minimum

Let  $(X, \leq)$  be a poset. Let  $E \subseteq X$ .

1.  $y \in X$  is a *lower bound* of  $E$  if  $y \leq x$  for all  $x \in E$ .
2.  $E$  is *bounded below* if it has at least one lower bound.
3. If  $E$  is nonempty and bounded below, then the *infimum*, if it exists, of  $E$ , denoted  $\inf E$ , is the greatest lower bound of  $E$ .
4.  $E$  has a *minimum* if there is  $y \in E$  such that  $y \leq x$  for all  $x \in E$ .

Going back to example 1.1.2, we can see that  $E$  is bounded below by  $\emptyset$ . The infimum of  $E$  is  $\emptyset$ . The minimum of  $E$  is also  $\emptyset$ .

### Definition 1.1.6: Complete

Let  $(X, \leq)$  poset.  $X$  is *complete* if every nonempty subset of  $X$  that is bounded above has a supremum.

### Example 1.1.3 ( $\mathbb{Q}$ )

$(\mathbb{Q}, \leq)$  is not complete.

### Claim 1.1.1 $\mathbb{R}$

There is a complete ordered field  $(\mathbb{R}, +, \cdot, \leq)$ . Its elements are called real numbers.

## 1.2 First Recitation, 1/18

### Exercise 1.2.1 Function Example

Let  $X$  be the set of all functions  $f : D_f \rightarrow Z$  with  $D_f \subseteq Y$ . We say that  $f \leq g$  if  $D_f \subseteq D_g$  and  $f(x) = g(x)$  for all  $x \in D_f$ . Is  $(X, \leq)$  a poset? Is it complete?

*Proof.* To show that  $(X, \leq)$  is complete, we need to show that every nonempty subset of  $X$  that is bounded above has a supremum. Let  $E \subseteq X$  be nonempty and bounded above. Let  $G = \bigcup_{f \in E} D_f$ .  $G$  is the union of all the domains of the functions in  $E$ .  $G$  is bounded above by the union of the upper bounds of the domains of the functions in  $E$ . Let  $H = \bigcup_{f \in E} f(D_f)$ .  $H$  is bounded above by the union of the upper bounds of the ranges of the functions in  $E$ . Let  $F : G \rightarrow H$  be defined as  $F(x) = f(x)$  for all  $x \in D_f$ .  $F$  is the supremum of  $E$ .  $\odot$

## 1.3 Natural Numbers

### Exercise 1.3.1

Take  $(X, +, \cdot, \leq)$  ordered field. Prove:

1. If  $0 \leq x$ , then  $-x \leq 0$ .
2. If  $x \leq y$ , and  $0 \leq z \neq 0$ , then  $xz \leq yz$ .
3. For all  $x \in X$ ,  $0 \leq x^2$ .
4. Prove  $0 < 1$ .

*Proof.* Fields have the following important properties:

- If  $a \leq b$ , then  $a + c \leq b + c$ .
  - If  $a, b \geq 0$ , then  $ab \geq 0$ .
1. Take the first property with  $a = 0$ ,  $b = x$ , and  $c = -x$ . Then  $0 \leq x \implies 0 + (-x) \leq x + (-x) \implies -x \leq 0$ .
  2. If  $x \leq y$ , then  $0 \leq y + (-x)$ . By the second property,  $0 \leq z \cdot (y + (-x)) = zy + (-zx)$ . Then  $0 \leq zy + (-zx) \implies zx \leq zy$ .
  3. We split into the three trichotomy cases:
    - If  $x = 0$ , then  $0 \leq 0^2$ .
    - If  $x < 0$  with  $x \neq 0$ , then  $0 \leq -x$ . By the second property,  $0 \leq (-x)^2 = (-x)(-x) = x^2$ .
    - If  $x > 0$ , then  $0 \leq x$ . By the second property,  $0 \leq x^2$ .
  4. FSOC, assume  $0 > 1$  and multiply both sides by 1. Then we get  $0 \cdot 1 > 1 \cdot 1 \implies 0 > (1)^2$ , which is a contradiction to the third property we proved.

☺

### Definition 1.3.1: Inductive

Take  $E \subseteq \mathbb{R}$ .  $E$  is *inductive* if  $1 \in E$  and  $x \in E$  implies  $x + 1 \in E$ .

### Example 1.3.1 (Inductive Sets)

- $\mathbb{R}$  is inductive.
- $\{x \in \mathbb{R} : 0 \leq x\}$

*Proof.*  $1 \in E$  because  $1 \geq 0$ . If  $x \in E$ , then  $x + 1 \geq 0$ , so  $x + 1 \in E$ .

☺

### Definition 1.3.2: Natural Numbers

The intersection of all inductive sets is denoted  $\mathbb{N}$ . The elements of  $\mathbb{N}$  are called *natural numbers*.

Properties of  $\mathbb{N}$ :

- $\mathbb{N} \neq \emptyset$ . Since  $1 \in$  every inductive set,  $1 \in \mathbb{N}$ .
- $\mathbb{N}$  is an inductive set.

**Theorem 1.3.1** Induction

For every  $n \in \mathbb{N}$ , let  $P(n)$  be a proposition such that:

1.  $P(1)$  is true.
2. If  $P(n)$ , then  $P(n + 1)$ .

Then  $P(n)$  is true for every  $n \in \mathbb{N}$

*Proof.*  $E = \{n \in \mathbb{N} : P(n)\}$  is inductive by 1. and 2. So,  $\mathbb{N} \subseteq E$ , but  $E \subseteq \mathbb{N}$  by definition of  $\mathbb{N}$ . Thus,  $E = \mathbb{N}$ . ☺

**Theorem 1.3.2** Archimedean Property

Let  $a, b \in \mathbb{R}$  with  $a > 0$ . Then there is  $n \in \mathbb{N}$  such that  $na > b$ .

*Proof.* If  $b \leq 0$ , then we take  $n = 1$ . Assume  $b > 0$ . For sake of contradiction, assume there does not exist  $n$  such that  $na > b$ . Then  $E = \{na : n \in \mathbb{N}\}$  is bounded above by  $b$ . Let  $c = \sup E$ .  $c - a \leq c$ , so  $c - a$  is not an upper bound of  $E$ . Thus, there is  $n \in \mathbb{N}$  such that  $c - a \leq na$ . Then  $c \leq (n + 1)a$ . But  $c$  is an upper bound of  $E$ , so  $c \geq (n + 1)a$ . Thus,  $c = (n + 1)a$ . But  $c \in E$ , so  $c = na$  for some  $n \in \mathbb{N}$ . Thus,  $na = (n + 1)a$ , so  $n = n + 1$ , which is a contradiction. ☹

**Definition 1.3.3: Integers**

$$\mathbb{Z} := \mathbb{N} \cup \{0\} \cup \{-n : n \in \mathbb{N}\}$$

**Theorem 1.3.3** Integer Part

For every  $x \in \mathbb{R}$ , there is a unique  $k \in \mathbb{Z}$  such that  $k \leq x < k + 1$ .

**Definition 1.3.4: Integer Part**

The  $k$  that satisfies the above theorem is called the *integer part* of  $x$ , denoted  $\lfloor x \rfloor$ .

*Proof.* Let  $E = \{k \in \mathbb{Z} : k \leq x\}$ . First we show that  $E$  is nonempty.

- If  $x \geq 0$ , then  $0 \in E$ , so  $E$  is nonempty.
- If  $x < 0$ , then  $-x > 0$ . By the Archimedean property, there is  $n \in \mathbb{N}$  such that  $n > -x$ . Thus,  $-n < x$ . So,  $-n \in E$ , so  $E$  is nonempty.

Now we show that  $E$  is bounded from above. Very clearly,  $x$  is an upper bound. By supremum property, there is  $L = \sup(E)$  and  $L \in \mathbb{R}$ .  $L - 1$  is not an upper bound, which means that there is an element  $k \in E$  such that  $L - 1 < k$ . But since  $L$  is the supremum,  $L \geq k$ . Thus,  $L - 1 < k \leq L$ . So,  $L < k + 1$  so  $k + 1 \notin E$ . Now,  $k \leq x$  since  $k \in E$ . Now we show that  $k$  is unique. Assume there is  $m \in \mathbb{Z}$  such that  $m \leq x < m + 1$ . Then  $m \in E$ , so  $m \leq L$ . But  $L$  is the supremum, so  $L \geq m$ . Thus,  $L = m$ . So,  $k = m$ . ☹

**Definition 1.3.5:  $\mathbb{Q}$** 

If  $p \in \mathbb{Z}$  with  $p \neq 0$ , then  $\exists p^{-1} \in \mathbb{R}$ . Define  $\mathbb{Q} = \{pq^{-1} : p, q \in \mathbb{Z}, p \neq 0\}$ .

## 1.4 Density of Rationals

### Theorem 1.4.1 Density of the Rationals

Let  $a, b \in \mathbb{R}$  with  $a < b$ . Then there is  $r \in \mathbb{Q}$  such that  $a < r < b$ .

*Proof.* We have  $a < b \implies 0 = a + (-a) < b - a \implies 0 < \frac{1}{b-a}$ . By the integer part theorem, there is  $q \in \mathbb{Z}$  such that  $\frac{1}{b-a} < q$ . So now,  $\frac{1}{q} < b - a \implies a < a + \frac{1}{q} < b$ . Multiply both sides by  $q > 0$  to get  $aq < a + 1 < bq$ . By the integer part theorem, there is  $p \in \mathbb{Z}$  such that  $p \leq qa < p + 1$  (i.e.  $p = \lfloor qa \rfloor$ ). Since  $qa < p + 1 \leq qa + 1 < qb$ . Getting rid of unnecessary stuff, we have  $qa < p + 1 < qb$ . Thus,  $a < \frac{p+1}{q} < b$ . Let  $r = \frac{p+1}{q}$ . Then  $r \in \mathbb{Q}$  and  $a < r < b$ .  $\odot$

### Definition 1.4.1: Irrational Numbers

$\mathbb{R} \setminus \mathbb{Q}$  is the set of *irrational numbers*.

### Exercise 1.4.1 TODO in Recitation 1/23

- Prove that there is no  $r \in \mathbb{Q}$  such that  $r^2 = 2$ .
- Prove that “ $\sqrt{2}$ ” exists in  $\mathbb{R}$ . (prove that there is at least one irrational number)
  - Have to play with the set  $E = \{x \in \mathbb{R} : x > 0, x^2 < 2\}$ .

### Theorem 1.4.2 Density of Irrationals

Let  $a, b \in \mathbb{R}$  with  $a < b$ . Then there is  $x \in \mathbb{R} \setminus \mathbb{Q}$  such that  $a < x < b$ .

*Proof.*  $a < b \implies a\sqrt{2} < b\sqrt{2}$ . By the density of rationals, there is  $r \in \mathbb{Q}$  such that  $a\sqrt{2} < r < b\sqrt{2}$ . Then  $a < \frac{r}{\sqrt{2}} < b$ . Let  $x = \frac{r}{\sqrt{2}}$ . If  $r = 0$ , then  $a\sqrt{2} < 0 < b\sqrt{2}$ . By previous theorem, we can find  $q \in \mathbb{Q}$  such that  $a\sqrt{2} < q < 0 < b\sqrt{2}$ . Then  $a < \frac{q}{\sqrt{2}} < b$ . Let  $x = \frac{q}{\sqrt{2}}$ . Then  $x \in \mathbb{R} \setminus \mathbb{Q}$  and  $a < x < b$ .  $\odot$

### Note:

Take  $x \in \mathbb{R}$ ,  $E = \{r \in \mathbb{Q} : r < x\}$ .  $x$  is the upper bound of  $E$ . This set is nonempty because we can take  $x - 1 < r < x$ . Now we prove that  $x = \sup E$ .

*Proof.* Assume  $\exists L$  upper bound of  $E$  such that  $L < x$ . Then  $L < x \implies$  there exists some  $r \in \mathbb{Q}$  such that  $L < r < x$ , but  $r \in E$ , so  $L$  is not an upper bound of  $E$ . Thus,  $L$  cannot be an upper bound of  $E$  and  $x$  is the least upper bound of  $E$ .  $\odot$

Since now we know that  $\sqrt{2} = \sup\{r \in \mathbb{Q} : r < \sqrt{2}\}$ , we can also define  $3^{\sqrt{2}} = \sup\{3^r : r \in \mathbb{Q}, r < \sqrt{2}\}$ .

### Definition 1.4.2: $x^0$

Let  $0 \neq x \in \mathbb{R}$ . We define  $x^0 = 1$ .

### Definition 1.4.3: $x^n$

Let  $x \in \mathbb{R}$ ,  $n \in \mathbb{N}$ . We start with  $x^1 := x$ . Then assume  $x^m$  has been defined. Then we say  $x^{m+1} := x^m \cdot x$ .

### Definition 1.4.4: $x^{p/m}$

Let  $x \in \mathbb{R}$ ,  $p \in \mathbb{Z}$ ,  $m \in \mathbb{N}$ . We say  $x^{p/m} = \sqrt[m]{x^p}$ .

### Exercise 1.4.2 Properties of Exponents

Let  $x \in \mathbb{R}$ ,  $r, q \in \mathbb{Q}$ , and  $x, r, q > 0$ . Prove the following:

- $x^r \cdot x^q = x^{r+q}$
- $(x^r)^q = (x^q)^r = x^{rq}$

*Proof.*



### Definition 1.4.5: Negative Exponent

Take  $x > 0$ ,  $r = -\frac{p}{m}$  for  $p, m \in \mathbb{N}$ . First, we have that  $x^{-r} := (x^{-1})^{p/m}$ .

### Exercise 1.4.3 More Properties of Exponents

Take  $x \in \mathbb{R}$ ,  $x > 0$ ,  $r, q \in \mathbb{Q}$ . Prove the following:

- If  $r > 0$ , prove that  $x^r > 1$ .
- If  $r < q$ , prove that  $x^r < x^q$ .

## 1.5 1/23 - Recitation - Proving Irrationality of $\sqrt{2}$

Existence of  $\sqrt{2}$ :

1. Let  $E = \{x \in \mathbb{R} : x > 0, x^2 < 2\}$ . Prove that  $E$  is non-empty and that  $E$  is bounded above.

*Proof.* We know that  $0 < 1$  and from that we get  $1^2 = 1 < 2$ , which can be checked by subtracting 1 from both sides. As such  $E$  is nonempty.

Now we show that  $E$  is bounded above. We know that  $2^2 = 4 > 2 > a^2 \in E$ , so  $2^2 > a^2 \Rightarrow 2 > a$ , so 2 is an upper bound of  $E$ . ☺

2. By the completeness of  $(\mathbb{R}, \leq)$ ,  $E$  has a supremum,  $L$ . Prove that  $L > 0$  and that  $L^2 = 2$ .

*Proof.* Since  $L$  is the least upper bound, it has to be greater than 1 which is in the set  $E$ . Therefore,  $L > 1 > 0 \Rightarrow L > 0$ .

Now we show that  $L^2 \geq 2$ . For sake of contradiction, assume  $L^2 < 2$ . Since  $L > 0$ , this means that  $L \in E$ . By the density of rationals, there exists  $r \in \mathbb{Q}$  such that  $L < r < \sqrt{2}$ . Since  $L$  is an upper bound of  $E$ ,  $r \notin E$ . But  $r \in \mathbb{Q}$ , so  $r^2 \neq 2$ . Thus,  $r^2 > 2$ . Since  $r > 0$ ,  $r^2 > 2 \Rightarrow r > \sqrt{2}$ . But  $r < \sqrt{2}$ , so we have a contradiction. Thus,  $L^2 \geq 2$ . ☺

3. Prove that if  $y \in \mathbb{R} \setminus E$  and  $y > 0$ , then  $y$  is an upper bound of  $E$ .

*Proof.* Assume  $y \in \mathbb{R} \setminus E$  and  $y > 0$ . We need to show that  $y$  is an upper bound of  $E$ . Assume for sake of contradiction that  $y$  is not an upper bound of  $E$ . Then there exists  $x \in E$  such that  $x > y$ . But  $x \in E \Rightarrow x^2 < 2$ . Since  $y > 0$ ,  $x^2 < 2 \Rightarrow y^2 < 2$ . But  $y \notin E$ , so  $y^2 \geq 2$ . But this would mean that  $y \in E$ . Contradiction. Thus,  $y$  is an upper bound of  $E$ . ☺

4. Prove that  $L^2 = 2$ .



*Proof.* We know that  $L^2 \geq 2$  from part 2. Now we show that  $L^2 \leq 2$ . Assume for sake of contradiction that  $L^2 > 2$ .

How small does  $\epsilon > 0$  need to be such that  $(L - \epsilon)^2 > 2$  as well.

Start with  $(L - \epsilon)^2 = L^2 - 2L\epsilon + \epsilon^2$ , which is greater than  $L^2 - 2L\epsilon$  since  $\epsilon > 0$ . So now, how small does  $\epsilon$  need to be such that  $L^2 > 2 \implies L^2 - 2L\epsilon > 2$  too.

$$\begin{aligned} 2L\epsilon &< 2 - L^2 \\ \epsilon &< \frac{2 - L^2}{2L} \end{aligned}$$

Since  $L^2 > 2$ , this means that an  $\epsilon$  can be found. This means that  $L$  is not the least upper bound. Contradiction. Thus,  $L^2 \leq 2$ .  $\odot$

## 1.6 Exponents

### Definition 1.6.1: $\sqrt{2}$

$$\sqrt{2} := \sup\{x \in \mathbb{R} : x > 0, x^2 < 2\}$$

### Exercise 1.6.1

For  $n \in \mathbb{N}, n \geq 2$ . Fix  $x > 0$ .

$$E = \{y \in \mathbb{R} : y > 0, y^n < x\}.$$

Prove that  $l = \sup E$  satisfies  $l^n = x$ .

*Proof.* We first need to show that  $\sup E$  exists. Let  $y = x/(1+x)$ . Then,  $0 \leq y < 1$ , so  $y^n \leq y < x$ . Thus,  $y \in E$ . So,  $E$  is nonempty.  $E$  is also bounded from above because  $x$  is an upper bound of  $E$ . Thus,  $\sup E$  exists by the completeness of  $\mathbb{R}$ . Let  $l = \sup E$ . We now show that  $l^n = x$ .

First we show that  $l^n \leq x$ . FSO, assume  $l^n > x$ . If you choose an  $\epsilon > 0$  that is small enough, then  $(l - \epsilon)^n > x$  as well. We can't do this because  $y > l - \epsilon$  for some  $y \in E$  since  $l$  is the supremum of  $E$ . As such, we arrive at a contradiction which means that  $l^n \leq x$ .

To show that  $l^n \geq x$ , assume FSO that  $l^n < x$ . Then we can choose an  $\epsilon$  such that  $(l + \epsilon)^n < x$ , meaning we have an element  $(l + \epsilon)$  which is in  $E$  but bigger than the supremum, which is a contradiction.

Thus,  $l^n = x$ . ⊕

### Definition 1.6.2: $\sqrt[n]{x}$

$$\sqrt[n]{x} := \sup\{y \in \mathbb{R} : y > 0, y^n < x\}$$

### Definition 1.6.3: $x^{p/q}$

$$x^{p/q} := \left(\sqrt[q]{x}\right)^p$$

### Definition 1.6.4: $x^q$

For  $q \in \mathbb{R}, q > 0$ , and  $x > 1$ .

$$x^q := \sup\{x^r : r \in \mathbb{Q}, 0 < r < q\}$$

### Example 1.6.1

$$\sqrt{2} = \sup\{r \in \mathbb{Q} : r > 0, r < \sqrt{2}\}$$

### Theorem 1.6.1

Take  $a, b \in \mathbb{R}, a, b > 0$  and  $x \in \mathbb{R} > 1$ . Then  $x^a \cdot x^b = x^{a+b}$ .

*Proof.* Let  $E_i = \{x^r : r \in \mathbb{Q}, r > 0, r < i\}$ . Consider  $E_a, E_b, E_{a+b}$ . Then let  $l_i = \sup(E_i)$ . Consider  $l_a, l_b, l_{a+b}$ . We want to show that  $l_a \cdot l_b = l_{a+b}$  by showing that both  $l_a \cdot l_b \leq l_{a+b}$  and  $l_a \cdot l_b \geq l_{a+b}$ .

Let  $r \in \mathbb{Q}$  with  $0 < r < a$ . Let  $s \in \mathbb{Q}$  with  $0 < s < b$ . Then we have that  $x^r \cdot x^s = x^{r+s}$  (from the exercise two days ago and since  $r, s \in \mathbb{Q}$ .) we know that  $0 < r + s < a + b$  and is rational. Thus,  $x^{r+s} \in E_{a+b}$ . Thus,  $x^r \cdot x^s \leq l_{a+b}$ .

We want to divide both sides by  $x^s$  while fixing  $r$ . So, we have that  $x^r \leq \frac{l_{a+b}}{x^s}$ , which is true for all  $r \in \mathbb{Q}$ , such that  $0 < r < a$ . Thus,  $\frac{l_{a+b}}{x^s}$  is an upper bound for  $E_a$ . Thus,  $l_a \leq \frac{l_{a+b}}{x^s}$ . Thus,  $x^s \leq \frac{l_{a+b}}{l_a}$ , meaning that  $\frac{l_{a+b}}{l_a}$  is an upper bound for  $E_b$ . Thus,  $l_b \leq \frac{l_{a+b}}{l_a}$ . Thus,  $l_a \cdot l_b \leq l_{a+b}$ .

Now we show that  $l_a \cdot l_b \geq l_{a+b}$ . Let  $t \in \mathbb{Q}$  with  $0 < t < a + b$ . We need  $0 < r \in \mathbb{Q} < a$  and  $0 < s \in \mathbb{Q} < b$  with  $t = r + s$ . We start by looking at  $t - a < b$ . By the density of  $\mathbb{Q}$ , find  $s \in \mathbb{Q}$  such that  $t - a < s < b$ . Take  $s > 0$  because  $b > 0$ . So  $t - s < a$ . By the density of  $\mathbb{Q}$ , find  $0 < p \in \mathbb{Q}$  such that  $t - s < p < a$ . So  $t < s + p$ . So,  $x^t < x^{s+p} = x^s x^p \leq l_a l_b$  since  $x^s \in E_b$  and  $x^p \in E_a$ . We know that  $l_a l_b$  is an upper bound of  $E_{a+b}$ , so  $l_{a+b} \leq l_a l_b$ .

Therefore  $l_a \cdot l_b = l_{a+b}$ .  $\odot$

#### Definition 1.6.5: Negative Exponents

Let  $x > 1$ ,  $a < 0$ . Then:

$$x^a := (x^{-a})^{-1}$$

#### Definition 1.6.6: Exponents between 0 and 1

Let  $x \in \mathbb{R}$  with  $0 < x < 1$  and  $a > 0$ . Then:

$$x^a := \left(\frac{1}{x}\right)^{-a}$$

An important note is that if we have  $E \subseteq (0, \infty)$  with a bounded  $E$ . Then if we define  $F = \{\frac{1}{x} : x \in E\}$ , then we have the following:

$$\begin{aligned} \sup E &= \frac{1}{\inf F} \\ \inf E &= \frac{1}{\sup F} \end{aligned}$$

## 1.7 1/25 - Recitation - Sequences of Set

#### Definition 1.7.1: Sequence of a Set

Given a set  $X$ , a sequence on  $X$  is a function  $f : \mathbb{N} \rightarrow X$ . We denote  $f(n)$  as  $x_n$ . We can also denote the sequence as  $\{x_n\}_{n=1}^{\infty}$ .

#### Definition 1.7.2

Let  $(X, \leq)$  be a poset and  $\{x_n\}_{n=1}^{\infty}$  be a sequence on  $X$ . Then  $E = \{x_n : n \in \mathbb{N}\}$  is a subset of  $X$ . We say that  $\{x_n\}_{n=1}^{\infty}$  is bounded from above if the set  $E$  is bounded from above. We say that  $\{x_n\}_{n=1}^{\infty}$  is bounded from below if the set  $E$  is bounded from below. We say that  $\{x_n\}_{n=1}^{\infty}$  is bounded if it is bounded from above and below.

#### Definition 1.7.3: Limit Superior

Let  $(X, \leq)$  be a poset. Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence on  $X$ . Suppose  $\{x_n\}_n$  is bounded from above. Then, we define the *limit superior* of  $x_n$  as  $n \rightarrow \infty$  as:

$$\limsup_{n \rightarrow \infty} x_n = \inf_{n \in \mathbb{N}} \sup_{k \geq n} x_k$$

**Definition 1.7.4: Limit Inferior**

Let  $(X, \leq)$  be a poset. Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence on  $X$ . Suppose  $\{x_n\}_n$  is bounded from below. Then, we define the *limit inferior* of  $x_n$  as  $n \rightarrow \infty$  as:

$$\liminf_{n \rightarrow \infty} x_n = \sup_{n \in \mathbb{N}} \inf_{k \geq n} x_k$$

### Exercise 1.7.1

1. Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence on  $\mathbb{R}$  bounded above. Prove that  $L \in \mathbb{R}$  is the limsup of  $\{x_n\}_{n=1}^{\infty}$  iff for every  $\epsilon > 0$ , there exists  $n_{\epsilon} \in \mathbb{N}$  such that:
  - (a)  $x_n < L + \epsilon$  for all  $n \geq n_{\epsilon}$ .
  - (b)  $L - \epsilon < x_n$  for infinitely many  $n$ .

*Proof.* Let  $L \in \mathbb{R}$  be the limsup of  $\{x_n\}_{n=1}^{\infty}$ . Let  $\epsilon > 0$ .  $L$  being the lim sup means that  $L = \inf_{n \in \mathbb{N}} \sup_{k \geq n} x_k$ . Thus,  $L \leq \sup_{k \geq n} x_k$  for all  $n \in \mathbb{N}$ . Thus,  $L - \epsilon < \sup_{k \geq n} x_k$  for all  $n \in \mathbb{N}$ . Then  $L - \epsilon$  is not an upper bound of  $\{x_n\}_{n=1}^{\infty}$ . Thus, there is  $n_{\epsilon} \in \mathbb{N}$  such that  $L - \epsilon < x_{n_{\epsilon}}$ . Thus,  $L - \epsilon < x_n$  for infinitely many  $n$ . Now we show that  $x_n < L + \epsilon$  for all  $n \geq n_{\epsilon}$ . Assume for sake of contradiction that there is  $n \geq n_{\epsilon}$  such that  $x_n \geq L + \epsilon$ . Then  $L + \epsilon$  is an upper bound of  $\{x_n\}_{n=1}^{\infty}$ . But  $L$  is the limsup, so  $L \geq L + \epsilon$ . Contradiction. Thus,  $x_n < L + \epsilon$  for all  $n \geq n_{\epsilon}$ .

Now we show the other direction. Assume that for every  $\epsilon > 0$ , there exists  $n_{\epsilon} \in \mathbb{N}$  such that  $x_n < L + \epsilon$  for all  $n \geq n_{\epsilon}$  and  $L - \epsilon < x_n$  for infinitely many  $n$ . We want to show that  $L$  is the limsup of  $\{x_n\}_{n=1}^{\infty}$ . We know that  $L$  is an upper bound of  $\{x_n\}_{n=1}^{\infty}$ . We need to show that  $L$  is the least upper bound. Assume for sake of contradiction that  $L$  is not the least upper bound. Then there is  $L' < L$  such that  $L'$  is an upper bound of  $\{x_n\}_{n=1}^{\infty}$ . Let  $\epsilon = L - L'$ . Then  $L' < L - \epsilon$ . But  $L - \epsilon < x_n$  for infinitely many  $n$ . But  $L' < L - \epsilon$ , so  $L'$  is not an upper bound of  $\{x_n\}_{n=1}^{\infty}$ . Contradiction.  $\odot$

## 1.8 Vector Spaces

### Example 1.8.1 (Vector Spaces)

- Euclidean Space  $\subseteq \mathbb{R}^n$ .  $x \in \mathbb{R}^n$  is a vector.  $x = (x_1, \dots, x_n)$ .
- Polynomial Space from  $\mathbb{R} \rightarrow \mathbb{R}$ .  $x \in \mathbb{R}^x$ .  $x = a_0 + a_1x + \dots + a_nx^n$ .
- $f : [a, b] \rightarrow \mathbb{R}$  continuous functions.

### Definition 1.8.1: Boundedness of Functions

Let  $E$  be a set and  $f : E \rightarrow \mathbb{R}$ .

1.  $f$  is bounded from above if the set  $f(E) = \{y \in \mathbb{R} : y = f(x), x \in E\}$  is bounded from above.
2.  $f$  is bounded from below if the set  $f(E) = \{y \in \mathbb{R} : y = f(x), x \in E\}$  is bounded from below.
3.  $f$  is bounded if  $f(E)$  is bounded.

### Definition 1.8.2: Inner Product

A function  $(\cdot, \cdot) : V \times V \rightarrow \mathbb{R}$  is an *inner product* if it satisfies the following properties:

- $(x, x) \geq 0$  for all  $x \in X$ .
- $(x, x) = 0$  iff  $x = 0$ .
- $(x, y) = (y, x)$  for all  $x, y \in X$ .
- $(sx + ty, z) = s(x, z) + t(y, z)$  for all  $x, y, z \in X$  and  $s, t \in \mathbb{R}$ .

### Example 1.8.2 (Examples of Inner Products)

- $\mathbb{R}^n$  with dot products.
- $f : [a, b] \rightarrow \mathbb{R}$  with  $(f, g) = \int_a^b f(x)g(x)dx$ . This is not an inner product because we can define:

$$f = \begin{cases} 1 & x = 0.5 \\ 0 & \text{otherwise} \end{cases}$$

which has an integral of 0. But  $f \neq 0$ . If we add that  $f$  is continuous, then it is an inner product.

### Definition 1.8.3: Norm

Let  $V$  be a vector space with an inner product  $(\cdot, \cdot)$ . Then the *norm* of  $x \in X$  is defined as  $\|\cdot\| : X \rightarrow [0, \infty)$  such that:

1.  $\|x\| = 0 \iff x = 0$
2.  $\|tx\| = |t|\|x\|$  for all  $x \in X$
3.  $\|x + y\| \leq \|x\| + \|y\|$  for all  $x, y \in X$

### Example 1.8.3 (Examples of Norms)

- $\|x\| = \sqrt{(x, x)}$  for  $x \in \mathbb{R}^n$
- $X = \{f : E \rightarrow \mathbb{R}, f \text{ bounded}\}$ .  $\|f\| = \sup_{x \in E} |f(x)|$ .
  - First property is obviously true.
  - For the second property, we use the fact that

$$\sup(tF) = \begin{cases} t \sup(F) & \text{if } t \geq 0 \\ t \inf(F) & \text{if } t < 0 \end{cases}$$

- For the third property, we use the triangle inequality:

$$\begin{aligned} \sup |f + g| &\leq \sup |f| + \sup |g| \\ |f(x) + g(x)| &\leq |f(x)| + |g(x)| \leq \sup |f| + \sup |g| \end{aligned}$$

#### Note:

Space of bounded functions denoted as  $\ell^\infty(E) = \{f : E \rightarrow \mathbb{R} : f \text{ bounded}\}$ .

### Theorem 1.8.1 Cauchy Schwarz Inequality

Let  $X$  be a vector space with an inner product  $(\cdot, \cdot)$ . Then for all  $x, y \in X$ , we have that  $|(x, y)| \leq \sqrt{(x, x)} \cdot \sqrt{(y, y)}$ .

*Proof.* Let  $y \neq 0$ . Consider  $(x + ty, x + ty) = (x, x + ty) + t(y, x + ty) = (x, x) + t(x, y) + t(y, x) + t^2(y, y)$ . We can

combine the middle terms to get  $t^2(y, y) + 2(x, y) + (x, x)$ , which is quadratic in  $t$ . Take  $t = -\frac{(x, y)}{(y, y)}$ .

$$\begin{aligned} 0 &\leq (x, x) - 2\frac{(x, x)^2}{(y, y)} + \frac{(x, y)^2}{(y, y)} \\ 0 &\leq (x, x)(y, y) - 2(x, y)^2 + (x, y)^2 \\ 0 &\leq (x, x)(y, y) - (x, y)^2 \\ (x, y)^2 &\leq (x, x)(y, y) \\ |(x, y)| &\leq \sqrt{(x, x)} \cdot \sqrt{(y, y)} \end{aligned}$$

⊕

## 1.9 Inner Products, Norms, and Metric Spaces

### Theorem 1.9.1

Let  $X$  be a vector space with an inner product  $(\cdot, \cdot)$ . Then  $\|x\| := \sqrt{(x, x)}$  is a norm.

*Proof.* We check the properties of norms:

1.  $\|x\| = 0 \iff \sqrt{(x, x)} = 0 \iff (x, x) = 0 \iff x = 0$ .
2.  $\|tx\| = \sqrt{(tx, tx)} = \sqrt{t^2(x, x)} = |t|\sqrt{(x, x)} = |t|\|x\|$ .
3.  $\|x + y\|^2 = (x + y, x + y) = (x, x) + 2(x, y) + (y, y) = \|x\|^2 + 2(x, y) + \|y\|^2 \leq \|x\|^2 + 2|(x, y)| + \|y\|^2 \leq \|x\|^2 + 2\|x\| \cdot \|y\| + \|y\|^2 = (\|x\| + \|y\|)^2$ .

⊕

### Corollary 1.9.1 Parallelogram Identity

Let  $X$  be a vector space with inner product  $(\cdot, \cdot)$ . Then for all  $x, y \in X$ , we have that

$$\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2$$

*Proof.*

$$\begin{aligned} \|x + y\|^2 + \|x - y\|^2 &= (x + y, x + y) + (x - y, x - y) \\ &= (x, x) + 2(x, y) + (y, y) + (x, x) - 2(x, y) + (y, y) \\ &= 2(x, x) + 2(y, y) \\ &= 2\|x\|^2 + 2\|y\|^2 \end{aligned}$$

⊕

If we subtract them instead, we get

$$\frac{\|x + y\|^2 - \|x - y\|^2}{4} = (x, y) \quad (*)$$

So, if  $\|\cdot\|$  is a norm, then if i want to define an inner product, I can use  $*$ .

### Exercise 1.9.1

Let  $\|\cdot\|$  be a norm. Then  $(x, y) := \frac{1}{4}(\|x + y\|^2 - \|x - y\|^2)$  is an inner product iff the parallelogram identity holds.

Linearity of inner products is the hard part to prove because we have to consider:

- $t \in \mathbb{N}$
- $t = \frac{1}{2}$
- $t \in \mathbb{Q}$
- $t \in \mathbb{R}$  (density of  $\mathbb{Q}$ )

**Note:**

For recitation:

1.  $X = \{f : E \rightarrow \mathbb{R} \text{ bounded}\}$ ,  $\|f\| = \sup_E |f|$ , does not satisfy the parallelogram identity.
2.  $x \in \mathbb{R}^N$ ,  $\|x\|_1 = |x_1| + |x_2| + \cdots + |x_N|$  does not satisfy the parallelogram identity.

**Definition 1.9.1: Metric**

Let  $X$  be a set. A *metric* on  $X$  is a function  $d : X \times X \rightarrow [0, \infty)$  such that:

1.  $d(x, y) = 0 \iff x = y$
2.  $d(x, y) = d(y, x)$  for all  $x, y \in X$
3.  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in X$

**Definition 1.9.2: Metric Space**

A set  $X$  with a metric  $d$  is called a *metric space* and is denoted as  $(X, d)$ .

**Example 1.9.1 (Metrics)**

Let  $X$  be a set. Then the following is a metric on  $X$ :

$$d(x, y) = \begin{cases} 0 & x = y \\ 1 & x \neq y \end{cases}$$

**Theorem 1.9.2** If  $X$  is a vector space with  $\|\cdot\|$  as a norm. Then

$$d(x, y) := \|x - y\|$$

is a metric on  $X$ .

*Proof.* We check all the properties of metrics.

- $d(x, y) = 0 = \|x - y\| \Rightarrow 0 = x - y \iff x = y$ .
- $d(x, y) = \|x - y\| = \|y - x\| = d(y, x)$ .
- $d(x, y) = \|x - y\| = \|x - z + z - y\| \leq \|x - z\| + \|z - y\| = d(x, z) + d(z, y)$ .





### Example 1.9.2

Let's define

$$d(x, y) = \left| \frac{x}{1 + |x|} - \frac{y}{1 + |y|} \right|$$

as a metric on  $\mathbb{R}$ . However, this is not a norm because  $d(tx, ty) \neq td(x, y)$ .

### Definition 1.9.3: Ball

Let  $(X, d)$  be a metric space. Let  $x \in X$  and  $r > 0$ . Then the *ball* of radius  $r$  centered at  $x$  is defined as  $B_r(x) = \{y \in X : d(x, y) < r\}$ .

### Example 1.9.3

- Take  $X = \mathbb{R}^2$  with  $(x, y) \in \mathbb{R}$ . Then define  $\|(x, y)\|_\infty = \max(|x|, |y|)$  is a norm. Take  $B((0, 0), 1) = \{(x, y) \in \mathbb{R}^2 : \|(x, y) - (0, 0)\|_\infty < 1\}$ . This is a square with vertices  $(1, 1), (-1, 1), (-1, -1), (1, -1)$ .
- If we have  $\|(x, y)\|_1 = |x| + |y|$ , then  $B((0, 0), 1) = \{(x, y) \in \mathbb{R}^2 : \|(x, y) - (0, 0)\|_1 < 1\}$ . This is a square with vertices  $(1, 0), (0, 1), (-1, 0), (0, -1)$ .

### Definition 1.9.4: Interior

Let  $(X, d)$  be a metric space and  $E \subseteq X$ .  $x \in E$  is called an *interior point* of  $E$  if there is  $B(x, r) \subseteq E$ . The set of all interior points of  $E$  is called the *interior* of  $E$  and is denoted as  $E^\circ$ .

### Definition 1.9.5: Open Set

$E$  is *open* if  $E = E^\circ$ .

## 1.10 Open Sets

### Example 1.10.1 (Balls)

$B(x, r)$  is open.

*Proof.* Let  $y \in B(x, r)$  and take  $B(y, r - d(x, y))$ . Let  $z \in B(y, r - d(x, y))$ . Then  $d(x, z) \leq d(x, y) + d(y, z) < d(x, y) + r - d(x, y) = r$ . Thus,  $z \in B(x, r)$ . Thus,  $B(y, r - d(x, y)) \subseteq B(x, r)$ . Thus,  $B(x, r)$  is open.  $\odot$

### Example 1.10.2 ( $\mathbb{R}$ )

1.  $E = (0, 1) \cap \mathbb{Q}$  is not open. Because the irrationals are dense, we can always find a rational number in any ball. Thus,  $E^\circ = \emptyset$ .
2.  $E = (3, 4)$  is open. Let  $x \in E$ . Take  $B(x, \min(x - 3, 4 - x))$ . Then  $B(x, \min(x - 3, 4 - x)) \subseteq E$ . Thus,  $E$  is open.
3.  $E = [3, 4)$  is not open.  $E^\circ = (3, 4)$ .
4.  $E = \{x \in \mathbb{R} : x^3 - 3x + 4 > 0\}$ . This is open and we'll be able to use continuity to prove this easily later.
5.  $l^\infty([0, 1]) = \{f : [0, 1] \rightarrow \mathbb{R} \text{ bounded}\}$ .  $\|f\|_\infty = \sup_{[0, 1]} |f|$ .  $d(f, g) = \|f - g\|_\infty$ .  $E = \{f \in l^\infty([0, 1]) : f(x) > 0 \forall x \in [0, 1]\}$  is open? (finish in recitation)

Properties of open sets  $(X, d)$ :

- $\emptyset$  is open.  $X$  is open.
- Infinite intersections of open sets are not necessarily open. For example, we have  $\bigcap_{n=1}^{\infty} (-1/n, 1/n) = \{0\}$ , which is not open.
- Finite intersections of open sets are open. Consider  $U_1, \dots, U_n$ . Let  $x \in \bigcap_{i=1}^n U_i$ . Then  $x \in U_i$  for all  $i$ . Since  $U_i$  is open, there exists  $r_i > 0$  such that  $B(x, r_i) \subseteq U_i$ . Let  $r = \min(r_1, \dots, r_n)$ . Then  $B(x, r) \subseteq U_i$  for all  $i$ . Thus,  $B(x, r) \subseteq \bigcap_{i=1}^n U_i$ .
- Unions of open sets are open because if a point in the union is contained in one of the open sets, then there is a ball in that set that is contained in the union.

#### Definition 1.10.1: Topological Space

Let  $X$  be a set. A *topology* on  $X$  is a collection  $\mathcal{T}$  of subsets of  $X$  such that:

1.  $\emptyset, X \in \mathcal{T}$ .
2. If  $U_1, \dots, U_n \in \mathcal{T}$ , then  $\bigcap_{i=1}^n U_i \in \mathcal{T}$ . (finite intersections)
3. If  $U_\alpha \in \mathcal{T}$  for all  $\alpha \in A$ , then  $\bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}$ . (arbitrary unions)

Elements of  $\mathcal{T}$  are called open sets.

#### Definition 1.10.2: Closed

Let  $(X, d)$  be a metric space. We say  $C \subseteq X$  is *closed* if  $X \setminus C$  is open.

Note that  $X$  and  $\emptyset$  are both open and closed.

#### Example 1.10.3 (Open and Closed Sets)

- $[0, 1)$  is not open or closed.
- $[0, 1]$  is closed.

Properties of closed sets:

- $\emptyset$  and  $X$  are closed.
- Infinite intersections of closed sets are closed. (De Morgan's Law)
- Finite unions of closed sets are closed. For example, if we have  $\bigcup_{m=1}^{\infty} (-\infty, -\frac{1}{m}) = (-\infty, 0)$  which is closed.

## 1.11 2/1 - Rectitation

Recall:

1. Let  $\{x_n\}$  be a sequence bounded above in  $\mathbb{R}$ . Then  $L \in \mathbb{R}$  is the limit superior of  $\{x_n\}$  if for every  $\epsilon > 0$ , there exists  $n_\epsilon \in \mathbb{N}$  such that:
  - (a)  $x_n < L + \epsilon$  for all  $n \geq n_\epsilon$ .
  - (b)  $x_n > L - \epsilon$  for infinitely many  $n$ .
2. Let  $\{x_n\}$  be a sequence bounded below in  $\mathbb{R}$ . Then  $L \in \mathbb{R}$  is the limit inferior of  $\{x_n\}$  if for every  $\epsilon > 0$ , there exists  $n_\epsilon \in \mathbb{N}$  such that:
  - (a)  $x_n < L + \epsilon$  for infinitely many  $n$ .
  - (b)  $x_n > L - \epsilon$  for all  $n \geq n_\epsilon$ .

Now consider the following sequence:

$$x_n = (-1)^n \frac{2n}{n+1} \in \mathbb{R}$$

Prove that  $\limsup_{n \rightarrow \infty} x_n = 2$ .

*Proof.* We need to show that for every  $\epsilon > 0$ , there exists  $n_\epsilon \in \mathbb{N}$  such that:

1.  $x_n < 2 + \epsilon$  for all  $n \geq n_\epsilon$ .
2.  $2 - \epsilon < x_n$  for infinitely many  $n$ .

Let  $\epsilon > 0$ . We need to find  $n_\epsilon \in \mathbb{N}$  such that  $x_n < 2 + \epsilon$  for all  $n \geq n_\epsilon$  and  $2 - \epsilon < x_n$  for infinitely many  $n$ . We can find  $n_\epsilon \in \mathbb{N}$  such that  $2 - \epsilon < x_n$  for all  $n \geq n_\epsilon$ . Then  $x_n < 2 + \epsilon$  for all  $n \geq n_\epsilon$ . Thus,  $\limsup_{n \rightarrow \infty} x_n = 2$ . ☺

Now prove that for any  $\{x_n\}$  in  $\mathbb{R}$ , prove that  $\liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n$ .

*Proof.* Comes quickly from properties of limits and that the inf is less than the sup. ☺

Now prove that  $\liminf_{n \rightarrow \infty} -x_n = -\limsup_{n \rightarrow \infty} x_n$  and that  $\limsup_{n \rightarrow \infty} -x_n = -\liminf_{n \rightarrow \infty} x_n$ .

*Proof.* We start by using the property that  $\inf(-E) = -\sup(E)$ . Then we use the property that  $\sup(-E) = -\inf(E)$ .  
So,

$$\begin{aligned} \liminf_{n \rightarrow \infty} -x_n &= \sup_{n \in \mathbb{N}} \inf_{k \geq n} -x_k \\ &= \sup_{n \in \mathbb{N}} -\sup_{k \geq n} x_k \\ &= -\inf_{n \in \mathbb{N}} \sup_{k \geq n} x_k \\ &= -\limsup_{n \rightarrow \infty} x_n \end{aligned}$$

☺

## 1.12 Closure

### Definition 1.12.1: Closure

Let  $(X, d)$  be a metric space with  $A \subset X$ . Then the *closure* of  $A$  is defined as  $\bar{A}$ , the intersection of all sets that contain  $A$ .

### Definition 1.12.2: Boundary Point

Let  $(X, d)$  be a metric space with  $E \subseteq X$ . Then  $x \in X$  is a *boundary point* of  $E$  if for every  $r > 0$ ,  $B(x, r) \cap E \neq \emptyset$  and  $B(x, r) \cap (X \setminus E) \neq \emptyset$ . The set of all boundary points is denoted as  $\partial E$ .

### Theorem 1.12.1

Let  $(X, d)$  be a metric space and  $E \subseteq X$ . Then  $\bar{E} = E \cup \partial E$ .

*Proof.* Let  $x \in \bar{E}$ . F.S.O.C., assume  $x \notin E \cup \partial E$ . Since  $x \notin \partial E$ , there exists  $r > 0$  such that  $B(x, r)$  that doesn't intersect with either  $E$  or complement of  $E$ . But since  $x \notin E$ , only the second option can occur. So there exists  $r$  such that  $B(x, r) \cap E = \emptyset$ . Because of that and the fact that  $B(x, r)$  is open, it follows that  $X \setminus B(x, r)$  is closed and contains  $E$ . By the definition of  $\bar{E}$ , we have that  $\bar{E} \subseteq X \setminus B(x, r)$ . But this is a contradiction because  $x \in \bar{E}$ .

Conversely, let  $x \in E \cup \partial E$  and assume  $x \notin \bar{E}$ . Since  $\bar{E}$  is closed,  $X \setminus \bar{E}$  is open. Using the fact that  $x \in E \cup \partial E$ , we have that we can find a  $B(x, r) \subseteq X \setminus \bar{E}$ . But this is a contradiction because  $B(x, r)$  is open and contains  $E$ . Thus,  $E \cup \partial E \subseteq \bar{E}$ .  $\odot$

### Definition 1.12.3: Accumulation Point

Let  $(X, d)$  be a metric space with  $E \subseteq X$ . Then  $x \in X$  is an *accumulation point* of  $E$  if for every  $r > 0$ , there exists  $y \in E$  such that  $y \neq x$  and  $d(x, y) < r$ .

### Definition 1.12.4: Interval

$I \subseteq \mathbb{R}$  is an *interval* if we have that  $z \in I$  for all  $x < z < y$ .

### Definition 1.12.5: Rectangle

$R \subseteq \mathbb{R}^N$  is a *rectangle* if  $R = I_1 \times \cdots \times I_N$  where  $I_1, \dots, I_N$  are intervals in  $\mathbb{R}$ .

### Definition 1.12.6: Sequence

Let  $X$  be a set. A *sequence* is a function  $f : \mathbb{N} \rightarrow X$ . We denote  $f(n)$  as  $x_n$ .

### Definition 1.12.7: Convergent Sequence

Let  $(X, d)$  be a metric space. A sequence  $\{x_n\}_{n=1}^{\infty}$  is *convergent* if there exists  $x \in X$  such that for every  $\epsilon > 0$ , there exists  $n_{\epsilon} \in \mathbb{N}$  such that  $d(x, x_n) < \epsilon$  for all  $n \geq n_{\epsilon}$ . We write  $x_n \rightarrow x$  as  $n \rightarrow \infty$  or  $\lim_{n \rightarrow \infty} x_n = x$ .

## 1.13 Bolzano-Weierstrass

### Theorem 1.13.1 Bolzano-Weierstrauss

If  $E \subset \mathbb{R}^N$  is bounded and contains infinitely many distinct points, then  $E$  has an accumulation point

*Proof.*

#### Lemma 1.13.1 1

If  $[a_n, b_n] \supseteq [a_{n+1}, b_{n+1}]$  for all  $n$ , then  $\bigcap_{n=1}^{\infty} [a_n, b_n] \neq \emptyset$ .

*Proof.* For all  $a_n$  and  $b_n$ , we have:

$$\begin{aligned} a_1 &\leq a_2 \leq \cdots \\ b_1 &\geq b_2 \geq \cdots \end{aligned}$$

Let

$$A := \{a_1, a_2, \dots\}.$$

We have that  $a_n \leq b_n \leq b_1$  for all  $n$ . So  $A$  is bounded above, so by the supremum property, there exists  $x = \sup A \in \mathbb{R}$  and  $a_n \leq x$  for all  $n \in \mathbb{N}$ . We claim that  $x \leq b_n$  as well. If not, then there exists  $m \in \mathbb{N}$  such that  $b_m < x$ . Since  $x$  is an upper bound of  $A$ , we'll have that there's an  $n \in \mathbb{N}$  such that  $b_m < a_n \leq x$ . Find  $k \geq m, n$ , then we have  $b_m < a_n \leq a_k \leq b_k \leq b_m$ , which is a contradiction. This proves the claim. Hence,  $x \in [a_n, b_n]$  for all  $n$ . Thus,  $x \in \bigcap_{n=1}^{\infty} [a_n, b_n]$ .  $\odot$

#### Lemma 1.13.2 2

Let  $R_n$  be a closed and bounded rectangle. Assume that  $R_1 \supseteq R_2 \supseteq \cdots$ . Then  $\bigcap_{n=1}^{\infty} R_n \neq \emptyset$ .

*Proof.* We know that

$$\begin{aligned} R_n &= [a_{1,n}, b_{1,n}] \times \cdots \times [a_{N,n}, b_{N,n}] \\ R_{n+1} &= [a_{1,n+1}, b_{1,n+1}] \times \cdots \times [a_{N,n+1}, b_{N,n+1}] \end{aligned}$$

We can apply lemma 1  $N$  times (for each of the components of  $R_n$ ) to find that  $x_1, x_2, \dots, x_N \in \mathbb{R}$  such that  $a_{i,n} \leq x_i \leq b_{i,n}$  for all  $1 \leq i \leq N$ . Then, if you take  $x = (x_1, \dots, x_N)$ , then  $x \in R_n$  for all  $n$ . Thus,  $x \in \bigcap_{n=1}^{\infty} R_n$ .  $\odot$

#### Lemma 1.13.3 3

Let  $(X, d)$  be a metric space with  $E \subseteq X$ . Then  $x \in X$  is an accumulation point of  $E$  if and only if there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  in  $E$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$ .

*Proof.* Let  $x \in X$  be an accumulation point of  $E$ . Take  $r = \frac{1}{n}$ . Find  $x_n \in B\left(x, \frac{1}{n}\right) \cap E$  with  $x_n \neq x$ . We claim  $x_n \rightarrow x$ . Given  $\epsilon > 0$ , find  $n_\epsilon \geq \frac{1}{\epsilon}$ . Then  $d(x, x_n) < \frac{1}{n} \leq \frac{1}{n_\epsilon}$  for all  $n \geq n_\epsilon$ . Thus,  $x_n \rightarrow x$  as  $n \rightarrow \infty$ .

Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence in  $E$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$ . We claim that  $x \in \text{acc}(E)$ . Let  $r > 0$  and take  $\epsilon = r$ . Then there exists  $n_\epsilon \in \mathbb{N}$  such that  $d(x, x_n) < \epsilon = r$  for all  $n \geq n_\epsilon$ . Thus,  $x_n \in B(x, r) \cap E$  for all  $n \geq n_\epsilon$ . Thus,  $x \in \text{acc}(E)$ .  $\odot$

Now we prove the actual theorem. Let  $E \subseteq \mathbb{R}^N$  be bounded.  $E \subseteq B(0, r)$  for some  $r$ . Let  $Q_1$  be the closed cube centered at 0 with sidelength  $2r$ . Pick some point  $x_1 \in E \subseteq Q_1$ . Subdivide  $Q_1$  into  $2^N$  closed cubes of sidelength  $\frac{2r}{2}$ . Let  $Q_2$  be the closed cube containing  $x_1$ . Pick some point  $x_2 \in E \cap Q_2$  with  $x_2 \neq x_1$ . Inductively, assume  $Q_1 \supseteq Q_2 \supseteq \cdots \supseteq Q_n$  have been chosen. Then  $Q_n$  is a closed cube of sidelength  $\frac{2r}{2^{n-1}}$  containing  $x_n$ . Each

$Q_n$  contains infinitely many elements of  $E$ . Assume also that  $x_1, x_2, \dots, x_n \in E$  have been chosen with  $x_i \in Q_i$  and  $x_i \neq x_j$  for  $i \neq j$ .

Now we can subdivide  $Q_n$  to get  $Q_{n+1}$  and continue this process infinitely.

By Lemma 2, we know that  $\bigcap_{n=1}^{\infty} Q_n \neq \emptyset$ . Let  $x \in \bigcap_{n=1}^{\infty} Q_n$ . Now we need to show there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  in  $E$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$  but  $x_i \neq x$  for any  $i$  because then the rest of the points won't converge to  $x$ . If  $x = x_i$  for some  $i$ , we can just pick another point.

So WLOG, assume  $x_n \neq x$  for any  $n$ . So we claim  $x_n \rightarrow x$  as  $n \rightarrow \infty$ . We know that in  $Q_n$ , the difference between any two points in this cube is given by:

$$\|x_n - x\| = \sqrt{(x_{n,1} - x_1)^2 + (x_{n,2} - x_2)^2 + \dots + (x_{n,N} - x_N)^2} \leq \sqrt{\frac{2r}{2^{n-1}} + \frac{2r}{2^{n-1}} + \dots + \frac{2r}{2^{n-1}}} = \sqrt{N} \frac{2r}{2^{n-1}}$$

This value is less than  $\epsilon$  for all large  $n$ , so this concludes the proof.  $\ominus$

## 1.14 2/6 - Recitation - Spaces

Let  $X = \{f : [0, 1] \rightarrow \mathbb{R} \text{ bounded}\}$ . Define  $\|f\| = \sup_{x \in [0, 1]} |f(x)|$ . Prove that  $(X, \|\cdot\|)$  does not suffice parallelogram identity. That is, show a counterexample to the parallelogram identity, which is

$$\|f + g\|^2 + \|f - g\|^2 = 2\|f\|^2 + 2\|g\|^2$$

*Proof.* Counterexample: Let  $f(x) = x$  and  $g(x) = 1$ .  $\ominus$

Now given a normed space which satisfies the parallelogram identity, can we define an inner product?

*Proof.* Yes. We can define  $(f, g) = \frac{1}{4}(\|f + g\|^2 - \|f - g\|^2)$ . We can prove that this is an inner product.

Linearity of products because the other properties are easy to prove. We need to show that  $(x + y, z) = (x, z) + (y, z)$ . I'm so lazy so I won't tth.

We now show that  $(tx, y) = t(x, y) \forall t \in \mathbb{Z}$ . We proceed with induction for  $t \in \mathbb{Z}^+$

Our two base cases are  $t = 0, 1$ . For  $t = 0$ , we have that  $(0x, y) = (0, y) = 0 = 0(0, y)$ . For  $t = 1$ , we have that  $(x, y) = (x, y) = 1(x, y)$ .

Now we assume that  $(tx, y) = t(x, y)$  for some  $t \in \mathbb{Z}^+$ . Then we have that  $(t + 1)x = tx + x$ . Then we have that  $(t + 1)x, y = (tx + x, y) = (tx, y) + (x, y) = t(x, y) + (x, y) = (t + 1)(x, y)$ . Thus, we have that  $(tx, y) = t(x, y)$  for all  $t \in \mathbb{Z}^+$ .

Now we have to deal with  $t \in \mathbb{Z}^-$ . We have that  $(tx, y) = -t(-x, y) = -t(x, y) = t(x, y)$ . Thus, we have that  $(tx, y) = t(x, y)$  for all  $t \in \mathbb{Z}$ .

To proceed, we deal with  $t \in \mathbb{Q}$ . We have that  $t = \frac{m}{n}$  for some  $m, n \in \mathbb{Z}$ . Then we have that  $n(tx, y) = (ntx, y) = (mx, y) = m(x, y) = t(mx, y) = t(n(x, y))$ . Thus, we have that  $n(tx, y) = t(n(x, y))$ . Thus, we have that  $(tx, y) = t(x, y)$  for all  $t \in \mathbb{Q}$ .  $\ominus$

## 1.15 Compactness

### Definition 1.15.1: Subsequence

Let  $X$  be a set and  $f : \mathbb{N} \rightarrow X$  a sequence. Let  $g : \mathbb{N} \rightarrow \mathbb{N}$  be strictly increasing. Then  $f \circ g : \mathbb{N} \rightarrow X$  is a *subsequence* of  $f$ . We denote  $m_k$  as  $g(k)$ , so  $f(g(k)) = f(m_k) = x_{m_k}$ . So we denote the whole sequence as  $\{x_{m_k}\}_k$ .

### Definition 1.15.2: Sequentially Compact

Let  $(X, d)$  be a metric space.  $K \subseteq X$  is *sequentially compact* if every sequence  $\{x_n\}_n$  in  $K$  and there exists a subsequence  $\{x_{n_k}\}_k$  such that  $x_{n_k} \rightarrow x$  as  $k \rightarrow \infty$  for some  $x \in K$ .

**Example 1.15.1** ( $\mathbb{R}$ )

1.  $(0, 1]$  is not sequentially compact. Consider the sequence  $x_n = \frac{1}{n}$ . This sequence has no convergent subsequence that tends to 0 since 0 is not in the set. The issue is that it's not closed.
2.  $[0, \infty)$  is not sequentially compact. Consider the sequence  $x_n = n$ . This sequence has no convergent subsequence that tends to  $\infty$  since  $\infty$  is not in the set. So,  $[0, \infty)$  is not sequentially compact. The issue is that it's not bounded.

**Theorem 1.15.1**

Let  $(X, d)$  be a metric space. If  $K \subseteq X$  is sequentially compact, then  $K$  is closed and bounded.

*Proof.* Claim:  $K$  is closed. We want  $X \setminus K$  to be open. Let  $x \in X \setminus K$ . We want  $B(x, r) \subseteq X \setminus K$  for some  $r > 0$ . By contradiction, for all  $r > 0$ , assume  $\exists y \in B(x, r) \cap K$ . Take  $r = \frac{1}{m} \Rightarrow y_m \in B(x, \frac{1}{m}) \cap K$ .  $d(y_m, x) < \frac{1}{m} \rightarrow 0$ , so  $y_m \rightarrow x$ . But  $x \notin K$  even though  $y_m \in K$ . This is a contradiction, so  $K$  is closed.

Claim:  $K$  is bounded. By contradiction, assume  $K$  is not bounded. Let  $x_0 \in X$ . Then  $K \not\subseteq B(x_0, r)$  for any  $r > 0$ . Take  $r = n$ . Then  $\exists x_n \in K$  such that  $d(x_n, x_0) \geq n$ . So  $\{x_n\}_n \in K$ .  $K$  is sequentially compact, so there exists a subsequence  $\{x_{n_k}\}_k$  such that  $x_{n_k} \rightarrow x$  as  $k \rightarrow \infty$  for some  $x \in K$ . But  $n_k \leq d(x_{n_k}, x_0) \leq d(x_{n_k}, x) + d(x, x_0)$ . But  $d(x_{n_k}, x) \rightarrow 0$  as  $k \rightarrow \infty$ , so  $n_k \rightarrow \infty < d(x_{n_k}, x_0) \leq d(x, x_0)$  which is a fixed number, so we have a contradiction. As such,  $K$  is bounded. ☺

**Theorem 1.15.2**

Let  $K \subseteq \mathbb{R}^N$ . Then  $K$  is sequentially compact if and only if  $K$  is closed and bounded.

*Proof.* We just showed the first direction. So, we need to show that if  $K$  is closed and bounded, then  $K$  is sequentially compact.

So now, assume  $K$  is closed and bounded. Let  $\{x_n\}_n$  be a sequence in  $K$ . We want to show that there exists a subsequence  $\{x_{n_k}\}_k$  such that  $x_{n_k} \rightarrow x$  as  $k \rightarrow \infty$  for some  $x \in K$ .

Consider the set  $E = \{x_n : n \in \mathbb{N}\} \subseteq \mathbb{R}_N$ . We now case on whether  $E$  has infinitely many distinct points or not.

If  $E$  doesn't have infinitely many distinct points, there exists  $x \in K$  such that  $x_n = x$  for infinitely many  $n$ . Then  $x_{n_k} = x$  for all  $k$ , so  $x_{n_k} \rightarrow x$  as  $k \rightarrow \infty$ .

Now we consider the case where Bolzano-Weierstrass applies. By B-W,  $E$  has an accumulation point  $x \in \mathbb{R}^N$ . So we can find a subsequence  $\{x_{n_k}\}_k$  such that  $x_{n_k} \rightarrow x$  as  $k \rightarrow \infty$ . But  $x \in K$  because  $K$  is closed. Thus,  $K$  is sequentially compact. ☺

**Note:**

Let  $(X, \|\cdot\|)$  be a normed space. If every closed and bounded set is sequentially compact, then  $X$  has finite dimension.

**Exercise 1.15.1**

Recall  $l^\infty([0, 1]) = \{f : [0, 1] \rightarrow \mathbb{R} \text{ bounded}\}$ . Define  $\|f\|_\infty = \sup_{x \in [0, 1]} |f(x)|$ .  $B(0, 1) = \{g \in l^\infty([0, 1]) : \|g\|_\infty < 1\}$ . Prove that  $B(0, 1) = \{g \in l^\infty([0, 1]) : |g(x)| < 1 \ \forall x \in [0, 1]\}$ . Also prove that this not sequentially compact.



## 1.16 2/8 - Recitation

Let  $n \in \mathbb{N}$ ,  $x, y \in \mathbb{R}$ .

1. Prove that  $x^n - y^n = (x - y)(x^{n-1} + x^{n-2}y + \cdots + xy^{n-2} + y^{n-1})$ .

*Proof.* Base case:  $n = 1$  is trivial.

Now assume that for any  $n \in \mathbb{N}$ ,  $x^n - y^n = (x - y)(x^{n-1} + x^{n-2}y + \cdots + xy^{n-2} + y^{n-1})$ . We want to show that this is true for  $n+1$ . We have that  $x^{n+1} - y^{n+1} = x(x^n - y^n) + y^n(x - y) = x(x - y)(x^{n-1} + x^{n-2}y + \cdots + xy^{n-2} + y^{n-1}) + y^n(x - y)$ . Then we get  $(x - y)(x^n + x^{n-1}y + \cdots + xy^{n-1} + y^n) = (x - y)(x^n + x^{n-1}y + \cdots + xy^{n-1} + y^n)$ .  $\odot$

2. Prove that when  $|x - y| \leq 1$ , then  $|x^n - y^n| \leq n(1 + |x|)^{n-1}|x - y|$ .

*Proof.* Let  $|x - y| \leq 1$ . Then we have that  $|x^n - y^n| = |(x - y)(x^{n-1} + x^{n-2}y + \cdots + xy^{n-2} + y^{n-1})| \leq |x - y|(|x^{n-1}| + |x^{n-2}y| + \cdots + |xy^{n-2}| + |y^{n-1}|) \leq |x - y|(|x|^{n-1} + |x|^{n-2}|y| + \cdots + |x||y|^{n-2} + |y|^{n-1}) \leq |x - y|(|x|^{n-1} + |x|^{n-2}|y| + \cdots + |x||y|^{n-2} + |y|^{n-1}) \leq |x - y|(|x|^{n-1} + |x|^{n-2} + \cdots + |x| + 1) \leq n(1 + |x|)^{n-1}|x - y|$ .  $\odot$

3. Let  $E = \{x \in \mathbb{R} : x^n > 3\}$  for a fixed  $n$ . Prove that  $E$  is open.

*Proof.* Let  $x \in E$ . We want to show that there is an  $r > 0$  such that  $B(x, r) \subseteq E$ . Take  $r = \frac{x^n - 3}{n(1 + |x|)^{n-1}}$  and take  $y \in B(x, r)$ . Then  $|x - y| < r \Rightarrow |x^n| - |y^n| \leq |x^n - y^n| \leq n(1 + |x|)^{n-1}|x - y| < n(1 + |x|)^{n-1}r < x^n - 3$ . Then  $y^n \geq x^n - n(1 + |x|)^{n-1}r > 3$ . Thus,  $y \in E$ . Thus,  $B(x, r) \subseteq E$ . Thus,  $E$  is open.  $\odot$

4. Consider the space  $l^\infty([0, 1]) = \{f : [0, 1] \rightarrow \mathbb{R} \text{ bounded}\}$ . Define  $\|f\|_\infty = \sup_{x \in [0, 1]} |f(x)|$ .

Let  $E = \{f \in l^\infty([0, 1]) : f(x) > 0 \forall x \in [0, 1]\}$ . Prove that  $E$  is not open.

*Proof.* Consider

$$f(x) = \begin{cases} x & x \in [0, 1) \\ 1 & x = 1 \end{cases}$$

Then let  $r > 0$  and consider  $g(x) = f(x) \cdot \frac{r}{2}$ . Then  $g(x) \in B(f, r)$ . But  $g(x) \notin E$  because  $g(1) = \frac{r}{2}$ . Thus,  $B(f, r) \not\subseteq E$ . Thus,  $E$  is not open.  $\odot$

## 1.17 Limits

### Definition 1.17.1: Limits

Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces,  $E \subseteq X$ ,  $f : E \rightarrow Y$ . Let  $x_0 \in \text{acc } E$ .

Take  $l \in Y$ .  $l$  is the *limit* of  $f$  as  $x \rightarrow x_0$ . We write  $\lim_{x \rightarrow x_0} f(x) = l$  if for every  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $0 < d_X(x, x_0) < \delta \Rightarrow d_Y(f(x), l) < \epsilon$ . We can also write it as  $f(x) \rightarrow l$  as  $x \rightarrow x_0$ .

### Note:

Even if  $x_0 \in E$ , you don't take in the definition for the limit.

### Theorem 1.17.1

Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces,  $E \subseteq X$ ,  $f : E \rightarrow Y$ , and  $x_0 \in \text{acc } E$ . If  $\lim_{x \rightarrow x_0} f(x)$  exists, then it is unique.

*Proof.* Assume that  $\lim_{x \rightarrow x_0} f(x) = l$  and  $\lim_{x \rightarrow x_0} f(x) = m$ . Take  $\epsilon = \frac{d_Y(l, m)}{2} > 0$ . Then there exists  $\delta_1 > 0$  such that  $0 < d_X(x, x_0) < \delta_1 \Rightarrow d_Y(f(x), l) < \epsilon$ . There also exists  $\delta_2 > 0$  such that  $0 < d_X(x, x_0) < \delta_2 \Rightarrow d_Y(f(x), m) < \epsilon$ . Take  $\delta = \min(\delta_1, \delta_2)$ . Then  $0 < d_X(x, x_0) < \delta \Rightarrow d_Y(f(x), l) < \epsilon$  and  $d_Y(f(x), m) < \epsilon$ . Then  $d_Y(l, m) \leq d_Y(l, f(x)) + d_Y(f(x), m) < 2\epsilon = d_Y(l, m)$ . This is a contradiction, so  $l = m$ .  $\odot$

**Example 1.17.1 ( $\mathbb{R}^2$ )**

Take  $(x_0, y_0) \in \mathbb{R}^2$  and  $y_0 \neq 0$ . Compute

$$\lim_{(x,y) \rightarrow (x_0,y_0)} \frac{x}{y}$$

We want to show that this is  $\frac{x_0}{y_0}$ . We have the set  $E = \{(x, y) \in \mathbb{R}^2 : y \neq 0\}$ . We also know that  $(x_0, y_0) \in \text{acc } E$ . What we know is that  $(x, y) \rightarrow (x_0, y_0)$ :  $|x - x_0|$  and  $|y - y_0|$  are going to be small. Then

$$\begin{aligned} \left| f(x, y) - \frac{x_0}{y_0} \right| &= \left| \frac{x}{y} - \frac{x_0}{y_0} \right| \\ &= \left| \frac{xy_0 - x_0y_0}{yy_0} \right| \\ &= \left| \frac{xy_0 - x_0y_0 + x_0y_0 - x_0y}{yy_0} \right| \\ &= \left| \frac{y_0(x - x_0) + x_0(y_0 - y)}{yy_0} \right| \\ &\leq \frac{|y_0||x - x_0| + |x_0||y_0 - y|}{|y||y_0|} \\ &= \frac{|x - x_0|}{|y|} + \frac{|x_0||y_0 - y|}{|y||y_0|} \end{aligned}$$

Then we have  $\delta < \frac{|y_0|}{2}$ . If  $|y - y_0| < \delta < \frac{|y_0|}{2}$ , then we get  $|y| \geq \frac{|y_0|}{2} \Rightarrow \frac{1}{|y|} \leq \frac{2}{|y_0|}$ .

$$\frac{|x - x_0|}{|y|} + \frac{|x_0||y_0 - y|}{|y||y_0|} \leq \frac{2|x - x_0|}{|y_0|} + \frac{2|x_0||y_0 - y|}{|y_0|^2}$$

Take  $\delta = \min \left\{ \epsilon, \frac{|y_0|}{2} \right\} > 0$ . Then  $0 < \|(x, y) - (x_0, y_0)\| < \delta$ .

$$\begin{aligned} |x - x_0| &= \sqrt{(x - x_0)^2} \leq \sqrt{(x - x_0)^2 + (y - y_0)^2} \\ |y - y_0| &\leq \delta \end{aligned}$$

So,

$$\left| f(x, y) - \frac{x_0}{y_0} \right| < \epsilon \left( \frac{2}{|y_0|} + \frac{2}{|y_0|^2} \right)$$

Say you can prove that for every  $\epsilon > 0$ ,  $\exists \delta > 0$  such that

$$d(f(x), l) < \epsilon |\log(\epsilon)| \text{ for all } x \in E \text{ such that } 0 < d(x, x_0) < \delta$$

For every  $\eta > 0$  ("my epsilon"), since  $\lim_{\epsilon \rightarrow 0^+} \epsilon |\log(\epsilon)| = 0$ ,  $\exists \delta_1 > 0$  such that  $\epsilon |\log(\epsilon)| < \eta$  for all  $0 < \epsilon < \delta_1$ .

So given  $\eta > 0$ , take  $0 < \epsilon < \delta_1$ . Find  $\eta$  from  $d(f(x), l) < \epsilon |\log(\epsilon)| < \eta$  for all  $x \in E$  such that  $0 < d(x, x_0) < \delta$ . This means that

$$d_Y(f(x), l) < \epsilon |\log(\epsilon)| < \eta$$

for all  $x \in E$ ,  $0 < d(x, x_0) < \delta$ . Thus,  $\lim_{x \rightarrow x_0} f(x) = l$ .

## 1.18 Limits Continued

### Definition 1.18.1: Restriction

Assume that  $\lim_{x \rightarrow x_0} f(x) = l$  exists. Let  $F \subseteq E$  such that  $x_0 \in \text{acc } F$ . The function  $f : F \rightarrow Y$  is called the *restriction* of  $f$  to  $F$ . It is denoted as  $f|_F$ .

### Note:

If  $\lim_{x \rightarrow x_0} f(x) = l$ , then  $\lim_{x \rightarrow x_0} f|_F(x) = l$ .

So to prove that the limit does not exist, you can conjure up two restrictions of the function and show that the limits are different.

### Example 1.18.1 (Limits that don't exist)

Consider  $\lim_{x \rightarrow 0} \sin\left(\frac{1}{x}\right)$ . We can find two restrictions of the function and show that the limits are different.

- $\frac{1}{x} = 2\pi n + \frac{\pi}{2}$
- $x_n = \frac{1}{2\pi n + \frac{\pi}{2}}$

So we have:

- $\sin x_n = \sin\left(\frac{\pi}{2} + 2\pi n\right) = 1$
- $\sin x_n = \sin\left(\frac{1}{2\pi n}\right) = 0$

Thus, the limit does not exist.

### Exercise 1.18.1 TODO in Recitation

- $\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2+y^2}$  (no)
- $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2y}{x^2+y^2}$  (yes, 0)
- $\lim_{(x,y) \rightarrow (0,0)} \frac{x^{1000000000}y}{y - \sin(x)}$  (no)

Now we talk about the composition of limits.

### Example 1.18.2

Consider

$$g(y) = \begin{cases} 1 & y \neq 0 \\ 2 & y = 0 \end{cases}.$$

The limit of  $g(y)$  as  $y \rightarrow 0$  is 1. Now consider  $f(x) = 0$ . The limit of  $f(x)$  as  $x \rightarrow x_0$  is 0. Now consider  $g(f(x))$ . The limit of  $g(f(x))$  as  $x \rightarrow x_0$  is 2.

### Theorem 1.18.1 Composition of Limits

Let  $(X, d_X)$ ,  $(Y, d_Y)$ , and  $(Z, d_Z)$  be metric spaces,  $E \subseteq X$ ,  $F \subseteq Y$ ,  $f : E \rightarrow F$ ,  $g : F \rightarrow Z$ , and  $x_0 \in \text{acc } E$ . Assume there exists  $\lim_{x \rightarrow x_0} f(x) = l \in Y$ . Assume  $l \in \text{acc } F$  and that there is  $\lim_{y \rightarrow l} g(y) = L \in Z$ . Assume that either  $f(x) \neq l$  for all  $x \in E$  or  $l \in F$  and  $g(l) = L$ . Then there is  $\lim_{x \rightarrow x_0} g(f(x)) = L$ .

*Proof.* Since  $\lim_{y \rightarrow l} g(y) = L$ , there for every  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $d_Z(g(y), L) < \epsilon$  for all  $y \in F$  with  $0 < d_Y(y, l) < \delta$ . We would like to take  $y = f(x)$ . Use  $\delta$  as “my epsilon” for the definition of the limit of  $f(x)$ . Then to find  $\eta > 0$  such that  $d_X(f(x), l) < \delta$  for all  $x \in E$  with  $0 < d_X(x, x_0) < \eta$ . Now we split into cases:

- Assume  $f(x) \neq l$  for all  $x \in E$ . Then  $0 < d_Y(f(x), l)$  so we can take  $y = f(x)$  to get  $d_Z(g(f(x)), L) < \epsilon$  for all  $x \in E$  with  $0 < d_X(x, x_0) < \eta$ . This means that there exists  $\lim_{x \rightarrow x_0} g(f(x)) = L$ .
- Assume  $l \in F$  and  $g(l) = L$ . If  $f(x) = l$ , then  $d_Z(g(f(x)), L) = d_Z(g(l), L) = 0$  for all  $x \in E$  with  $0 < d_X(x, x_0) < \eta$ . If  $f(x) \neq l$ , then take  $y = f(x)$  to get  $0 < d_Y(f(x), l)$  so we can take  $y = f(x)$  to get  $d_Z(g(f(x)), L) < \epsilon$  for all  $x \in E$  with  $0 < d_X(x, x_0) < \eta$ . This means that there exists  $\lim_{x \rightarrow x_0} g(f(x)) = L$ . This means that there exists  $\lim_{x \rightarrow x_0} g(f(x)) = L$ .

⊖

### Corollary 1.18.1 Limits of the Sum/Products/Quotients

Let  $(X, d)$  be a metric space and  $E \subseteq X$ . Then take  $f : E \rightarrow \mathbb{R}$  and  $g : E \rightarrow \mathbb{R}$ . Let  $x_0 \in X$  and  $x_0 \in \text{acc } E$ . Assume  $\lim_{x \rightarrow x_0} f(x) = l$  and  $\lim_{x \rightarrow x_0} g(x) = m$ . Then we have the following results:

- $\lim_{x \rightarrow x_0} (f + g)(x) = l + m$ .
- $\lim_{x \rightarrow x_0} (f \cdot g)(x) = l \cdot m$ .
- $\lim_{x \rightarrow x_0} \frac{f}{g}(x) = \frac{l}{m}$ .

*Proof.* We can use the composition of limits to prove this. We'll just proceed with the quotient case. Consider  $x \rightarrow (f(x), g(x))$ . Then consider the function that takes  $(s, t) \rightarrow \frac{s}{t}$  and call it  $h$ . Then we have  $\frac{f(x)}{g(x)} = h(f(x), g(x))$ . We then have  $\lim_{x \rightarrow x_0} (f(x), g(x)) = (l, m)$  and  $\lim_{(s,t) \rightarrow (l,m)} h(s, t) = \frac{l}{m} = h(l, m)$ . So now we can use the composition of limits to get  $\lim_{x \rightarrow x_0} \frac{f}{g}(x) = \frac{l}{m}$ .

The other two cases are similar. For products, you need to show that the limit as  $(x, y) \rightarrow (x_0, y_0)$  of  $xy$  is  $x_0 y_0$  and similarly for sum. ⊖

## 1.19 Squeeze Theorem

### Theorem 1.19.1 Squeeze Theorem

Let  $(X, d_X)$  be a metric space,  $E \subseteq X$ ,  $f : E \rightarrow \mathbb{R}$ ,  $g : E \rightarrow \mathbb{R}$ , and  $h : E \rightarrow \mathbb{R}$ . Let  $x_0 \in \text{acc } E$  and have  $f \leq g \leq h$ . Assume that  $\lim_{x \rightarrow x_0} f(x) = l = \lim_{x \rightarrow x_0} h(x)$ . Then  $\lim_{x \rightarrow x_0} g(x) = l$ .

*Proof.* Assume  $\lim_{x \rightarrow x_0} f(x) = l = \lim_{x \rightarrow x_0} h(x)$ . Then for every  $\epsilon > 0$ , there exists  $\delta_1 > 0$  such that  $0 < d_X(x, x_0) < \delta_1 \Rightarrow |f(x) - l| < \epsilon$  and  $0 < d_X(x, x_0) < \delta_2 \Rightarrow |h(x) - l| < \epsilon$ . Take  $\delta = \min(\delta_1, \delta_2)$  and  $x \in E$  with  $0 < d_X(x, x_0) < \delta$ . Then  $l - \epsilon < f(x) < g(x) < h(x) < l + \epsilon$ . Then  $|g(x) - l| < \epsilon$  for all  $x \in E$  with  $0 < d_X(x, x_0) < \delta$ . Thus,  $\lim_{x \rightarrow x_0} g(x) = l$ . ⊖

### Example 1.19.1

$$\lim_{x \rightarrow 0} |x|^a \sin \frac{1}{x} = 0$$

for all  $a > 0$ .

Let  $Q > 0$ . Then  $0 \leq |x|^Q \sin \frac{1}{x} \leq |x|^Q$ . Since both sides tend to 0 as  $x \rightarrow 0$ , then the middle does as well.

### Definition 1.19.1: Increasing

$f : E \rightarrow \mathbb{R}$  is *increasing* if  $f(x) \leq f(y)$  for all  $x \leq y$ . It is *strictly increasing* if  $f(x) < f(y)$  for all  $x < y$ .

**Definition 1.19.2: Decreasing**

$f : E \rightarrow \mathbb{R}$  is *decreasing* if  $f(x) \geq f(y)$  for all  $x \leq y$ . It is *strictly decreasing* if  $f(x) > f(y)$  for all  $x < y$ .

**Definition 1.19.3: Divergent**

Let  $(X, d_X)$  be a metric space with  $E \subseteq X$ ,  $x_0 \in \text{acc } E$ , and  $f : E \rightarrow \mathbb{R}$ . We say that  $f$  diverges to  $+\infty$  as  $x \rightarrow x_0$  if for every  $M > 0 \in \mathbb{R}$ , there exists  $\delta > 0$  such that  $f(x) > M$  for all  $x \in E$  with  $0 < d_X(x, x_0) < \delta$ . We say that  $f$  diverges to  $-\infty$  as  $x \rightarrow x_0$  if for every  $M < 0 \in \mathbb{R}$ , there exists  $\delta > 0$  such that  $f(x) < M$  for all  $x \in E$  with  $0 < d_X(x, x_0) < \delta$ .

**Theorem 1.19.2**

Let  $E \subseteq \mathbb{R}$  and  $f : E \rightarrow \mathbb{R}$  be increasing. Let  $x_0 \in \mathbb{R}$ . Assume  $x_0$  is an accumulation point of  $E \cap (-\infty, x_0)$ . Then there is

$$\lim_{x \rightarrow x_0^-} f(x) = \sup_{E \cap (-\infty, x_0)} f(x)$$

Now if  $x_0$  is an accumulation point of  $E \cap (x_0, \infty)$ , then there is

$$\lim_{x \rightarrow x_0^+} f(x) = \inf_{E \cap (x_0, \infty)} f(x)$$

*Proof.*

- Case 1: Assume  $f$  is bounded from above on  $E \cap (-\infty, x_0)$ . Let  $l = \sup_{E \cap (-\infty, x_0)} f(x)$ . Then for every  $\epsilon > 0$ , there exists  $x_1 \in E \cap (-\infty, x_0)$  such that  $l - \epsilon < f(x_1) \leq l$ . Then for every  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $l - \epsilon < f(x_1) \leq l$ . Take  $\delta = x_0 - x_1 > 0$ . Let  $x \in E$  with  $x_0 - \delta < x < x_0$ . Since  $f$  is increasing, we have  $l - \epsilon < f(x_1) \leq f(x) \leq l < l + \epsilon$ . Thus,  $\lim_{x \rightarrow x_0^-} f(x) = l$ .
- Case 2: If  $f$  is not bounded from above, then for every  $M > 0$ , there exists  $x_1 \in E \cap (-\infty, x_0)$  such that  $f(x_1) > M$ . Let  $\delta = x_0 - x_1 > 0$ . Then for every  $x \in E$  with  $x_1 = x_0 - \delta < x < x_0$ , we have  $f(x) \geq f(x_1) > M$ . Thus,  $\lim_{x \rightarrow x_0^-} f(x) = +\infty$ .

The other case is similar. ⊕

**Definition 1.19.4: Infinite Sum**

Let  $X$  be a set and take  $f : X \rightarrow [0, \infty]$ . The *infinite sum* is defined as:

$$\sum_{x \in X} f(x) = \sup \left\{ \sum_{x \in F} f(x) : F \subseteq X \text{ finite} \right\}.$$

**Lemma 1.19.1**

Let  $X$  be nonempty with  $f : X \rightarrow [0, \infty]$ . Assume that  $\sum_{x \in X} f(x) < \infty$ . Then  $\{x \in X : f(x) > 0\}$  is countable.

*Proof.* Take  $n \in \mathbb{N}$  and define  $X_n = \{x \in X : f(x) \geq \frac{1}{n}\}$ . Let  $E \subseteq X_n$  be finite. Then  $\frac{1}{n}|E| < \sum_{x \in E} f(x) \leq M$ . Then  $|E| < nM$ . Thus,  $X_n$  is countable. Then  $\bigcup_{n \in \mathbb{N}} X_n = \{x \in X : f(x) > 0\}$  is countable. ⊕

**Theorem 1.19.3**

Take  $I \subseteq \mathbb{R}$  to be an interval with  $f : I \rightarrow \mathbb{R}$  increasing. Then for all but countably many  $x_0 \in I$ , there is  $\lim_{x \rightarrow x_0^-} f(x) = \lim_{x \rightarrow x_0^+} f(x) = f(x_0)$ .

*Proof.* Let  $x_0 \in I$ .

⊖