

MATHEMATICAL ANALYSIS I TUTORING

4TH WEEK

GLOBAL PROPERTIES OF CONTINUOUS FUNCTIONS - LANDAU SYMBOLS - LOCAL COMPARISON INFINITESIMAL, INFINITE FUNCTIONS, ORDER AND PRINCIPAL PART - ASYMPTOTES

PROPOSED EXERCISES

1. For each of the following statements, say if it is true or false, justifying by means of theoretical references.

- (a) If a function is continuous in x_0 , then it is bounded in every neighborhood of x_0 .

A function $f(x)$ is continuous in x_0 if $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, that is, if for every neighborhood I of $f(x_0)$ **there exists a neighborhood J of x_0** , such that for every $x \in J$, it holds that $f(x) \in I$; hence $f(x)$ is bounded if $x \in J$.

It is not guaranteed that f is bounded in **every** neighborhood of x_0 : it is sufficient to consider the following counterexample: $f(x) = \tan x$, it is continuous in $x_0 = 0$ but it is not bounded in every neighborhood of x_0 (take for instance a neighborhood with radius 5).

Therefore the statement is FALSE.

- (b) If a function is continuous in $I=[a, b]$, then it admits maximum in I .

Weierstrass Theorem - Let f be a continuous function in a closed and bounded interval $[a, b]$. Then f is bounded in $[a, b]$ and it admits maximum and minimum.

Therefore the statement is TRUE by Weierstrass Theorem.

- (c) Let $f : [-2, 3] \subseteq \mathbb{R} \rightarrow \mathbb{R}$, continuous and such that $f(-2) = -1$ and $f(3) = 2$; then necessarily the equation $f(x) = \lambda$ admits at least a solution in $-2 \leq \lambda \leq 3$.

This statement applies (in a wrong way) the Mean Values Theorem (Corollary of Existence of Zeros Theorem) that states:

Mean Values Theorem - Let f be a continuous function on a closed and bounded interval $[a, b]$. Then $f(x)$ assumes all the values between $f(a)$ and $f(b)$.

The equation $f(x) = \lambda$ admits at least a solution if $-2 \leq \lambda \leq 3$ is equivalent to say that $f(x)$ assumes any value λ such that $-2 \leq \lambda \leq 3$.

Therefore the statement is FALSE, since the function $f(x)$ takes all the values $-1 \leq \lambda \leq 2$, and not necessarily the values $-2 \leq \lambda \leq 3$.

- (d) There exists a function $f : [3, 7] \rightarrow \mathbb{R}$ continuous and surjective.

FALSE. If the function is continuous on $[3, 7]$ then the function is bounded on such interval (by Weierstrass Theorem, it assumes maximum M and minimum m and thus $\text{Im}(f) = [m, M]$). It follows that the image of the function on such interval cannot be the whole set \mathbb{R} , i.e. the function cannot be surjective.

- (e) The function $f(x) = \begin{cases} x \sin \frac{1}{x} & x \neq 0 \\ 0 & x = 0 \end{cases}$ is not continuous in $x = 0$.

The function is continuous for $x \neq 0$ by composition and product of continuous functions. We have to investigate the continuity at the origin, that is, we have to verify the equality $\lim_{x \rightarrow 0} f(x) = f(0)$. By direct computation:


$$\lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0$$

applying the substitution $\frac{1}{x} = t$, we reduce to the known limit $\lim_{t \rightarrow +\infty} \frac{\sin t}{t} = 0$. Hence the function is continuous on \mathbb{R} .

Now we investigate if the function is bounded or not.
 Since f is even, we study the function only for $x > 0$.
 Observe that

- $-1 \leq \sin \frac{1}{x} \leq 1, \forall x \in \mathbb{R}$
- $\lim_{x \rightarrow +\infty} x \sin \frac{1}{x} = \lim_{t \rightarrow 0} \frac{\sin t}{t} = 1.$
- We have that $-1 \leq x \sin \frac{1}{x} \leq 1$

Indeed, $\forall t \in \mathbb{R}, -t \leq \sin t \leq t$; thus $-\frac{1}{x} \leq \sin \frac{1}{x} \leq \frac{1}{x}$, and finally $-1 \leq x \sin \frac{1}{x} \leq 1$.



We can conclude that the function is continuous and bounded, therefore the statement is TRUE.

2. a) Prove that for $x \rightarrow 0$, given two powers of x , the negligible one has higher exponent.

$$\lim_{x \rightarrow 0} \frac{x^n}{x^m} = \lim_{x \rightarrow 0} x^{n-m} = 0 \Leftrightarrow n - m > 0 \Leftrightarrow n > m$$

- b) Prove that for $x \rightarrow \pm\infty$, between two powers of x , the negligible one has lower degree.

$$\lim_{x \rightarrow \pm\infty} \frac{x^n}{x^m} = \lim_{x \rightarrow \pm\infty} \frac{1}{x^{m-n}} = 0 \Leftrightarrow m - n > 0 \Leftrightarrow n < m$$

3. Verificare che per $x \rightarrow +\infty$, $2^x + 2^{-x} = o(2^{3x} + 1)$.

Per $x \rightarrow x_0$ $f(x) = o(g(x))$ se $\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 0$.

Il limite $\lim_{x \rightarrow +\infty} \frac{2^x + 2^{-x}}{2^{3x} + 1}$ presenta la forma indeterminata $\frac{\infty}{\infty}$.

Utilizziamo il principio di eliminazione dei termini trascurabili: poiché, per $x \rightarrow +\infty$, si ha $2^{-x} = o(2^x)$ e $1 = o(2^{3x})$, il limite si semplifica:

$$\lim_{x \rightarrow +\infty} \frac{2^x + 2^{-x}}{2^{3x} + 1} = \lim_{x \rightarrow +\infty} \frac{2^x}{2^{3x}} = \lim_{x \rightarrow +\infty} \frac{1}{2^{2x}} = 0$$

4. Given $f(x) = x^3$, $g(x) = x^3 - x$, $h(x) = x^3 - 1$.

Verify that for $x \rightarrow +\infty$, $f \sim g \sim h$; say if $f - g \sim f - h$.

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow +\infty} \frac{x^3}{x^3 - x} = \lim_{x \rightarrow +\infty} \frac{x^3}{x^3 \left(1 - \frac{1}{x^2}\right)} = 1 \Rightarrow f \sim g$$

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{h(x)} = \lim_{x \rightarrow +\infty} \frac{x^3}{x^3 - 1} = \lim_{x \rightarrow +\infty} \frac{x^3}{x^3 \left(1 - \frac{1}{x^3}\right)} = 1 \Rightarrow f \sim h$$

$$\lim_{x \rightarrow +\infty} \frac{g(x)}{h(x)} = \lim_{x \rightarrow +\infty} \frac{x^3 - x}{x^3 - 1} = \lim_{x \rightarrow +\infty} \frac{x^3 \left(1 - \frac{1}{x^2}\right)}{x^3 \left(1 - \frac{1}{x^3}\right)} = 1 \Rightarrow g \sim h$$

$$\lim_{x \rightarrow +\infty} \frac{f(x) - g(x)}{f(x) - h(x)} = \lim_{x \rightarrow +\infty} \frac{x^3 - (x^3 - x)}{x^3 - (x^3 - 1)} = \lim_{x \rightarrow +\infty} \frac{x}{1} = +\infty \Rightarrow g \not\sim h$$

5. For each couple of functions, check if they are infinite or infinitesimal with the same order:

- $f_1(x) = \sqrt{2x^2 + 3x + 5}, \quad f_2(x) = 2x - 1, \quad \text{for } x \rightarrow +\infty$

$$\begin{aligned} f_1(x) &= \sqrt{2x^2 + 3x + 5} = \sqrt{x^2 \left(2 + 3\frac{x}{x^2} + \frac{5}{x^2}\right)} \\ &= |x| \sqrt{2 + 3\frac{1}{x} + \frac{5}{x^2}} \\ &\sim \sqrt{2}|x| = \sqrt{2}x, \quad \text{for } x \rightarrow +\infty \\ f_2(x) &\sim 2x, \quad \text{for } x \rightarrow +\infty \end{aligned}$$

Hence $f_1(x)$ and $f_2(x)$ are infinite functions with same order (not equivalent), for $x \rightarrow +\infty$.

- $g_1(x) = \sqrt{x+3} - \sqrt{3}, \quad g_2(x) = 2x, \quad \text{for } x \rightarrow 0$

$$\lim_{x \rightarrow 0} \frac{g_1(x)}{g_2(x)} = \lim_{x \rightarrow 0} \frac{\sqrt{x+3} - \sqrt{3}}{2x} = \lim_{x \rightarrow 0} \frac{x+3-3}{2x(\sqrt{x+3} + \sqrt{3})} = \lim_{x \rightarrow 0} \frac{x}{2x(\sqrt{x+3} + \sqrt{3})} = \lim_{x \rightarrow 0} \frac{1}{2(\sqrt{x+3} + \sqrt{3})} = \frac{1}{2\sqrt{3}}$$

Therefore $g_1(x)$ and $g_2(x)$ have the same order, and thus they are infinitesimal functions with same order, for $x \rightarrow 0$.

6. As $\alpha \in \mathbb{R}$, compute the limit:

$$\lim_{x \rightarrow 0^+} \frac{1 - e^x}{\frac{2}{\pi} \arctan x^\alpha}$$

Recall that, $e^t - 1 \sim t$ and $\arctan t \sim t$, for $t \rightarrow 0$, by substitution:

$$\lim_{x \rightarrow 0^+} \frac{1 - e^x}{\frac{2}{\pi} \arctan x^\alpha} = \lim_{x \rightarrow 0^+} \frac{-x}{\frac{2}{\pi} x^\alpha} = -\frac{\pi}{2} \lim_{x \rightarrow 0^+} x^{1-\alpha} = \begin{cases} -\frac{\pi}{2} & \text{if } \alpha = 1 \\ -\infty & \text{if } \alpha > 1 \\ 0 & \text{if } \alpha < 1 \end{cases}$$

7. Compute the following limits:

a) $\boxed{\lim_{x \rightarrow 0} \frac{1 - \cos 5x}{\sin^2 3x}}$

Recall that $\sin t \sim t$ and $1 - \cos t \sim \frac{1}{2}t^2$, for $t \rightarrow 0$. By substitution:

$$\lim_{x \rightarrow 0} \frac{1 - \cos 5x}{\sin^2 3x} = \lim_{x \rightarrow 0} \frac{\frac{1}{2}(5x)^2}{(3x)^2} = \lim_{x \rightarrow 0} \frac{\frac{1}{2}25x^2}{9x^2} = \frac{25}{18}$$

b) $\boxed{\lim_{x \rightarrow 0} \frac{2x \cos x - 2x}{x \sin^2 x}}$

Recall that $\sin t \sim t$ and $1 - \cos t \sim \frac{1}{2}t^2$, for $t \rightarrow 0$. By substitution:

$$\lim_{x \rightarrow 0} \frac{2x \cos x - 2x}{x \sin^2 x} = \lim_{x \rightarrow 0} \frac{2x(\cos x - 1)}{x \cdot x^2} = \lim_{x \rightarrow 0} \frac{2x \left(-\frac{1}{2}x^2\right)}{x^3} = \lim_{x \rightarrow 0} \frac{-x^3}{x^3} = -1$$

c) $\lim_{x \rightarrow 0} \frac{1 - \cos^4 x}{x \tan x}$

Recall that $\tan t \sim t$ and $1 - \cos t \sim \frac{1}{2}t^2$, for $t \rightarrow 0$. By substitution:

$$\lim_{x \rightarrow 0} \frac{1 - \cos^4 x}{x \tan x} = \lim_{x \rightarrow 0} \frac{(1 - \cos x)(1 + \cos x)(1 + \cos^2 x)}{x \cdot x} = 4 \lim_{x \rightarrow 0} \frac{(1 - \cos x)}{x^2} = 4 \lim_{x \rightarrow 0} \frac{\frac{1}{2}x^2}{x^2} = 2$$

d) $\lim_{x \rightarrow 0} \frac{\tan x - \sin x}{x^3}$

Recall that $\sin t \sim t$ and $1 - \cos t \sim \frac{1}{2}t^2$, per $t \rightarrow 0$. By substitution:

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\tan x - \sin x}{x^3} &= \lim_{x \rightarrow 0} \frac{\frac{\sin x}{\cos x} - \sin x}{x^3} \\ &= \lim_{x \rightarrow 0} \frac{\sin x \left(\frac{1}{\cos x} - 1 \right)}{x^3} \\ &= \lim_{x \rightarrow 0} \frac{\sin x (1 - \cos x)}{\cos x x^3} \\ &= \lim_{x \rightarrow 0} \frac{x \cdot \frac{1}{2}x^2}{x^3} \\ &= \lim_{x \rightarrow 0} \frac{x \frac{1}{2}x^2}{x^3} = \frac{1}{2} \end{aligned}$$

e) $\lim_{x \rightarrow 0} \frac{(1 + \sin x)^5 - 1}{1 - \cos \sqrt{x}}$

Recall that $(1 + t)^\alpha - 1 \sim \alpha t$, $1 - \cos t \sim \frac{1}{2}t^2$ and $\sin t \sim t$, for $t \rightarrow 0$. By substitution:

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{(1 + \sin x)^5 - 1}{1 - \cos \sqrt{x}} &= \lim_{x \rightarrow 0} \frac{\frac{1}{5} \sin x}{\frac{1}{2}(\sqrt{x})^2} \\ &= \lim_{x \rightarrow 0} \frac{\frac{1}{5}x}{\frac{1}{2}x} = 10 \end{aligned}$$

f) $\lim_{x \rightarrow 1} \frac{\sqrt{2 - x^2} - 1}{(x - 1)^2}$

If $x \rightarrow 1$, then $x - 1 \rightarrow 0$; we apply the substitution $t = x - 1$ (and thus $x = t + 1$), and we recall that $(1 + t)^\alpha - 1 \sim \alpha t$, for $t \rightarrow 0$, then:

$$\begin{aligned} \lim_{x \rightarrow 1} \frac{\sqrt{2 - x^2} - 1}{(x - 1)^2} &= \lim_{t \rightarrow 0} \frac{\sqrt{2 - (t + 1)^2} - 1}{t^2} \\ &= \lim_{t \rightarrow 0} \frac{\sqrt{2 - t^2 - 2t - 1} - 1}{t^2} \\ &= \lim_{t \rightarrow 0} \frac{(1 + (-t^2 - 2t))^{1/2} - 1}{t^2} \\ &= \lim_{t \rightarrow 0} \frac{\frac{1}{2}(-t^2 - 2t)}{t^2} \\ &= \lim_{t \rightarrow 0} \frac{\frac{1}{2}(-2t)}{t^2} \\ &= \lim_{t \rightarrow 0} \frac{-1}{t} \end{aligned}$$

The limit does not exist.

g) $\lim_{x \rightarrow \frac{1}{2}} \frac{3^{2x-1} - 1}{\sin(2x - 1)}$

If $x \rightarrow \frac{1}{2}$, then $x - \frac{1}{2} \rightarrow 0$; imposing $t = x - \frac{1}{2}$, (and thus $x = t + \frac{1}{2}$), we have that $t \rightarrow 0$; therefore:

$$\begin{aligned} \lim_{x \rightarrow \frac{1}{2}} \frac{3^{2x-1} - 1}{\sin(2x-1)} &= \lim_{x \rightarrow \frac{1}{2}} \frac{3^{2(x-\frac{1}{2})} - 1}{\sin(2(x-\frac{1}{2}))} \\ &= \lim_{t \rightarrow 0} \frac{3^{2t} - 1}{\sin(2t)} \\ &= \lim_{t \rightarrow 0} \frac{2t \log(3)}{2t} = \log(3) \end{aligned}$$

(we applied the equivalences, for $t \rightarrow 0$, $a^t - 1 \sim t \log a$ and $\sin t \sim t$)

h) $\boxed{\lim_{x \rightarrow 0} \frac{\sqrt{2x^3} - x^6}{4x^6 - \sqrt{x^4 + x^3}}}$

It holds $x^6 = o(x^{3/2})$ for $x \rightarrow 0$ and $x^4 = o(x^3)$ for $x \rightarrow 0$; therefore we eliminate the negligible terms as follows:

$$\lim_{x \rightarrow 0} \frac{\sqrt{2x^3} - x^6}{4x^6 - \sqrt{x^4 + x^3}} = \lim_{x \rightarrow 0} \frac{\sqrt{2}x^{3/2}}{4x^6 - \sqrt{x^3}} = \lim_{x \rightarrow 0} \frac{\sqrt{2}x^{3/2}}{-x^{3/2}} = -\sqrt{2}$$

i) $\boxed{\lim_{x \rightarrow 0} \frac{\sin 3x + x^4}{6x + 5 \log(1 + x^2)}}$

It holds $x^4 = o(x)$ for $x \rightarrow 0$, and $\log(1 + 5x^2) = o(x)$ for $x \rightarrow 0$ (since $\log(1 + 5x^2) \sim 5x^2$); therefore

$$\lim_{x \rightarrow 0} \frac{\sin 3x + x^4}{6x + 5 \log(1 + x^2)} = \lim_{x \rightarrow 0} \frac{3x}{6x} = \frac{1}{2}$$

j) $\boxed{\lim_{x \rightarrow 0} \frac{1 - \cos(x^3 + 2x^2)}{x^4 - x^7}}$

It holds $x^7 = o(x^4)$ for $x \rightarrow 0$ and $x^3 = o(x^2)$ for $x \rightarrow 0$; moreover $1 - \cos t \sim \frac{1}{2}t^2$ for $x \rightarrow 0$; hence:

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x^3 + 2x^2)}{x^4 - x^7} = \lim_{x \rightarrow 0} \frac{1 - \cos(2x^2)}{x^4} = \lim_{x \rightarrow 0} \frac{\frac{1}{2}(4x^4)}{x^4} = 2$$

k) $\boxed{\lim_{x \rightarrow 0} \frac{\log(2 - \cos x)}{\sin^2 x}}$

Recall that $\sin t \sim t$, $\log(1 + t) \sim t$ and $1 - \cos t \sim \frac{1}{2}t^2$, for $t \rightarrow 0$. It holds:

$\log(2 - \cos x) = \log(1 + (1 - \cos x)) \sim (1 - \cos x)$ (set $t = 1 - \cos x$, then if $x \rightarrow 0$ then $t \rightarrow 0$); thus:

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\log(2 - \cos x)}{\sin^2 x} &= \lim_{x \rightarrow 0} \frac{\log(1 + (1 - \cos x))}{x^2} \\ &= \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} \\ &= \lim_{x \rightarrow 0} \frac{\frac{1}{2}x^2}{x^2} = \frac{1}{2} \end{aligned}$$

l) $\boxed{\lim_{x \rightarrow +\infty} 2x(\log(x+1) - \log x)}$ Recall that $\lim_{x \rightarrow +\infty} \left(1 + \frac{1}{x}\right)^x = e$, it holds:

$$\lim_{x \rightarrow +\infty} 2x(\log(x+1) - \log x) = \lim_{x \rightarrow +\infty} 2x \log\left(\frac{x+1}{x}\right) = \lim_{x \rightarrow +\infty} 2 \log\left(1 + \frac{1}{x}\right)^x = 2$$

9. Verify that $f(x) = \sqrt{x+5} - \sqrt{5}$ and $g(x) = \sqrt{x+7} - \sqrt{7}$ are infinitesimal with the same order for $x \rightarrow 0$ and find $k \in \mathbb{R} : f \sim kg \ (x \rightarrow 0)$. Find the principal part of $f(x)$ for $x \rightarrow 0$, with respect to the infinitesimal test function $\varphi(x) = x$.

Remember that for $t \rightarrow 0$, it holds $(1+t)^\alpha - 1 \sim \alpha t$, and thus $\sqrt{1+t} - 1 \sim \frac{1}{2}t$, we have:

$$f(x) = \sqrt{x+5} - \sqrt{5} = \sqrt{5 \left(1 + \frac{x}{5}\right)} - \sqrt{5} = \sqrt{5} \left(\sqrt{1 + \frac{x}{5}} - 1 \right) \sim \sqrt{5} \left(\frac{1}{2} \frac{x}{5} \right), \text{ for } x \rightarrow 0$$

Therefore $f(x) \sim \frac{\sqrt{5}}{10} x$, for $x \rightarrow 0$

$$g(x) = \sqrt{x+7} - \sqrt{7} = \sqrt{7 \left(1 + \frac{x}{7}\right)} - \sqrt{7} = \sqrt{7} \left(\sqrt{1 + \frac{x}{7}} - 1 \right) \sim \sqrt{7} \left(\frac{1}{2} \frac{x}{7} \right), \text{ for } x \rightarrow 0$$

Hence $g(x) \sim \frac{\sqrt{7}}{14} x$, for $x \rightarrow 0$.

$$f \sim kg \ (x \rightarrow 0) \Leftrightarrow \lim_{x \rightarrow 0} \frac{f(x)}{kg(x)} = 1 \Leftrightarrow \lim_{x \rightarrow 0} \frac{\frac{\sqrt{5}}{10} x}{k \frac{\sqrt{7}}{14} x} = 1 \Leftrightarrow \frac{\frac{\sqrt{5}}{10}}{k \frac{\sqrt{7}}{14}} = 1 \Leftrightarrow k = \frac{\sqrt{5}}{5} \frac{7}{\sqrt{7}} = \frac{\sqrt{7}}{\sqrt{5}}$$

The principal part of $f(x)$ as $x \rightarrow 0$ w.r.t. the test function $\varphi(x) = x$ is $p(x) = \frac{\sqrt{5}}{10} x$

Find the order of infinitesimal and the principal part w.r.t. (with respect to) the test function $\varphi(x) = \frac{1}{x}$, for $x \rightarrow +\infty$, for the functions:

$$f_1(x) = \frac{3x^2 + \sqrt[3]{x}}{2x^3}$$

$$f_1(x) = \frac{3x^2 + \sqrt[3]{x}}{2x^3} \sim \frac{3x^2}{2x^3}, \text{ for } x \rightarrow +\infty$$

Therefore $f_1(x) \sim \frac{3}{2x}$, for $x \rightarrow +\infty$

W.r.t. (with respect to) the infinitesimal test function $\varphi(x) = \frac{1}{x}$, for $x \rightarrow +\infty$, the order of infinitesimal of $f_1(x)$ is 1 and its principal part is $p(x) = \frac{3}{2} \frac{1}{x}$.

$$f_2(x) = \sqrt{x^2 + 2} - x$$

$$f_2(x) = \sqrt{x^2 + 2} - x = \sqrt{x^2 \left(1 + \frac{2}{x^2}\right)} - x = |x| \sqrt{1 + \frac{2}{x^2}} - x = x \left(\sqrt{1 + \frac{2}{x^2}} - 1 \right) \sim x \left(\frac{1}{2} \frac{2}{x^2} \right), \text{ as } x \rightarrow +\infty$$

Thus $f_2(x) \sim \frac{1}{x}$, as $x \rightarrow +\infty$.

W.r.t. the infinitesimal test function $\varphi(x) = \frac{1}{x}$, as $x \rightarrow +\infty$, the order of infinitesimal of $f_2(x)$ is 1 and its principal part is $p(x) = \frac{1}{x}$.

$$f_3(x) = \log \left(1 + \sin \frac{2}{x} \right)$$

$$f_3(x) = \log \left(1 + \sin \frac{2}{x} \right) \sim \sin \frac{2}{x} \sim \frac{2}{x}, \text{ for } x \rightarrow +\infty$$

Thus $f_3(x) \sim \frac{2}{x}$, for $x \rightarrow +\infty$

W.r.t. the infinitesimal test function $\varphi(x) = \frac{1}{x}$, as $x \rightarrow +\infty$, the order of infinitesimal of $f_3(x)$ is 1 and its principal part is $p(x) = 2 \frac{1}{x}$.

$$g(x) = \tan \frac{3}{x^2}$$

$$g(x) = \tan \frac{3}{x^2} \sim \frac{3}{x^2}, \text{ for } x \rightarrow +\infty$$

W.r.t. the infinitesimal test function $\varphi(x) = \frac{1}{x}$, per $x \rightarrow +\infty$, the order of infinitesimal of $g(x)$ is 2 and its principal part is $p(x) = 3 \frac{1}{x^2}$.

Which of them is $o(\frac{1}{x})$ (as $x \rightarrow +\infty$) ?

The functions $f_1(x), f_2(x), f_3(x)$ have the same order as $\frac{1}{x}$, for $x \rightarrow +\infty$.

On the other hand $g(x) = o(\frac{1}{x})$: indeed $\lim_{x \rightarrow +\infty} \frac{g(x)}{\frac{1}{x}} = \lim_{x \rightarrow +\infty} \frac{\frac{3}{x^2}}{\frac{1}{x}} = \lim_{x \rightarrow +\infty} \frac{3}{x} = 0$.

Find the order of infinity and the principal part w.r.t. $\varphi(x) = x$, as $x \rightarrow +\infty$, for the functions:

$$f_1(x) = 2x - \sqrt{4x^2 + x^4}$$

$$f_1(x) = 2x - \sqrt{4x^2 + x^4} \sim 2x - \sqrt{x^4}, \text{ for } x \rightarrow +\infty \sim 2x - x^2, \text{ for } x \rightarrow +\infty \sim -x^2, \text{ for } x \rightarrow +\infty$$

W.r.t. the infinite test function $\varphi(x) = x$, as $x \rightarrow +\infty$, the order of infinity of $f_1(x)$ is 2 and its principal part is $P(x) = -x^2$.

$$f_2(x) = \frac{x^3 + x + 2}{x^2 - x - 1}$$

$$f_2(x) = \frac{x^3 + x + 2}{x^2 - x - 1} = \frac{x^3 \left(1 + \frac{1}{x^2} + \frac{2}{x^3}\right)}{x^2 \left(1 - \frac{1}{x} - \frac{1}{x^2}\right)} = \frac{x^3}{x^2}, \text{ as } x \rightarrow +\infty \sim x, \text{ as } x \rightarrow +\infty$$

W.r.t. the infinite test function $\varphi(x) = x$, as $x \rightarrow +\infty$, the order of infinity of $f_2(x)$ is 1 and its principal part is $P(x) = x$.

$$f_3(x) = \frac{x^2 + 1}{\sqrt{x^2 - 1}}$$

$$f_3(x) = \frac{x^2 + 1}{\sqrt{x^2 - 1}} = \frac{x^2 \left(1 + \frac{1}{x^2}\right)}{\sqrt{x^2 \left(1 - \frac{1}{x^2}\right)}} \sim \frac{x^2}{|x|}, \text{ as } x \rightarrow +\infty \sim x, \text{ as } x \rightarrow +\infty$$

W.r.t. the infinite test function $\varphi(x) = x$, for $x \rightarrow +\infty$, the order of infinity of $f_3(x)$ is 1 and its principal part is $P(x) = x$.

Compute now

$$\lim_{x \rightarrow +\infty} \frac{f_1(x)}{x^2 + \sin x} = \lim_{x \rightarrow +\infty} \frac{-x^2}{x^2 \left(1 + \frac{\sin x}{x^2}\right)} = -1$$

$$\lim_{x \rightarrow +\infty} \frac{x - \sqrt{1 + e^{-x}}}{f_2(x)} = \lim_{x \rightarrow +\infty} \frac{x - 1}{x} = 1$$

$$\lim_{x \rightarrow +\infty} \frac{f_3(x)(2 + \cos x)}{x^2 + x} = \lim_{x \rightarrow +\infty} \frac{x(2 + \cos x)}{x(x + 1)} = \lim_{x \rightarrow +\infty} \frac{(2 + \cos x)}{(x + 1)} = 0$$

10. Let $a_n = \frac{(n+4)! - n!}{(n+2)!}$.

a) Verify that $a_n \sim n^2$, $n \rightarrow +\infty$

$$\begin{aligned} a_n &= \frac{(n+4)! - n!}{(n+2)!} \\ &= \frac{(n+4)(n+3)(n+2)(n+1)n! - n!}{(n+2)(n+1)n!} \\ &= \frac{(n+4)(n+3)(n+2)(n+1) - 1}{(n+2)(n+1)} \sim \frac{n^4}{n^2}, \text{ as } n \rightarrow +\infty \end{aligned}$$

Thus $a_n \sim n^2$, as $n \rightarrow +\infty$.

b) Compute $\lim_{n \rightarrow +\infty} \frac{a_n}{\left(1 + \frac{1}{n}\right)^{n^2}}$

$$\lim_{n \rightarrow \infty} \frac{a_n}{\left(1 + \frac{1}{n}\right)^{n^2}} = \lim_{n \rightarrow \infty} \frac{n^2}{\left(\left(1 + \frac{1}{n}\right)^n\right)^n} = \lim_{n \rightarrow \infty} \frac{n^2}{e^n} = 0$$

11. Compute the principal part as $x \rightarrow +\infty$ and as $x \rightarrow 0$ (w.r.t. the standard test functions) for $f(x) = \frac{5x^2 + 2x}{3}$.

Given the previous computations, solve

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x^3 \sin\left(\frac{1}{x}\right)} \quad ; \quad \lim_{x \rightarrow 0} f(x) \frac{\sin^2 x}{1 - \cos x}$$

It holds:

$$\frac{5x^2 + 2x}{3} \sim \frac{5}{3}x^2, \quad \text{as } x \rightarrow +\infty, \quad \frac{5x^2 + 2x}{3} \sim \frac{2}{3}x, \quad \text{as } x \rightarrow 0$$

Hence the principal part of $f(x)$ as $x \rightarrow +\infty$ is $P(x) = \frac{5}{3}x^2$, while the principal part of $f(x)$ as $x \rightarrow 0$ is the function $p(x) = \frac{2}{3}x$.

Using these results, compute the limits:

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x^3 \sin\left(\frac{1}{x}\right)} = \lim_{x \rightarrow +\infty} \frac{\frac{5}{3}x^2}{x^3 \frac{1}{x}} = \lim_{x \rightarrow +\infty} \frac{\frac{5}{3}x^2}{x^2} = \frac{5}{3}$$

$$\lim_{x \rightarrow 0} f(x) \frac{\sin^2 x}{1 - \cos x} = \lim_{x \rightarrow 0} \frac{2}{3}x \frac{x^2}{\frac{1}{2}x^2} = 0$$

SOLUTIONS - EXERCISES from WRITTEN EXAMS

1. (19 September 2017)

- (a) Suppose that a function assumes value with opposite sign in the two points a and b . State which are the conditions to have 0 in the range of the function. If the function is continuous on the interval $I = [a, b]$, by the Zeros Existence Theorem, f necessarily takes value 0 in I .
- (b) Consider the function $g(x) = \text{sgn}((x-1)(x+3))$, where

$$\text{sgn}(t) = \begin{cases} 1 & \text{if } t > 0 \\ 0 & \text{if } t = 0 \\ -1 & \text{if } t < 0 \end{cases}$$

Say if the previous function annihilates in the interval $[0, 4]$ and say if, in such interval, it satisfies conditions given at point (a).

Study the sign of $(x-1)(x+3)$, then

$$\begin{cases} (x-1)(x+3) > 0 & \text{if } x < -3 \vee x > 1 \\ (x-1)(x+3) = 0 & \text{if } x = -3 \vee x = 1 \\ (x-1)(x+3) < 0 & \text{if } -3 < x < 1 \end{cases}$$

hence:

$$\text{sgn}((x-1)(x+3)) = \begin{cases} 1 & \text{if } x < -3 \vee x > 1 \\ 0 & \text{if } x = -3 \vee x = 1 \\ -1 & \text{if } -3 < x < 1 \end{cases}$$

The function is zero in $x = 1$, the belongs to the interval $[0, 4]$.

Since $g(x)$ is not continuous in $[0, 4]$, we cannot apply the Zeros Existence Theorem; nevertheless, we have that the function annihilates: indeed continuity is a sufficient condition, but not necessary for the existence of zeros.

2. (1 February 2017 - I)

- (a) Write the definition for

$$f(x) = o(2x^2 - 1) \quad \text{as } x \rightarrow +\infty,$$

and show the latter with an example.

The definition is: $\lim_{x \rightarrow +\infty} \frac{f(x)}{2x^2 - 1} = 0$.

Consider the examples

Example 1:

$$f(x) = x.$$

In this case, both the functions $f(x) = x$ and $g(x) = 2x^2 - 1$ are infinite (as $x \rightarrow +\infty$) and **it holds** $f(x) = o(2x^2 - 1)$.

Example 2:

$$f(x) \equiv 1.$$

In this case $g(x) = 2x^2 - 1$ is infinite for $x \rightarrow +\infty$, whereas $f(x)$ is not and **it holds** $f(x) = o(2x^2 - 1)$.

- (b) Establish if, from the hypothesis $f(x) = o(2x^2 - 1)$ for $x \rightarrow +\infty$, it follows that

$$e^{f(x)} = o(e^{2x^2 - 1}) \quad \text{as } x \rightarrow +\infty$$

If it is true, prove it, otherwise provide a counterexample.

Let

$$g(x) = 2x^2 - 1.$$

Note that

$$f(x) - g(x) = g(x) \left(\frac{f(x)}{g(x)} - 1 \right).$$

Given $f = o(g)$ we have that $f/g \rightarrow 0$ and thus

$$\left(\frac{f(x)}{g(x)} - 1 \right) \rightarrow -1.$$

Given that $\lim_{x \rightarrow +\infty} g(x) = +\infty$ it holds

$$\lim_{x \rightarrow +\infty} g(x) \left(\frac{f(x)}{g(x)} - 1 \right) = -\infty \text{ and therefore } \lim_{x \rightarrow +\infty} \frac{e^{f(x)}}{e^{g(x)}} = \lim_{x \rightarrow +\infty} e^{g(x) \left(\frac{f(x)}{g(x)} - 1 \right)} = 0.$$

Then, $e^{f(x)} = o(e^{g(x)})$ for $x \rightarrow +\infty$.

3. (28 January 2016 - II)

- (a) Write the definition of order of infinitesimal of $f(x)$ with respect to $\frac{1}{x}$, for $x \rightarrow +\infty$.

See the textbook.

- (b) Prove that

$$a_n = \sqrt[7]{4n - \frac{2}{\sqrt{n} \log^3 n}} - \sqrt[7]{4n} = o\left(\frac{1}{n}\right), \quad \text{for } n \rightarrow +\infty.$$

It holds

$$\begin{aligned} \sqrt[7]{4n} \left[\left(1 - \frac{1}{2n^{3/2} \log^3 n} \right)^{1/7} - 1 \right] &= \sqrt[7]{4n} \left[-\frac{1}{14n^{3/2} \log^3 n} + o\left(\frac{1}{n^{3/2} \log^3 n}\right) \right] \\ &= -\frac{\sqrt[7]{4}}{14 n^{19/14} \log^3 n} + o\left(\frac{1}{n^{19/14} \log^3 n}\right) \quad \text{for } n \rightarrow +\infty. \end{aligned}$$

Therefore

$$\frac{a_n}{\frac{1}{n}} = a_n n = -\frac{\sqrt[7]{4}}{14 n^{5/14} \log^3 n} + o\left(\frac{1}{n^{5/14} \log^3 n}\right) = o(1) \quad \text{as } n \rightarrow +\infty.$$

- (c) Say if the following statement is true or false. If it is true, prove it; otherwise provide a counterexample. If a_n is such that $1 \leq a_n \leq 3$, for every $n \in \mathbb{N}$, then $\lim_{n \rightarrow +\infty} a_n$ exists and it is a real number $l \in [1, 3]$.

The statement is false. Take the following counterexample: $a_n = (-1)^n + 2$ such that $1 \leq a_n \leq 3$ but it does not admit limit as $n \rightarrow +\infty$.