

Galaxy Zoo: Evidence for rapid, recent quenching across a population of AGN host galaxies

R. J. Smethurst,¹ C. J. Lintott,¹ B. D. Simmons,¹ K. Schawinski,² S. Bamford,³
C. Cardamone,⁴ S. J. Kruk,¹ K. Masters,⁵ M. Urry,⁶ K. W. Willett,⁷ O. I. Wong⁸

¹ *Oxford Astrophysics, Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford, OX1 3RH, UK*

² *Institute for Astronomy, Department of Physics, ETH Zurich, Wolfgang-Pauli Strasse 27, CH-8093 Zurich, Switzerland*

³ *School of Physics and Astronomy, The University of Nottingham, University Park, Nottingham, NG7 2RD, UK*

⁴ *Math & Science Department, Wheelock College, 200 The Riverway, Boston, MA 02215, USA*

⁵ *Institute of Cosmology and Gravitation, University of Portsmouth, Dennis Sciama Building, Barnaby Road, Portsmouth, PO1 3FX, UK*

⁶ *Department of Physics and Yale Center for Astronomy and Astrophysics, Yale University, PO Box 208121, New Haven, CT 06520-8121, USA*

⁷ *School of Physics and Astronomy, University of Minnesota, 116 Church St SE, Minneapolis, MN 55455, USA*

⁸ *International Centre for Radio Astronomy Research, UWA, 35 Stirling Highway, Crawley, WA 6009, Australia*

20 July 2015

ABSTRACT

We present a population study of the star formation history (SFH) of 1,244 Type 2 AGN host galaxies, compared to 123,243 inactive galaxies using a Bayesian method. We find evidence for the Type 2 AGN host galaxies to have undergone a recent and rapid drop in their star formation rate. This result is not seen for the inactive galaxies whose SFHs are dominated by the effects of downsizing at earlier epochs, a secondary effect for the AGN host galaxies. We show that rapid quenching histories cannot account fully for the quenching of all the star formation across the population of AGN host galaxies and that slower quenching histories, attributed to non-violent processes of evolution are also key in their evolution. This is contrary to the merger driven scenario of the co-evolution of black holes and their galaxies. The availability of gas in the reservoirs of a galaxy, and its ability to be replenished, appears to be the key driver behind this co-evolution.*

1 INTRODUCTION

The nature of the observed co-evolution of galaxies and their central supermassive black holes (Magorrian et al. 1998; Marconi & Hunt 2003; Haring & Rix 2004) and the effects of AGN feedback on galaxies are two of the most important open issues in galaxy evolution. AGN feedback was first suggested as a mechanism for regulating star formation in simulations (Croton et al. 2006; Bower et al. 2006; Somerville et al. 2008) and some indirect evidence has been observed for both positive and negative feedback in various systems (see the comprehensive review from Fabian 2006).

The strongest observational evidence for AGN feedback is that the largest fraction of AGN are found in the green valley (Cowie & Barger 2008; Hickox et al. 2009; Schawinski et al. 2010), suggesting some link to the process of quenching which moves a galaxy from the blue cloud to the red sequence. However, concrete statistical evidence for the effect of AGN feedback on the host galaxy population has yet to be found.

Here we present a large observational population study of the quenching of the host galaxies of Type 2 AGN, using a new PYTHON routine, STARPYPY¹, which implements a Bayesian method to effectively model the SFH of a galaxy with two parameters, time of quenching and exponential rate, given the observed near ultra-violet (NUV) and optical colours. We aim to determine the following: (i) Are galaxies currently hosting an AGN undergoing quenching? (ii) If so, when and at what rate does this quenching occur?

The zero points of all magnitudes are in the AB system and where necessary, we adopt the WMAP Seven-Year Cosmology (Jarosik et al. 2011) with $(\Omega_m, \Omega_\lambda, h) = (0.26, 0.73, 0.71)$.

2 DATA & METHODS

In this investigation we use visual classifications of galaxy morphologies from the Galaxy Zoo 2² citizen science project (Willett et al. 2013), which obtains multiple independent

* This investigation has been made possible by the participation of more than 250,000 users in the Galaxy Zoo project. Their contributions are individually acknowledged at <http://authors.galaxyzoo.org>

¹ <http://github.com/zoouniverse/starpy>

² <http://zoo2.galaxyzoo.org/>




Figure 1. Randomly selected SDSS *gri* composite images from the sample of 1,244 Type 2 AGN showing the continuous probabilistic nature of the Galaxy Zoo sample from a redshift range $0.04 < z < 0.05$. The debiased ‘disc or featured’ vote fraction (see Willett et al. 2013) for each galaxy is shown. The scale for each image is 0.099 arcsec/pixel.

classifications for each optical galaxy image. The full question tree for each image is shown in Figure 1 of Willett et al. 2013. The Galaxy Zoo 2 (GZ2) project used 304,022 images from the SDSS DR8 (York et al. 2000; Aihara et al. 2011) all classified by *at least* 17 independent users, with a mean number of classifications of ~ 42 .

Further to this, we required NUV photometry from the GALEX survey, within which $\sim 42\%$ of the GZ2 sample was observed, giving 126,316 galaxies total ($0.01 < z < 0.25$). This will be referred to as the GZ2-GALEX sample. The completeness of this sample ($-22 < M_u < -15$) is shown in Figure 2 of Smethurst et al. (2015) with the *u*-band absolute magnitude against redshift for this sample in comparison to the SDSS data set.

2.1 STARPY

STARPY³ is a PYTHON code which allows the user to derive the quenched star formation history (SFH) of a galaxy through a Bayesian Markov Chain Monte Carlo method with the input of two observed photometric colours, a redshift, and the use of the SSP models of Bruzual & Charlot (2003). The star formation history template is an exponential decline of the SFR and is described by two parameters $[t_q, \tau]$, where t_q is the time at which the onset of quenching begins [*Gyr*] and τ is the exponential rate at which quenching occurs [*Gyr*]. Under the simplifying assumption that all galaxies formed at $t = 0$ Gyr with an initial burst of star formation, the SFH can be described as:

$$SFR = \begin{cases} i_{sfr}(t_q) & \text{if } t < t_q \\ i_{sfr}(t_q) \times \exp\left(\frac{-(t-t_q)}{\tau}\right) & \text{if } t > t_q \end{cases} \quad (1)$$

where i_{sfr} is an initial constant star formation rate dependent on t_q (Schawinski et al. 2014; Smethurst et al. 2015). A smaller τ value corresponds to a rapid quench, whereas a larger τ value corresponds to a slower quench. The output of STARPY is probabilistic in nature and provides the posterior probability distribution across the entirety of the two parameter space for each individual galaxy. The probabilistic fitting methods to this SFH for an observed galaxy are described in full detail in Smethurst et al. (2015) wherein the STARPY code was run on the GZ2-GALEX sample using the observed $u - r$ and $NUV - u$ colours from SDSS and GALEX photometry.

2.2 AGN Sample

We select Type 2 AGN using a BPT diagram (Baldwin, Phillips & Terlevich 1981) using line and continuum strengths for [OIII], [NII], [SII] and [OII] obtained from the MPA-JHU catalogue (Kauffman et al. 2003a; Brinchmann et al. 2004) which matched those in the GZ2-GALEX; we then required the $S/N > 3$ for each emission line as in Schawinski et al. (2010). Those galaxies which satisfy all of the inequalities defined in Kewley et al. (2001), Kauffman et al. (2003b) and Kewley et al. (2006) are selected as Type 2 AGN, giving 1,244 host galaxies ($\sim 9\%$ of the GZ2-GALEX sample). Sarzi et al. (2010); Yan & Blanton (2012) and Singh et al. (2013) have all demonstrated that LINERs are not powered by AGN, therefore for purity, we exclude these galaxies from the sample (55 galaxies in total) with no change to the results. This will be referred to as the AGN-HOST sample.

Images from SDSS for ten randomly selected Type 2 AGN host galaxies from this sample are shown in Figure 1 and the entire sample is shown in Figure 2 with those galaxies selected as Type 2 AGN marked in red and the rest of the GZ2-GALEX in black. For this sample the mean $\log L[OIII] \sim 45.0$ with a range of $\log L[OIII]$ luminosities of $42.9 - 46.7$.

We construct a sample of inactive galaxies by removing from the GZ2-GALEX sample all galaxies in the agn-host sample, as well as sources identified as Type 1 AGN by the presence of broad emission lines (?). We refer to this sample below as the INACTIVE sample.

Since this investigation is focussed on whether an AGN can have an impact on the SF of their host galaxy, we must consider whether there is a selection effect present in this identification method. The possibility that an AGN in a host galaxy in the blue cloud could be obscured by the emission from star formation was addressed by Schawinski et al. (2010). They showed, through a simple empirical experiment which simulated the addition of an AGN of certain luminosity to a star forming galaxy, that this method produces a complete sample, even in the blue cloud, at luminosities of $L[OIII] > 10^{40} \text{ erg s}^{-1}$. We can therefore assume that this method is producing a complete sample of AGN, fully sampling the range of star formation rates of their host galaxies.

Type 2 AGN were used in this analysis as opposed to Type 1 due to their photometric obscuration. Type 1 AGN contaminate their galaxy’s photometric measurements which would need to be removed through aperture matching. Due to the requirement for NUV colours from GALEX, in order to be sensitive to any recent star formation, the aperture matching to SDSS becomes a non-trivial task.

Simmons et al. (2011) showed that the obscuration of a Type 2 AGN is more efficient in the NUV than in the optical. Residual NUV light from the AGN can be neglected

³ <http://github.com/zoouniverse/starpy>




Figure 2. BPT diagrams for galaxies in the GZ2-GALEX sample (black crosses) with $S/N > 3$ for each emission line. Inequalities defined in: Kewley et al. (2001) to separate star forming galaxies from AGN (dashed lines), Kauffman et al. (2003b) to separate star forming from composite SF-AGN galaxies (solid line) and Kewley et al. (2006) to separate LINERS and Seyferts (dotted lines). Galaxies were included in the AGN-HOST sample (red circles) if they satisfied all the inequalities from Kewley et al. (2001) and Kewley et al. (2006).

Table 1. Table showing the number of galaxies in each of the three mass bins for both the AGN-HOSTS and INACTIVE galaxy samples.

Mass Bin	AGN-HOSTS	INACTIVE
$\log M_* < 10.25 M_\odot$	165 (13.3%)	41197 (33.4%)
$10.25 < \log M_* [M_\odot] < 10.75$	630 (50.6%)	46428 (37.7%)
$\log M_* > 10.75 M_\odot$	449 (36.1%)	35618 (28.9%)

in comparison to that of the galaxy, however there is often some residual optical light that can affect the measurements of the host galaxy’s photometry. We can easily subtract this central optical AGN flux using the PSF magnitudes provided by SDSS, however the change in the colours of these galaxies after this correction is negligible, with $\Delta(u-r) \sim 0.09$. In order to reduce the amount of uncertainty that will progress into the SFH population likelihoods, we will not use these corrected optical colours in favour of reducing unnecessary complexity. We also split both the AGN-HOST and INACTIVE samples into low, medium and high masses (see Table 1) to control for any degeneracies. Masses were calculated using the $u-r$ colour and absolute r-band magnitude with the method outlined in Baldry et al. (2006).

3 RESULTS

We apply the method outlined in Section 2.1 to obtain a 2-dimensional likelihood distribution across the $[t_q, \tau]$ parameter space. We combine individual galaxy likelihood distributions within the AGN-HOST and INACTIVE galaxy samples, additionally weighting by GZ2 morphologies to produce quenching parameter likelihood distributions for both disc- and smooth-dominated galaxy populations in the 3 mass bins defined in Table 1. We stress that this portion is purely for visualisation purposes and is no longer a Bayesian method.

These 2-dimensional likelihoods are summed across each parameter axis and normalised to produce the one dimensional histograms shown in Figures 3 and 4 for the quenching time, t_q and exponential quenching rate, τ , respectively. In each figure the summed 1-dimensional nor-

malised probability distribution across the given parameter is shown for smooth and disc dominated galaxies across three mass bins for the AGN-HOST and INACTIVE samples. In Figure 4 the percentage likelihood in each region of quenching rate, shown by the dashed lines for rapid ($\tau < 1$ Gyr), intermediate ($1 < \tau$ [Gyr] < 2) and slow ($\tau > 2$ Gyr) quenching timescales are shown.

It is immediately apparent from Figures 3 and 4 that there is a distinct difference between the distribution of likelihood for AGN-HOST and INACTIVE populations for both parameters. For the INACTIVE sample, the likelihood for quenching at later times decreases with increasing mass, until for the lower mass galaxies the quenching is roughly constant with time for both smooth and disc dominated populations (right panel Figure 3). This is observational evidence of downsizing across the generic galaxy population whereby stars in massive galaxies form first and quench early (Cowie et al. 1996; Thomas et al. 2010).

The distribution of likelihood for the AGN-HOST population across the quenching time t_q parameter (left panels of Figure 3), is obviously different from that of the inactive galaxies (right panels of Figure 3). Recent quenching is the dominant history across all three mass bins, particularly for low mass galaxies. However, this effect is dampened in higher mass galaxies where quenching at earlier times also has significant likelihood.

The distribution of likelihoods for the rate of quenching, τ , in Figure 4, show the likelihood for rapid quenching ($\tau < 1$ Gyr) decreases for the INACTIVE population of increasing mass (see percentage likelihoods in the right hand panels of Figure 4) and the likelihood for slow quenching ($\tau > 2$ Gyr) increases for the INACTIVE disc population with increasing mass.

These trends in the distribution of likelihood of the rate of quenching are also seen for the AGN-HOST population; the likelihood for rapid quenching decreases with increasing mass for both smooth and disc dominated galaxies, but the overall distribution of the likelihood is very different. The likelihood distribution for the AGN-HOST galaxies shows evidence for the dominance of rapid, recent quenching having occurred across the entire population.




Figure 3. Likelihood distribution for the quenching time, t_q parameter normalised so that the areas under the curves are equal. AGN-HOST (left) and INACTIVE (right) galaxies are split into low (top), medium (middle) and high (bottom) mass for smooth (red dashed) and disc (blue solid) galaxies. A low value of t_q corresponds to the early Universe and a high value to the recent Universe.

Interestingly, there is little difference between the two morphological types of AGN-HOST galaxies for either quenching parameter.

4 DISCUSSION

The differences between the distribution of likelihood for the AGN-HOST and INACTIVE galaxy populations reveals that AGN have a significant effect on the SFH of their host galaxy and can be associated with both recent rapid quenching and earlier, slower quenching histories.

As well as the evidence for downsizing in the INACTIVE population (Figure 3), we can also see its effect on the AGN-HOST population. Although recent quenching is the dominant history, quenching at earlier times has significant likelihood, increasing with mass. These galaxies may have been affected by downsizing earlier in life with the current AGN also having an effect on the SFR through feedback, causing a recent, rapid quench of any residual star formation.

Tortora et al. (2009) model the effects of negative AGN feedback on a typical early type (i.e. smooth) galaxy and find the time between the current galaxy age, t_{gal} and the time that the feedback began t_{AGN} , peaks at $t_{gal} - t_{AGN} \sim 0.85$ Gyr. This agrees with the location of the peak in Figure 3 for low mass galaxies, where the difference between the peak of the likelihood and the average age of the population (calculated from the redshift and assuming all galaxies form at $t = 0$) is ~ 0.83 Gyr. This suggests that this dominant recent quenching history is caused directly by AGN feedback, as opposed to the AGN being a consequence of an alternative quenching mechanism. Smethurst et al. (2015) showed quenching is highly dependant on the morphology of the galaxy; since the AGN-HOST population does not show any morphological dependencies, this also suggests that this quenching is caused by AGN feedback.

If a slowly ‘dying’ or ‘dead’ galaxy has an infall of gas




Figure 4. Likelihood distribution for the quenching rate, τ normalised so that the areas under the curves are equal. AGN-HOST (left) host and INACTIVE (right) galaxies are split into low (top), medium (middle) and high (bottom) mass for smooth (red dashed) and disc (blue solid) galaxies. The black dashed lines show the separation between rapid ($\tau < 1.0$ Gyr), intermediate ($1.0 < \tau$ [Gyr] < 2.0) and slow ($\tau > 2.0$ Gyr) quenching timescales with the percentage likelihood distribution in each region for disc (blue) and smooth (red) populations. A small (large) value of τ corresponds to a rapid (slow) quench.

either through a minor merger, galaxy interaction or environmental change, this can trigger further star formation and feed the central black hole, igniting an AGN (Kaviraj et al. 2014). In turn this AGN can then quench the recent boost in star formation. This track is similar to the evolution history theorised for blue ellipticals (Kaviraj et al. 2013; McIntosh et al. 2014; Haines et al. 2015). As for the disc galaxies, this preference for slower evolution timescales follows from the ideas of previously isolated discs evolving slowly by the Kennicutt-Schmidt (KS; Schmidt 1959; Kennicutt 1997) law which can then undergo an interaction or merger to reinvigorate star formation and feed the central black hole.

The difference between the AGN-HOST and INACTIVE population distribution of likelihood in Figure 4 for the rate of quenching, τ , tells a story of gas reservoirs. Smethurst et al. (2015) speculate that rapid quenching rates could be attributed to mergers of galaxies, therefore we expect the trend of increasing likelihood for rapid quenching rates with increasing mass for the INACTIVE population, as mergers are thought to be responsible for creating the most massive smooth galaxies (Conselice et al. 2003; Springel, Di Matteo & Hernquist 2005; Hopkins et al. 2008).

The trends with likelihood of τ in Figure 4 are reversed however for the AGN-HOST population. The likelihood for rapid quenching decreases with increasing mass for both smooth and disc dominated populations. There is also more dominance of slower quenching timescales in the higher mass population across both morphologies. It appears that the most massive disc galaxies, with the most massive gas reservoirs, evolve with a slow quench of star formation through the KS law and also have enough gas to feed the central black hole to trigger a current AGN (Varela et al. 2004; Emsellem et al. 2015). This challenges the typical merger

driven theory for the co-evolution of black holes and their host galaxies.

Conversely, a rapid quench, possibly caused by the AGN itself through negative feedback, is the most dominant history for low mass galaxies with lower gravitational potentials which allow gas to be expelled more easily (or heated by the AGN) across the entire galaxy (Tortora et al. 2009). This rapid quenching is still apparent but at a lower likelihood across the smooth and disc populations in higher mass galaxies, which have larger potentials, making it more difficult for the AGN to have an impact on the galaxy wide SFR (Ishibashi et al. 2012; Zinn et al. 2013).

For the medium mass AGN-HOST population we see a bimodal distribution of likelihood between these two dominant quenching histories, highlighting the strength of this method that it is capable of detecting such variation in the SFHs across a population of galaxies.

We have used morphological classifications from the Galaxy Zoo 2 project to determine the morphology-dependent star formation histories of a population of 1,244 Type 2 Seyfert AGN host galaxies in comparison to a ‘typical’ inactive galaxy population via a Bayesian analysis of an exponentially declining star formation history model. We determined the most likely parameters for the quenching onset time, t_q and quenching timescale τ and find clear differences in the combined population likelihoods between inactive and AGN host galaxies and for a link between a galaxy currently hosting an AGN and its SFR. There is strong evidence for downsizing in massive inactive galaxies, which appears as a secondary effect in AGN host galaxies with higher masses. Our main finding is the detection of the dominance of rapid, recent quenching across a population of AGN host galaxies.

ACKNOWLEDGEMENTS

RS acknowledges funding from the STFC Grant Code ST/K502236/1. BDS gratefully acknowledges support from Balliol College, Oxford. KS gratefully acknowledges support from Swiss National Science Foundation Grant PP00P2_138979/1. SJK acknowledges funding from the STFC Grant Code ST/MJ0371X/1. KLM acknowledges funding from The Leverhulme Trust as a 2010 Early Career Fellow. KWW acknowledges funding from a Grant-in-Aid from the University of Minnesota. OIW acknowledges a Super Science Fellowship from the Australian Research Council. The development of Galaxy Zoo was supported in part by the Alfred P. Sloan Foundation and The Leverhulme Trust. Based on observations made with the NASA Galaxy Evolution Explorer⁴ and the SDSS⁵.

REFERENCES

- Aihara, H. et al., 2011, *ApJSS*, 193, 29
 Baldry, I. et al., 2006, *MNRAS*, 373, 469

- Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, *PASP*, 93, 5
 Bamford, S. et al., 2009, *MNRAS*, 393, 1324
 Bower, R. et al., 2006, *MNRAS*, 370, 645
 Brinchmann, J. et al., 2004, *MNRAS*, 351, 1151
 Bruzual, G. & Charlot, S., 2003, *MNRAS*, 344, 1000
 Cisternas, M. et al., 2011, *ApJ*, 726, 57
 Conselice, C. J. et al., 2003, *AJ*, 126, 1183
 Cowie, L. et al., 1996, *AJ*, 112, 839
 Cowie, L. & Barger, A. J., 2008, *ApJ*, 686, 72
 Croton, D. J. et al., 2006, *MNRAS*, 365, 11
 Emsellem, E. et al. 2015, *MNRAS*, 446, 2468
 Fabian, A. C. 2006, *ARA&A*, 50, 455
 Haines, T. et al., 2015, *arXiv:1505.01493*
 Haring, N. & Rix, H-W., 2004, *ApJ*, 604, 89
 Hickox, R. C., et al., 2009, *ApJ*, 696, 891
 Hopkins, F. et al., 2008, *ApJSS*, 175, 390
 Ishibashi, W. et al., 2012, *MNRAS*, 427, 2998
 Kauffman, G. et al., 2003, *MNRAS*, 341, 33
 Kauffman, G. et al., 2003, *MNRAS*, 346, 1055
 Kaviraj, S. et al., 2013, *MNRAS*, 428, 925
 Kaviraj, S. et al., 2014, *MNRAS*, 440, 2944
 Kennicutt, R. C., 1997, *ApJ*, 498, 491
 Kewley, L. J. et al., 2001, *ApJ*, 556, 121
 Kewley, L. J. et al., 2006, *MNRAS*, 372, 961
 Kormendy, J. & Kennicutt, R. J., 2004, *ARA&A*, 42, 603
 Lintott, C. J. et al., 2011, *MNRAS*, 410, 166
 Magorrian, J. et al., 1998, *AJ*, 115, 2285
 Marconi, A. & Hunt, L. K., 2003, *ApJ*, 589, 21
 McIntosh, D. et al., 2014, *MNRAS*, 442, 533
 Oh, K. et al. 2015, *arXiv: 1504.07247*
 Yan, R. & Blanton, M. R. 2012, *ApJ*, 747, 61
 Sarzi, M. et al., 2010, *MNRAS*, 402, 2187
 Schawinski, K. et al., 2010, *MNRAS*, 711, 284
 Schawinski, K. et al., 2014, *MNRAS*, 440, 889
 Schmidt, M., 1959, *ApJ*, 129, 243
 Simmons, B. D. et al., 2011, *ApJ*, 734, 121
 Singh, R. et al., 2013, *A&A*, 558, 43
 Smethurst, R. J. et al., 2015, *MNRAS*, 450, 435
 Somerville, R. S. et al., 2008, *MNRAS*, 391, 481
 Springel, V., Di Matteo, T. & Hernquist, L., 2005, *ApJ*, 620, L79
 Taylor, M. B., 2005, *ASP Conference Series*, 347
 Thomas, D. et al., 2010, *MNRAS*, 404, 1775
 Tortora, C. et al., 2009, *MNRAS*, 369, 61
 Varela, J. et al., 2004, *A&A*, 420, 873
 Willett, K. et al., 2013, *MNRAS*, 435, 2835
 York, D. G. et al., 2000, *AJ*, 120, 1579
 Zinn, P. et al., 2013, *ApJ*, 774, 66

⁴ <http://galex.stsci.edu/GR6/>

⁵ <https://www.sdss3.org/collaboration/boiler-plate.php>