#### What the p-Addicts Did This Week

Alexandra Michel, Joseph Rennie, Andrew Miller

©2014 the p-Addicts. All rights reserved.

July 9th, 2014


#### Outline

Birth of the p-Addicts

Adventures of the p-Addicts


What does the future have in store for the *p*-Addicts?

## Eric's Paper about $\{3^n\}$




#### Example

This is a picture of the sequence  $\{3^n\}$  in base 2, where the nth row is the binary digits (in black and white) of  $\{3^n\}$ .




### Example

Now we have the sequence  $\{3^{2^n}\}$  in base 2, where the *n*th row is the binary digits of  $\{3^{2^n}\}$ .



#### Example

This is the same sequence, but sheared so that the white space is gone and the columns line up. Using p-adic power series, Eric was able to prove that the limit of this sequence was  $\log 3$ .




#### Definition of Convergence


Remember that the definition of convergence is that for every k > 0 there is a N such that for n, m > N we have:

$$|C(2^n) - C(2^m)|_2 \le 2^{-k}$$

# Catalan(n)



## Catalan(n)



Conjecture: None

### Automata




#### Automata



Conjecture:  $\{\nu_2(C(2^n))\}\$  is constant.

#### Automata



Conjecture:  $\{\nu_2(C(2^n)-2)\}\$  is constant.

We used Mathematica to investigate the sequence  $\{C(2^n)\}$ . It seems to converge 2-adically. Below are the first 25 terms of the sequence in reversed base-2 representation:



We used Mathematica to investigate the sequence  $\{C(2^n)\}$ . It seems to converge 2-adically. Below are the first 25 terms of the sequence in reversed base-2 representation:



Unfortunately, we couldn't collect enough data to make a conjecture about *what* the limit of the sequence is.

We were, however, able to prove that the sequence converges.

We were, however, able to prove that the sequence converges. To do so, we used the following equivalent condition to p-adic convergence:

$$|C(p^n) - C(p^m)|_p \le p^{-k} \text{ iff}$$

$$\nu_p(C(p^n) - C(p^m)) \ge k \text{ iff}$$

$$C(p^n) - C(p^m) \equiv 0 \pmod{p^k} \text{ iff}$$

$$C(p^n) \equiv C(p^m) \pmod{p^k}.$$

We were, however, able to prove that the sequence converges. To do so, we used the following equivalent condition to p-adic convergence:

$$|C(p^n) - C(p^m)|_p \le p^{-k} \text{ iff}$$

$$\nu_p(C(p^n) - C(p^m)) \ge k \text{ iff}$$

$$C(p^n) - C(p^m) \equiv 0 \pmod{p^k} \text{ iff}$$

$$C(p^n) \equiv C(p^m) \pmod{p^k}.$$

And we used a fact from an existing paper: that for all n > k - 1,  $C(2^n) \equiv C(2^{k-1}) \pmod{2^k}$ .

### Convergence of other sequences of Catalan numbers

We also showed that the sequences  $\{C(2^n-1)\}$  converges.

### Convergence of other sequences of Catalan numbers

We also showed that the sequences  $\{C(2^n-1)\}$  converges. Here we used the fact that for any k, the sequence  $\{C(2^n-1)\}$  is constant modulo  $2^k$  for sufficiently large n.

### Convergence of other sequences of Catalan numbers

We also showed that the sequences  $\{C(2^n - 1)\}$  converges. Here we used the fact that for any k, the sequence  $\{C(2^n - 1)\}$  is constant modulo  $2^k$  for sufficiently large n.

And we have a poorly-understood proof that  $\{C(p^n)\}$  converges, using an existing result that has to do with the number of terms in the numerator of  $\binom{2p^n}{p^n}$  that are congruent to the elements of the multiplicative group  $\mathbb{Z}_{p^k}^{\times}$  modulo  $p^k$ . Don't ask.

What does the future have in store for the *p*-Addicts?

We want to find the actual value of

$$\lim_{n\to\infty}C(2^n),$$

### What does the future have in store for the *p*-Addicts?

We want to find the actual value of

$$\lim_{n\to\infty} C(2^n),$$

and we need to keep looking at  $\{C(p^n)\}$ , because our current "proof" just doesn't cut the mustard!

What does the future have in store for the *p*-Addicts?

We want to find the actual value of

$$\lim_{n\to\infty} C(2^n),$$

and we need to keep looking at  $\{C(p^n)\}\$ , because our current "proof" just doesn't cut the mustard!

The methods we used to prove convergences this week were all number-theoretic. We want to find some useful analytic methods as well.

