

Coma & Meningitis

Learning Objectives

- Definitions & terminology
- Etiologies
- Evaluation
- Initial management


What is coma?

- Greek (koma) → deep sleep
- Definition → unarousable unresponsiveness + eyes closed

States of Arousal

- Alert → aware of self and surroundings
 - Avoid “Confusion,” “Lethargy,” “Stupor”
 - Be descriptive
 - **Coma → cannot be aroused by external stimuli or inner need**
- Acute alteration in arousal → emergency

Anatomy of Arousal



Coma → dysfunction of:

- Upper brainstem
- B/L thalami
- B/L cerebral hemispheres


Chapter: Impaired Consciousness and Coma

Author(s): S. Arthur Moore and Eelco F. M. Wijdicks

From: Mayo Clinic Neurology Board Review Clinical Neurology for Initial Certification and MOC: Clinical Neurology for Initial Certification and MOC

Etiologies of Coma

- **Non-Structural:
Toxic/Metabolic/Infectious**
 - >50%
 - symmetric presentation
 - **Structural**
 - cerebral lesion
 - asymmetric (focality)
 - **Psychiatric**
 - Diagnosis of exclusion
- History/exam (& repeat exams) → differentiate between causes
- Multifactorial






Non-Structural



- Toxic
 - Sedatives
 - Anticholinergics
 - Opiates, Barbituates
 - ETOH
 - Poisoning
 - Lead, CO, Methanol
- Infections
 - Meningoencephalitis
 - Sepsis
 - UTI
 - PNA
- Metabolic
 - \uparrow or \downarrow Temp
 - Hypoxia
 - Hypercapnia
 - \uparrow or \downarrow Glucose, Na, Ca
 - \uparrow Mg
 - Uremia, \uparrow Ammonia
 - Hypothyroidism



Structural



- Asymmetric Presentation
 - Unilateral hemispheric mass lesion + herniation
 - Brainstem lesion
 - Ischemic stroke, ICH/SDH
- Symmetric Presentation
 - SAH***
 - Hydrocephalus***
 - Cerebral venous thrombosis***
 - Basilar occlusion
 - Midline brainstem lesion

Approach to Coma

- ABCs + Glucose
- History
 - Collateral
 - Time course of LOC? (abrupt, gradual, fluctuating)
 - Focal symptoms/signs precede LOC?
 - Prior neurologic hx? Recent illness? PMH?
 - Med list? Toxic habits?

Exam

- General Exam
 - Vitals
 - Ventilation pattern
 - Neck stiffness
 - HEENT, skin/mucosal, CV, pulm, abd
- Neuro
 - LOC
 - Brainstem reflexes
 - Sensorimotor response
 - Reflexes



Level of consciousness

- Describe spontaneous behavior + responses to stimuli

- Stimulation

1. Loud vocal commands
2. Shoulder shaking, trapezius squeeze
3. Noxious stimulation
 - Supraorbital ridge pressure
 - TMJ compression
 - Nailbed pressure







Figure 1: The FOUR scale.

Brainstem Reflexes

- Pupillary light reflex (CN 2,3)
 - Impairment → Dilation
 - Herniation
 - Brainstem Lesion
 - State size & reactivity
 - *L pupil 6 mm and non-reactive, R pupil 5 mm → 3 mm*
 - Sedatives can affect reactivity
 - Pinpoint pupils
 - Opiate overdose
 - Disruption of descending sympathetic outflow (pontine pupils)
- Funduscopic exam
 - Papilledema



- Eye Movements (CN 3,4,6)

- Eye position

- Deviation
 - Dysconjugate
 - Nystagmus

- Oculocephalic reflex

- C-spine cleared?

Oculocephalic (Doll's eyes)

Central



Eyes to left




Head to right

Eyes to right




Head to left


Cold caloric response



A normal tonic response



An asymmetrical response



A negative response

(A) Right-way eyes



(B) Wrong-way eyes



Remember that the frontal eye fields initiate gaze to the opposite side!! So, if the right frontal eye field is damaged, the eyes will be "stuck" towards the right - unable to look to the left!

- Caloric testing of OCR
 - Stronger stimulus
 - Tympanic membrane intact?
 - Syringe with catheter + >50 cc ice water → deviation of both eyes toward stimulated ear

Oculocephalic (Doll's eyes)

Central



Eyes to left



Head to right


Eyes to right



Head to left


Cold caloric response

0°C




A normal tonic response

0°C




An asymmetrical response


0°C




A negative response



- Corneal reflex (CN 5,7)
 - Touch edge of cornea
→ blink






- Gag reflex (CN 9,10)
 - Touch posterior pharyngeal wall → elevation of soft palate + pharyngeal muscle contraction





Sensorimotor Response


- Muscle tone
- No spontaneous movement → noxious stimulation
 - Purposeful (withdrawal away from stimulus)
 - Posturing
 - Reflexive
 - Abnormal movements




Reflexes




(A) Normal plantar response



(B) Extensor plantar response (Babinski sign)





(C)





Diagnostic Approach

- ABCs + Glucose
- History/Exam
- HCT & Labs
 - focality, fever, papilledema → STAT HCT
 - CMP, CBC, Coags, Lactate
 - ABG, Ammonia, Cultures, Tox
 - Acute onset → CTA head
 - Fever → CT and then LP
 - Seizure(s) → EEG

* Life-threatening & treatable causes



- Initial evaluation unrevealing →
 - MRI brain
 - Prolonged EEG monitoring
 - Serum drug levels, TSH/FT4, poisoning



Management

- ABCs
 - Stabilize C-spine
 - GCS<8 → intubation
 - Treat hypotension (MAP<70mm Hg), fever, hypo-/hyperglycemia

* Empiric thiamine (before D50)



- Overdose
 - Narcotic → naloxone
 - Benzo → flumazenil
- Herniation → mannitol
- Seizure → benzodiazepine + phenytoin/fosphenytoin
- CNS infection →
vancomycin/ceftriaxone + acyclovir
 - Blood cx + empiric treatment

Mimics of Coma


- Locked-In Syndrome
 - alert/aware, vertical eye movements & blink preserved
 - quadriplegic, lower cranial nerve palsies
 - horizontal gaze palsy
 - facial diplegia
 - palate/tongue weakness
 - ventral pontine lesion
 - basilar artery thrombosis
 - central pontine myelinolysis
 - mass lesion



Fig. 3. T2-weighted axial MRI showing massive and bilateral pontine infarction.

Mimics of Coma

- Persistent Vegetative State
 - loss of awareness & cognitive function
 - vegetative fxn (cardiorespiratory drive) preserved
- Premotor/Prefrontal Pathology
 - akinetic mutism or abulia
 - impaired motor initiation/planning
- Catatonia, Psychogenic
 - EEG
 - cold calorics



Checklist for the Evaluation of Acute Coma

History

- ◆ Sudden onset? Witnessed onset?
- ◆ Previous level of function
- ◆ Comorbid conditions
- ◆ Medications and toxic exposures
- ◆ Previous episodes of altered consciousness?
- ◆ Location (out of hospital, hospital floor, intensive care unit)

Physical Examination

- ◆ Vital signs
- ◆ Brainstem reflexes
- ◆ Funduscopy
- ◆ Motor responses to pain
- ◆ Lateralizing or other focal findings?
- ◆ Meningeal signs?
- ◆ Adventitious movements?

Blood (and Urine) Tests

- ◆ Glucose, electrolytes, blood urea nitrogen/creatinine, complete blood cell count
- ◆ Consider pH, PaCO₂, liver enzymes, ammonia, ethanol, toxicologic screen (urine and sometimes serum), levels of prescribed drugs, thyroid-stimulating hormone (TSH), cortisol
- ◆ Infectious workup (when pertinent)
- ◆ Autoimmune encephalitis panel (when pertinent)

Brain Imaging?

- ◆ Brain parenchyma (CT versus MRI)
- ◆ Brain vessels (arteries, veins)

Lumbar Puncture?


- ◆ Glucose, protein, nucleated cells
- ◆ Infectious workup (when pertinent)
- ◆ Autoimmune encephalitis panel (when pertinent)

EEG?

- ◆ Spot versus continuous

Meningitis


- Inflammation of meninges
 - Arachnoid, Pia (leptomeningitis)
 - Dura (pachymeningitis)
- **Infectious (LM)**
- Autoimmune (PM)
- Medications
- Neoplasm




© MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH. ALL RIGHTS RESERVED

Clinical Features

- Fever
- Meningismus
 - Kernig/Brudzinski → specific, NOT sensitive
- Altered mental status
 - meningoencephalitis



* <50% have all 3 features



Etiology

Acute

- Bacterial
 - Community-acquired
 - Streptococcus pneumoniae
 - Neisseria meningitidis (purpuric rash)
 - Listeria monocytogenes
 - >50 years, cell-mediated immunity deficiency
 - Healthcare-associated / Trauma
 - Staphylococci
 - Aerobic GNR
- Viral
 - HSV
 - VZV
 - Enterovirus
 - Arboviruses, HIV

Subacute/Chronic

- TB
 - Fungal
 - Cryptococcal, Coccidioides, Histoplasma, Blastomyces, Candida
 - Neoplastic
 - carcinomatosis
 - Autoimmune
 - Sarcoidosis, SLE
- Syphilis, Lyme, Parasites
- Drug-induced (NSAIDs, IVIG)

Evaluation

- CBC, BMP, Coags
- Blood Cx
- LP
 - CT before LP:
 - ↓ LOC or focal deficit
 - ↑ ICP or new seizure
 - Immunocompromised
 - Delay → blood cx & empiric tx
 - Opening pressure, cell count, glucose, protein, gram stain, cultures




Table 2. Cerebrospinal Fluid Findings In Central Nervous System Infections³

Infection	White Blood Cell Count (cells/mL)	Glucose Level	Protein Level
Bacterial Meningitis	Elevated (100-5000) Polymorphonuclear leukocytes predominate*	Decreased	Elevated
Viral Meningitis	Elevated (10-500) Lymphocytes predominate	Normal	Elevated
Fungal Meningitis	Normal to elevated (0-500) Lymphocytes predominate	Normal to decreased	Elevated
Tuberculous Meningitis	Normal to elevated (0-1000) Lymphocytes predominate	Decreased	Elevated
Brain Abscess	Normal to elevated (0-500) Mixed differential	Normal	Elevated

Note: In 10% of cases, lymphocytes predominate - see the **Lumbar Puncture** section on page 8 for details.

Management

- Empiric treatment → vancomycin/ceftriaxone + acyclovir
- Bacterial
 - Acute community-acquired → Vanc/CTX
 - Amp: >50 years, immunocompromised
 - NSGY/trauma → Vanc + Cefepime/meropenem
 - * Dex → mortality benefit in *S. Pneumoniae*

Complications (bacterial)

- Seizures
- Infarction
 - infectious vasculitis
- CVST
- Cerebral Edema & ↑ ICP
- Abscess

- Chronic → hydrocephalus, hearing loss, epilepsy, cognitive impairment