

COVER PAGE

STAT 608 Homework 03, Summer 2017

Please write your name and email address clearly below, then convert to PDF and attach as the first page of your homework upload.

NAME: RAJAN KAPOOR

EMAIL: r.kapoor@tamu.edu

45/49

(1) $H_0: \gamma_1 = \gamma_2$

$H_A: \gamma_1 \neq \gamma_2$

or $H_0: Y_j = \gamma_0 + \gamma_1 x_{j1} + \gamma_1 x_{j2} + \epsilon_j$

$H_A: Y_j = \gamma_0 + \gamma_1 x_{j1} + \gamma_2 x_{j2} + \epsilon_j$

$$F = \frac{RSS_{\text{reduced}} - RSS_{\text{full}}}{df_{\text{reduced}} - df_{\text{full}}} \div \frac{RSS_{\text{full}}}{df_{\text{full}}}$$

$$df_{\text{full}} = 60 - 3 = 57$$

$$df_{\text{reduced}} = 60 - 2 = 58$$

$$F \sim F_{2, 57}$$

If F is statistically significant, we favor the full model and reject the null hypothesis. i.e. p -value less than threshold (eg 0.001).

(2)

(2.1) Unbiased estimate of the error variance

$$s^2 = \frac{RSS}{df} = \frac{100}{30-4} = \frac{100}{26} = 38.46$$

(2.2) Possible numerical value of $\hat{\beta}_1$

$$t_c = \frac{\hat{\beta}_1 - 0}{se(\hat{\beta}_1)} \sim t\text{-distribution with } df = 26$$

$$p\text{-value} = 0.02$$

$$\Rightarrow P(T > |t_c|) = 0.02$$

For p-value of 0.02, the critical value is given by

$$\Rightarrow t_c = 2.479 \text{ or } t_c = -2.479$$

$$\frac{\hat{\beta}_1}{se(\hat{\beta}_1)} = 2.479 \text{ or } \frac{\hat{\beta}_1}{se(\hat{\beta}_1)} = -2.479$$

$$\Rightarrow \hat{\beta}_1 = 2.479 \times 0.5 \text{ or } \hat{\beta}_1 = -2.479 \times 0.5$$

$$\Rightarrow \hat{\beta}_1 = 1.2395 \text{ or } \hat{\beta}_1 = -1.2395$$

$$\begin{aligned} (2.3) \text{ 95\% CI for } \beta_2 &= \hat{\beta}_2 \pm t\left(\frac{0.05}{2}, 26\right) \times se(\hat{\beta}_2) \\ &= 1 \pm 0.3385 (0.25) \\ &\quad (0.915375, 1.084625) \end{aligned}$$

$$(2.4) \quad R^2 = 1 - \frac{RSS}{SST} = 0.9$$

$$\Rightarrow \frac{RSS}{SST} = 0.1 \Rightarrow SST = 1000$$

$$\Rightarrow RSS + SS_{reg} = 1000$$

$$\Rightarrow SS_{reg} = 900 \quad (\because RSS = 100)$$

$$\begin{aligned} (2.5) \quad F &= \frac{SS_{reg}/p}{RSS/(n-p-1)} \\ &= \frac{900/3}{100/26} = 3 \times 26 = 78 \sim F_{3, 26} \end{aligned}$$

At 5% level, $f_c = 2.975$
($\alpha = 0.05$)

F value $> f_c \Rightarrow$ Reject H_0

(3) (3.1) $H = X(X'X)^{-1}X'$ (Uppercase letters denote Matrix)

For idempotent matrix $AA' = A$

$$\begin{aligned}
 HH' &= X(X'X)^{-1}X' [X(X'X)^{-1}X']' \\
 &= X(X'X)^{-1}X' X'' (X'X)^{-1}X' \quad [\because (X'X)' = X'X \\
 &\quad \& (X^{-1})' = (X')^{-1}] \\
 &= X \underbrace{(X'X)^{-1}X'X}_{X''} (X'X)^{-1}X' \quad [\because X'' = X] \\
 &= X I (X'X)^{-1}X' \quad [\because X^{-1}X = I] \\
 &= X (X'X)^{-1}X' \quad [\because XI = X] \\
 &= H
 \end{aligned}$$

(3.2) $\text{Var}(\vec{\hat{e}} | X) = \text{Var}((I-H)\vec{y} | X)$

$$\begin{aligned}
 &= (I-H) \text{Var}(\vec{y} | X) (I-H)' \\
 &\quad [\because H = f(X) \& \\
 &\quad \text{Var}(A\vec{y}) = A \text{Var}(\vec{y}) A'] \\
 &= (I-H) \Sigma (I-H)' \\
 &= (I-H) \sigma^2 I (I-H)' \\
 &= (I-H) (I-H)' \sigma^2 \\
 &= (I-H) (I-H)' \sigma^2 \\
 &= (I - HI - IH' + HH') \sigma^2 \\
 &= (I - H - H + H) \sigma^2 \quad \because HH' = H \\
 &= (I-H) \sigma^2
 \end{aligned}$$

$$(4) \quad \vec{\hat{e}} = (I - H) \vec{y}$$

$$\vec{\hat{e}} = (I - X(X'X)^{-1}X') \vec{y} \quad \text{--- (1)}$$

To generate given expression, pre multiply $\vec{\hat{e}}$ by X'

$$\text{LHS} = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 1 & 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} \hat{e}_1 \\ \hat{e}_2 \\ \hat{e}_3 \\ \hat{e}_4 \end{bmatrix} = \begin{bmatrix} \hat{e}_1 + \hat{e}_3 \\ \hat{e}_1 + 2\hat{e}_2 + \hat{e}_4 \end{bmatrix}$$

$$\text{RHS} = X' (I - X(X'X)^{-1}X') \vec{y}$$

$$= (X'I - X'X(X'X)^{-1}X') \vec{y}$$

$$= (X' - I X') \vec{y}$$

$$= (X' - X') \vec{y} = 0_{2 \times 3}$$

Comparing second row in LHS & RHS.

$$\Rightarrow \hat{e}_1 + 2\hat{e}_2 + \hat{e}_4 = 0. \quad \text{Yes.}$$

(5)

$$\vec{y} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 2 & 1 \\ 1 & 2 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} \beta_A \\ \beta_B \end{bmatrix} + \vec{\hat{e}}$$

(6) Yes. The $\Phi\Phi$ plots of all predictor variables follow straight line, so no transformation is needed.

(a) For response variable, the $\Phi\Phi$ plot is not a straight line but becomes straight after log transformation.

The Y vs each predictor variable plots show increasing variance without transformation. Also box-cox transformation plot shows $\lambda=0$ lies within 95% Confidence interval of max. likelihood.

Since $\lambda=0$ corresponds to log transformation, the recommendation is correct.


(b) See page -

(c) From leverage plot, effects of high leverage points should be looked into before finalizing the model.


(d) Statistically insignificant predictors can be removed.

(e) Removing all predictors with insignificant p values at once is not recommended because then we'll be comparing model with subset of predictor variables with original model. p -value can only be used to remove one predictor variable at a time. Partial F -test should be used to decide if a subset of predictor variables should be removed at once.


Normal Q-Q Plot




Normal Q-Q Plot




Normal Q-Q Plot




Normal Q-Q Plot




Normal Q-Q Plot

















Normal Q-Q Plot




Normal Q-Q Plot










Normal Q-Q Plot



Normal Q-Q Plot





Call:

```
lm(formula = log(PrizeMoney) ~ DrivingAccuracy + GIR + PuttingAverage +
    BirdieConversion + SandSaves + Scrambling + PuttsPerRound)
```

Residuals:

	Min	1Q	Median	3Q	Max
Residuals	-1.71949	-0.48608	-0.09172	0.44561	2.14013

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.194300	7.777129	0.025	0.980095
DrivingAccuracy	-0.003530	0.011773	-0.300	0.764636
GIR	0.199311	0.043817	4.549	9.66e-06 ***
PuttingAverage	-0.466304	6.905698	-0.068	0.946236
BirdieConversion	0.157341	0.040378	3.897	0.000136 ***
SandSaves	0.015174	0.009862	1.539	0.125551
Scrambling	0.051514	0.031788	1.621	0.106788
PuttsPerRound	-0.343131	0.473549	-0.725	0.469601

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.6639 on 188 degrees of freedom

Multiple R-squared: 0.5577, Adjusted R-squared: 0.5412

F-statistic: 33.87 on 7 and 188 DF, p-value: < 2.2e-16