

COVER PAGE

STAT 608 Homework 07 Summer 2017

Please TYPE your name and email address below, then convert to PDF and attach as the first page of your homework upload.

NAME: RAJAN KAPOOR

EMAIL: r.kapoor@tanu.edu

35/35

Solution 1)

Assuming unrelated regression lines model

$$Y = \text{TOTALCHG}, x = \text{STAGE}$$

$$M_1 \begin{cases} Y = \beta_0 + \beta_1 x + e & \text{TRT} = 0 \\ Y = \beta_0 + \beta_2 + (\beta_1 + \beta_3)x + e & \text{TRT} = 1 \end{cases}$$

We call model M_1 as full model.

Now to test null hypothesis

$$H_0: \beta_3 = 0 \quad (\text{identical slopes})$$

$$H_A: \beta_3 \neq 0$$

The following is the reduced model

$$M_2 \begin{cases} Y = \beta_0 + \beta_1 x + e & (\text{TRT} = 0) \\ Y = \beta_0 + \beta_2 + \beta_2 x + e & (\text{TRT} = 1) \end{cases}$$

The null hypothesis can be tested using partial-F test

$$F = \frac{\text{RSS}(\text{reduced}) - \text{RSS}(\text{full})}{\text{df}(\text{red}) - \text{df}(\text{full})} \div \frac{\text{RSS}(\text{full})}{\text{df}_{\text{full}}}$$

For model M1

$$Y = 5.4760 + 2.4934x \quad \text{TRT}=0$$

$$Y = 7.7143 + 0.4785x \quad \text{TRT}=1$$

For reduced model

$$Y = 7.2298 + 1.5113x \quad \text{TRT}=0$$

$$Y = 5.91248 + 1.5113x \quad \text{TRT}=1$$

$$F = \frac{\frac{187.59 - 155.22}{1}}{\frac{155.22}{19}} = 3.9623$$

$$p\text{-value} = 0.001786 < \alpha (= 0.05)$$

\Rightarrow Reject Null Hypothesis

Solution 2

$$2.1) \quad \log \frac{p(x)}{1-p(x)} = \hat{\beta}_0 + \hat{\beta}_1 x$$

$$\Rightarrow p(x) = \frac{\exp(\hat{\beta}_0 + \hat{\beta}_1 x)}{\exp(\hat{\beta}_0 + \hat{\beta}_1 x) + \exp(\hat{\beta}_0 - \hat{\beta}_1 x)} p(x)$$

$$\Rightarrow p(x) = \frac{\exp(\hat{\beta}_0 + \hat{\beta}_1 x)}{1 + \exp(\hat{\beta}_0 - \hat{\beta}_1 x)} = \frac{1}{\exp(-\hat{\beta}_0 - \hat{\beta}_1 x) + 1}$$

at $x=3$

$$p(x) = \frac{1}{\exp[-(-2.643 + 0.674 \times 3)] + 1}$$

$$\Rightarrow P(x) = 0.3495$$

∴ probability that the insect will survive $= 1 - p(x)$
 $= 0.6504$

$$\text{Expected no. of surviving insects} = 200 \times (1 - p(x)) \\ = 130.$$

2.2 Odds on dying $= \frac{p(x)}{1 - p(x)} = \exp(\hat{\beta}_0 + \hat{\beta}_1 x)$

if x increases by 1 unit, odds on dying increases by $\phi = \exp(0.674)$
 $= 1.96207$

2.3 90% CI for $\phi = \exp(\hat{\beta}_1 \pm Z_{0.05} * \text{se}(\hat{\beta}_1))$ $\alpha = 0.1$
 $\alpha/2 = 0.05$

$$\exp(0.674 \pm 1.6448 \times 0.039)$$

$$(1.840, 2.092)$$

Solution 3

$$\log(\text{Sales}) = \beta_0 + \text{Time} + \text{Month}_i \quad i = 2, \dots, 12$$

The diagnostic plots and summary from R is attached.

Weakness - Variables Time, Month_2, Month_7 are not statistically significant
 - Case 32, 33, 41, 89 have $\text{standard res} > 2$.

Call:

```
lm(formula = log(Sales) ~ Month_2 + Month_3 + Month_4 + Month_5 +  
    Month_6 + Month_7 + Month_8 + Month_9 + Month_10 + Month_11 +  
    Month_12 + Time)
```

Residuals:

	Min	1Q	Median	3Q	Max
	-0.57520	-0.05166	-0.00739	0.07133	0.28116

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	5.8723775	0.0518913	113.167	< 2e-16	***
Month_2	-0.0522214	0.0635314	-0.822	0.413533	
Month_3	0.1907035	0.0635154	3.002	0.003572	**
Month_4	0.2474630	0.0635029	3.897	0.000201	***
Month_5	0.2270750	0.0634940	3.576	0.000595	***
Month_6	0.2138238	0.0634887	3.368	0.001167	**
Month_7	0.0690719	0.0634869	1.088	0.279874	
Month_8	0.2208295	0.0634887	3.478	0.000819	***
Month_9	0.3708060	0.0634940	5.840	1.07e-07	***
Month_10	0.4133160	0.0635029	6.509	6.13e-09	***
Month_11	0.7023337	0.0655759	10.710	< 2e-16	***
Month_12	1.4169818	0.0655707	21.610	< 2e-16	***
Time	0.0007670	0.0004756	1.613	0.110755	


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.1227 on 80 degrees of freedom


Multiple R-squared: 0.9111, Adjusted R-squared: 0.8978

F-statistic: 68.33 on 12 and 80 DF, p-value: < 2.2e-16


Diagnostic Plots:




$\text{lm}(\log(\text{Sales}) \sim \text{Month}_2 + \text{Month}_3 + \text{Month}_4 + \text{Month}_5 + \text{Month}_6 + \text{Month}_7 + \dots$



$\text{lm}(\log(\text{Sales}) \sim \text{Month}_2 + \text{Month}_3 + \text{Month}_4 + \text{Month}_5 + \text{Month}_6 + \text{Month}_7 + \dots$



Fitted values
 $\text{lm}(\log(\text{Sales}) \sim \text{Month}_2 + \text{Month}_3 + \text{Month}_4 + \text{Month}_5 + \text{Month}_6 + \text{Month}_7 + \dots)$



Leverage
 $\text{lm}(\log(\text{Sales}) \sim \text{Month}_2 + \text{Month}_3 + \text{Month}_4 + \text{Month}_5 + \text{Month}_6 + \text{Month}_7 + \dots)$

(4)

Solution 4

$$(4.1) \quad \begin{bmatrix} e_2 \\ e_3 \\ e_4 \\ \vdots \\ e_n \end{bmatrix}_Y = \rho \begin{bmatrix} e_1 \\ e_2 \\ e_3 \\ \vdots \\ e_{n-1} \end{bmatrix}_X + \begin{bmatrix} \epsilon_2 \\ \epsilon_3 \\ \epsilon_4 \\ \vdots \\ \epsilon_n \end{bmatrix} \quad \text{--- (M1)}$$

Clearly a linear regression of Y on X through the origin will estimate ρ . Since all 4 assumptions of Regression Model -

- Y is related to X by simple reg. model ($\because e_t = \rho e_{t-1} + \epsilon_t$)
 - ϵ_t are independent of each other
 - ϵ_t have common variance
 - ϵ_t are normally distributed with mean zero & common variance ($=1 \because$ standard normal). (least sq estimate)
- distribution given for ϵ_t

$$\hat{\rho} = \text{slope} = 0.412$$

- (4.2) YES. the valid S.E. for $\hat{\rho}$ should be 0.133.
Since model M1 is valid regression model (Linear).