



Why are we all using biased estimates of autocorrelation? and

Why are most time series simulation studies poor?


Rebecca Killick

Joint work with Colin Gallagher, Xiyan Tan, and Xinyi Li (Clemson)

Clemson Nov 2025



$$X_t = \phi X_{t-1} + N_t$$






- Provide better estimates of the (p)acf by adaptive shrinkage
- Give more intuition on causality
- Motivate better time series simulation studies.

Results




$$X_t = \phi X_{t-1} + N_t$$



Results



$$X_t = \theta N_{t-4} + N_t$$





For $y_t = x_t - \bar{x}$, sample ACF at lag h , $\rho(h)$ minimizes:

$$Q_h(\rho_h) = \sum_{t=h+1}^{n+h} (y_t - \rho_h y_{t-h})^2,$$

To shrink toward zero (reduce variance) add a penalty:

$$Q_h(\rho_h) + \lambda_h \left(\sum_{t=1}^n y_t^2 \right) (\rho_h)^2$$

To move toward a bias corrected estimator, $\hat{\rho}_b$:

$$Q_h(\rho_h) + \lambda_h \left(\sum_{t=1}^n y_t^2 \right) (\rho_h - \hat{\rho}_b)^2$$

Let the data tell us which one to use and how much to shrink.



We minimize

$$Q_h(\rho_h) + \lambda_h \left(\sum_{t=1}^n y_t^2 \right) (\rho_h - \rho_\tau(h))^2$$

The solution is

$$\tilde{\rho}(h) = w_h \rho_\tau(h) + (1 - w_h) \hat{\rho}(h),$$

where $w_h = \lambda_h / (1 + \lambda_h)$, and $\hat{\rho}(h)$ is the ordinary ACF at lag h .

Note: General framework includes, ordinary ACF, shrinkage toward zero, bias correction, Banded & Tapered, ...



We use plug-in estimators of the tuning parameters.

Shrink toward zero if $\hat{\rho}_b(h)$ is small, move toward bias correction if larger.

$$\rho_\tau(h) = \begin{cases} 0 & |\hat{\rho}_b(h)| < \ell_h \\ \hat{\rho}_b(h) & |\hat{\rho}_b(h)| > \ell_h \end{cases}$$

Select $\lambda_h(w)$ to move more aggressively near boundaries, and to smoothly transition between making smaller and increasing magnitude:

$$\lambda_h = \begin{cases} \frac{h(\ell_h - r)}{r^2} & r < \ell_h \\ \frac{10 \log_{10}(n) h(r - \ell_h)(1 - \ell_h)}{(1 - r)^2} & r > \ell_h, \end{cases}$$

where $r = |\hat{\rho}_b(h)|$.



ℓ_h controls the asymptotic distribution of $\tilde{\rho}(h)$ for $\rho(h) = 0$:

1. If $\sqrt{n}\ell_h \rightarrow 0$ as $n \rightarrow \infty$, $\hat{\rho}$ and $\tilde{\rho}(h)$ have same limiting distribution.
2. If $\sqrt{n}\ell_h \rightarrow C$ as $n \rightarrow \infty$, $\sqrt{n}\tilde{\rho}(h)$ has a non-normal limit with smaller variance than $\sqrt{n}\hat{\rho}(h)$.
3. If $\ell_h \rightarrow 0$ and $\sqrt{n}\ell_h \rightarrow \infty$ as $n \rightarrow \infty$, $\sqrt{n}\tilde{\rho}(h) \rightarrow 0$ in probability.

$\sqrt{n}(\tilde{\rho}(h) - \rho(h))$ has same limit as $\sqrt{n}(\hat{\rho}(h) - \rho(h))$, when $\rho(h) \neq 0$, \\\ We use

$$\ell_h = c_h \sqrt{\log(n)/n},$$

where c_h depends on the estimated ACF.

Similar development for the PACF at lag h , $\pi(h)$ gives:

$$\tilde{\pi}(h) = w_h \pi_\tau(h) + (1 - w_h) \hat{\pi}(h),$$

where $w_h = \lambda_h / (1 + \lambda_h)$, and $\hat{\pi}(h)$ is the ordinary PACF at lag h . Use ℓ_h , $\hat{\pi}_b(h)$, and λ_h as above (but using PACF).


$$\pi_\tau(h) = \begin{cases} 0 & |\hat{\pi}_b(h)| < \ell_h \\ \hat{\pi}_b(h) & |\hat{\pi}_b(h)| > \ell_h. \end{cases}$$

Asymptotic results again controlled by ℓ_h (we use our plug-in tuning parameters below).

Results




$$X_t = \phi X_{t-1} + N_t$$



Results




$$X_t = \theta N_{t-4} + N_t$$



Results




(c) $X_t = \phi X_{t-1} + N_t$





Results



(d) $X_t = \phi X_{t-4} + N_t$



(a) $X_t = 0.95N_{t-4} + N_t$ 





- Want to simulate across the causal space
- For a wide set of AR, MA and ARMA processes


HOW?



- Want to simulate across the causal space
- For a wide set of AR, MA and ARMA processes


HOW?

- JTSA published 44 papers in 2024, 39 had an AR simulation
- Only 2 papers selected coefficients randomly (in AR(1))
- Of the remaining 37, 25 used a single AR parameter setting!
- Only 12/39 considered orders larger than 1!






Degree of Correlation: AR(3), phi3= 0.3





Degree of Correlation: AR(3), phi3= 0.9





AR(p):

$$Y_t = \sum_{j=1}^p \phi_j Y_{t-j} + \epsilon_t \quad t = 0, \pm 1, \pm 2, \dots,$$

Causal:

$$Y_t = \sum_{j=0}^{\infty} c_j \epsilon_{t-j} \quad \sum_{j=0}^{\infty} |c_j| < \infty;$$

How to check causality?



AR(p):

$$Y_t = \sum_{j=1}^p \phi_j Y_{t-j} + \epsilon_t \quad t = 0, \pm 1, \pm 2, \dots,$$

Causal:

$$Y_t = \sum_{j=0}^{\infty} c_j \epsilon_{t-j} \quad \sum_{j=0}^{\infty} |c_j| < \infty;$$

How to check causality?

Roots of the polynomial:

$$\Phi(x) = 1 - \phi_{p,1}x - \phi_{p,2}x^2 - \cdots - \phi_{p,p}x^p$$

Are the roots outside the unit circle?


Known solution spaces



AR1: $\phi_{1,1} \in (-1, 1)$

AR2:

What about higher orders?



PACF at lag h : $\alpha(h)$.

$$|\alpha(h)| < 1 \quad \text{for} \quad 1 \leq h \leq p \quad \text{and} \quad \alpha(h) = 0 \quad \text{for} \quad h > p.$$

Ramsey (1974) noted a 1:1 mapping. Any point in $(-1, 1)^p$ corresponds to a unique point in the causal parameter space:

$$(\phi_{p,1}, \dots, \phi_{p,p}) = g(\alpha(1), \dots, \alpha(p))$$

$$(\alpha(1), \dots, \alpha(p)) = g^{-1}(\phi_{p,1}, \dots, \phi_{p,p}).$$



The map g can be described via the Durbin-Levinson algorithm, which provides

$$\phi_{k,k} = \alpha(k)$$

$$\phi_{k,j} = \phi_{k-1,j} - \phi_{k,k}\phi_{k-1,k-j} \quad k > 1, 1 < j < k-1.$$

Equivalent approaches



Let R_p denote the causal parameter space for an order p AR. Any of the following conditions can be used to assess if $\phi \in R_p$.


1. The roots of $\Phi(x) = 1 - \phi_{p,1}x - \phi_{p,2}x^2 - \cdots - \phi_{p,p}x^p$ are outside the unit circle.
2. $|\phi_{h,h}| < 1$ for $1 \leq h \leq p$, and for each $k = 2, \dots, p$ and $j = 1, \dots, k$,
 $\phi_{k,j} = \phi_{k-1,j} - \phi_{k,k}\phi_{k-1,k-j}$.
3. $|\phi_{p,p}| < 1$, $\phi_{2i,2i} > -1$ for $i < p/2$, and for each $k = 2, \dots, p$ and
 $j = 1, \dots, k$,

$$\sum_{j=1}^k \phi_{k,j} < 1 \quad \text{and} \quad \sum_{j=1}^k (-1)^{j+1} \phi_{k,j} > -1,$$


with $\phi_{k,j} = \phi_{k-1,j} - \phi_{k,k}\phi_{k-1,k-j}$.

4. For $p > 1$, $(\phi_{p-1,1}, \dots, \phi_{p-1,p-1}) \in R_{p-1}$, $|\phi_{p,p}| < 1$ and
 $\phi_{p,j} = \phi_{p-1,j} - \phi_{p,p}\phi_{p-1,p-j}$ for $j = 1, \dots, p-1$.

Causal Region AR3



Causal Region AR3





Dropbox Link



Need to summarise: $R_n(i, j) = \text{corr}(Y_i, Y_j)$, for $1 \leq i, j \leq n$

Measure of total correlation:


$$\rho(\{Y_t\}) = 1 - \det(R_n) = 1 - \prod_{k=1}^p (1 - \alpha(k)^2)^{p+1-k}.$$

where \det denotes the determinant.

$\rho(\{Y_t\}) \rightarrow 1$, high correlation

$\rho(\{Y_t\}) \rightarrow 0$, low correlation

AR2 Total Correlation





1. Generate $\{X_i\}_{i \in (1, \dots, p)}$ independently having, $0 \leq \ell, u \leq 1$
Uniform density


$$f(x) = \frac{1}{(u - \ell)} \mathbb{I}_{\ell < x < u}.$$

2. Generate $\{R_i\}$ independent of $\{X_i\}$,
 $P(R = 1) = P(R = -1) = 0.5$.
3. Let each $\alpha(i) = R_i X_i$ and solve for $\{\phi_{p,j}\}$.

Larger values of (ℓ, u) for one or more i correspond to more correlation.




$\ell = 0.8, u = 0.9, 10,000$ simulations, $\rho(\{Y_t\}) > 0.9$, in 90% of simulations.






MSE for YW AR(p) fits based on $\hat{\pi}$ vs $\tilde{\pi}$, 10,000 replications

Parameters randomly generated over parameter space.





Pairwise MSE for YW AR(1) fits based on $\hat{\pi}$ vs $\tilde{\pi}$.





- Provided a new approach for (P)ACF estimation adaptively shrinking towards zero and target at each lag
 - Improves over sample ACF, often beats Banded and Tapered too
 - Computationally fast
 - Adapts to a wide set of correlation structures
-
- Showed how PACF and AR coefficients are linked
 - Used this link to describe causality regions in higher dimensions
 - Visualized up to AR(4)
 - Utilized the new description for simulation study generation

Available in PenalizedCor on Github.