

Regional and Global Warming and Statistics: A statisticians view of the surge in warming debate

Rebecca Killick(r.killick@lancs.ac.uk)

Joint work with Claudie Beaulieu (Uni California Santa Cruz)

Parts with: Adelia Johnson, Colin Gallagher, Robert Lund, Xueheng Shi

Warwick April 2026




Detection of changes in variance of oceanographic
time-series using changepoint analysis

Rebecca Killick

work with Idris Eckley, Phil Jonathan and Kevin Ewans


Lancaster University





Rebecca Killick (Lancaster University)

GOMOS Talk


1 / 23









Model development





Model development



- What is the warming debate in climate science?
- What are changepoints?
- The story of a statistical solution
- Some frustrations


Where my interest began

School of
Mathematical Sciences

Lancaster
University




Global Mean Temp 2016



Global Mean Temp




2016



Global Mean Temp



Today



Is climate change speeding up? Here's what the science says.

This year's record temperatures have some scientists concerned that the pace of warming may be accelerating. But not everyone agrees.

By [Chris Mooney](#) and [Shannon Osaka](#)

December 26, 2023 at 6:30 a.m. EST


For the past several years, a small group of scientists has warned that sometime early this century, the rate of global warming — which has remained largely steady for decades — might accelerate. Temperatures could rise higher, faster. The drumbeat of weather disasters may become more insistent.

Where is Statistics?




The increased rate of global warming

Values are relative to the 1951-1980 global mean temperature, in degrees Celsius



Note: 2023 is an estimate based on values from January through November.

Where is Statistics? NYT




What are changepoints?



For data $y_1, \dots, y_n$, if a changepoint exists at τ , then $y_1, \dots, y_\tau$ differ from $y_{\tau+1}, \dots, y_n$ in some way.

There are many different types of change.




Change in mean



Assume we have time-series data where

$$Y_t | \theta_t \sim N(\theta_t, 1),$$

but where the means, θ_t , are piecewise constant through time.



We want to infer the number and position of the points at which the mean changes. One approach:

Likelihood Ratio Test

To detect a single changepoint we can use the likelihood ratio test statistic:

$$LR = \max_{\tau} \{ \ell(y_{1:\tau}) + \ell(y_{\tau+1:n}) - \ell(y_{1:n}) \}.$$

We infer a changepoint if $LR > \beta$ for some (suitably chosen) β . If we infer a changepoint its position is estimated as

$$\tau = \arg \max \{ \ell(y_{1:\tau}) + \ell(y_{\tau+1:n}) - \ell(y_{1:n}) \}.$$

This can test can be repeatedly applied to new segments to find multiple changepoints.

Define m to be the number of changepoints, with positions $\tau = (\tau_0, \tau_1, \dots, \tau_{m+1})$ where $\tau_0 = 0$ and $\tau_{m+1} = n$.


Then one application of the Likelihood ratio test can be viewed as

$$\min_{m \in \{0, 1\}, \tau} \left\{ \sum_{i=1}^{m+1} [-\ell(y_{\tau_{i-1}:\tau_i})] + \beta m \right\}$$

Repeated application is thus aiming to minimise

$$\min_{m, \tau} \left\{ \sum_{i=1}^{m+1} [-\ell(y_{\tau_{i-1}:\tau_i})] + \beta m \right\}$$

Problem




- How many changes?
- Where are the changes? 2^{n-1} possible solutions!

PELT in a nutshell



- Dynamic programming allows us to only worry about the location of the *last* change.
- Pruning means that as we go through the data we are smart about which locations are potential last change locations.






Let $0 < t < s < T$, if


$$F(t) + \mathcal{C}(y_{(t+1):s}) < F(s)$$

then at any future time $T > s$, t can never be the optimal last changepoint prior to T .

We can prove that, under certain regularity conditions, the expected computational complexity will be $\mathcal{O}(n)$.




More complicated changes






- From a publication in Marine Ecology (not the only one)
- Used the Rodionov (2004) method.
- Very popular but cannot deal with trend or autocorrelation.







- potentially hundreds or thousands of series
- no time to consider the format of change for each
- need to include both the potential for trends and also red noise (autocorrelation).






EnvCpt: select the most parsimonious but accurate model for the data. Simple to extend with other types of models.





Join-pin Models



Fearnhead, Maidstone, Letchford, JCGS (2019)

$$Y_t = \theta_i + \frac{\theta_{i+1} - \theta_i}{\tau_{i+1} - \tau_i}(t - \tau_i) + Z_t$$

where $Z_t \sim N(0, \sigma^2)$.

Implemented in the `cpop` R package.


$$Y_t = \theta_i + \frac{\theta_{i+1} - \theta_i}{\tau_{i+1} - \tau_i}(t - \tau_i) + Z_t$$

but now Z_t is AR(p), $Z_t = \phi_1 Z_{t-1} + \phi_2 Z_{t-2} + \dots + \phi_p Z_{t-p} + \epsilon_t$.

Challenge:

- Definition of join-pin
- Fitting AR parameter across segments:
 - Fixed we use EM algorithm
 - Varying we can embed in PELT


Join-pin Models



So what?



What would we need to see (in the statistically preferred model).





Global warming is NOT surging, scientists say - despite record-breaking temperatures

- Scientists find no evidence for a 'surge' in global temperatures
- [READ MORE: 2023 was officially the hottest year on RECORD](#)

By JONATHAN CHADWICK FOR MAILONLINE

PUBLISHED: 14:01, 15 October 2024 | UPDATED: 07:52, 16 October 2024



Share



138

shares




407

View comments


From the **UK's hottest day** to the **hottest year on record globally**, there's no doubt some worrying temperature records have been broken in recent years.

Many people think the rate of **global warming** has dramatically accelerated or 'surged' over the past 15 years – and is a cause of more extreme weather.


But a new study says there is not any statistical evidence for this so-called 'surge' or 'leap'.



Extension to gridded




Extension to gridded




Extension to gridded




B Southeast Mexico




C Gulf of Mexico




D Bolivia




E East Mediterranean




F Southeast China



G New Zealand



H North Pacific



- Detecting and documenting changepoints improves analyses ...
- ... and can help answer pertinent questions in different domains
- Extending to more complex model structures is interesting statistically
- Extending to monthly/grid point is interesting climatically

- I enjoy working with different disciplines ...
- ... as often it sparks my next research challenge.

PELT: <https://doi.org/10.1080/01621459.2012.737745>

Model Choice: <https://doi.org/10.1175/JCLI-D-17-0863.1> &
<https://doi.org/10.1002/qre.2712>

LMvsCpts: <https://doi.org/10.1007/s11222-017-9731-0> &
<https://doi.org/10.1002/env.2568>

Warming Surge: <https://doi.org/10.1038/s43247-024-01711-1>

Regional Surge:
<https://www.researchsquare.com/article/rs-7731926/v1>

