

Sampling and Aliasing

Rkka

July 30, 2024

Abstract

This article is about sampling and aliasing problems that I have encountered while studying numerical simulation of wave optics with computer to implement computer generated holography (CGH).

1 The Nyquist-Shannon sampling theorem

1.1 Statement of the theorem

If a function $x(t)$ contains no frequency higher than B hertz, then it can be completely determined from its ordinates at a sequence of points spaced less than $1/(2B)$ seconds apart.

We can rewrite the above theorem with the following expressions : $f_s \geq 2B$. In other words, when we denote a sampling interval as Δx and a period of an original signal as λ , the theorem is : $\Delta x < (\lambda/2)$. Intuitively, it says that we need to sample the original signal sufficiently densely in order to represent the signal without loss of information.

1.2 Motivation

The sampling theorem is required to avoid a type of distortion called *aliasing*.

2 Sampling theory

2.1 Dirac comb

The *Dirac comb* function is defined as :

$$\text{comb}_T(t) \equiv \sum_{n=-\infty}^{\infty} \delta(t - nT)$$

where $\delta(t)$ is the Dirac delta function. In the above definition the comb function has a period T . This function is important in sampling theory because every periodically sampled function $f_s(x)$ can be expressed as : $f_s(x) = f(x)\text{comb}_T(x)$ where T is a sampling period.

To understand a useful fourier transform property of the comb function, we need the following lemma :

$$\textbf{Lemma. } \delta(u) = \int_{-\infty}^{\infty} dx \cdot e^{-i2\pi ux}$$

$$\textbf{Proof. } \mathcal{F}^{-1}[\delta(u)] = \int_{-\infty}^{\infty} du \cdot \delta(u) e^{i2\pi ux} = 1$$

$$\rightarrow \mathcal{F}[\mathcal{F}^{-1}[\delta(u)]] = \delta(u) = \int_{-\infty}^{\infty} dx \cdot 1 \cdot e^{-i2\pi ux} = \int_{-\infty}^{\infty} dx \cdot e^{-i2\pi ux}$$

$$\text{Hence, } \delta(u) = \int_{-\infty}^{\infty} dx \cdot e^{-i2\pi ux}$$

(QED)

Now, we can prove the following useful theorem :

$$\textbf{Theorem. } \mathcal{F}[\text{comb}_T(t)] = \frac{1}{T} \text{comb}_{\frac{1}{T}}(u)$$

Proof. First, consider a Fourier series of $\text{comb}_T(t)$:

$$\text{comb}_T(t) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi \frac{n}{T} t} \text{ with } c_n = \frac{1}{T} \int_{-T/2}^{T/2} dt \cdot \text{comb}_T(t) e^{-i2\pi \frac{n}{T} t}$$

$$\rightarrow \text{comb}_T(t) = \frac{1}{T} \sum_{n=-\infty}^{\infty} e^{i2\pi \frac{n}{T} t}$$

Then, with a Fourier transform :

$$\mathcal{F}[\text{comb}_T(t)] = \int_{-\infty}^{\infty} dt \cdot \text{comb}_T(t) e^{-i2\pi ut} = \int_{-\infty}^{\infty} dt \cdot \frac{1}{T} \sum_{n=-\infty}^{\infty} e^{i2\pi \frac{n}{T} t} e^{-i2\pi ut}$$

$$= \frac{1}{T} \sum_{n=-\infty}^{\infty} \int_{-\infty}^{\infty} dt \cdot e^{-i2\pi(u - \frac{n}{T})t} = \frac{1}{T} \sum_{n=-\infty}^{\infty} \delta(u - \frac{n}{T})$$

$$= \frac{1}{T} \text{comb}_{\frac{1}{T}}(u)$$

(QED)

3 Aliasing

3.1 Definition

In signal processing, aliasing is the overlapping of frequency components[1]. For example, suppose we sample a high frequency(f_1) signal $x_1(t)$ with a sampling frequency f_s lower than the Nyquist frequency($f_s < f_{Nyquist} = 2f_1$). And denote the resulting sampled signal $x_1[n]$. The problem is, there is a lower frequency(f_2) signal $x_2(t)$ such that when we sample $x_2(t)$ with the same sampling frequency f_s , the sampled signal $x_2[n]$ becomes identical to $x_1[n]$: $x_1[n] = x_2[n]$.

In other words, when we sample a signal sparsely, then high-frequency information is lost and thus appears as a lower-frequency signal.




Figure 1: Aliasing. Red line represents original signal and dotted line represents sampled signal. You can see the original signal is sampled as a lower frequency signal.

3.2 Mathematical description

References

- [1] Nyquist–Shannon sampling theorem. In Wikipedia, The Free Encyclopedia.