

Boost 2013 Report Title

Report of BOOST2013, hosted by the University of Arizona, 12th-16th of August 2013.

D. Adams¹, A. Arce², L. Asquith³, J. Backus Mayes⁴, E. Bergeaas Kuutmann⁵, J. Berger⁶, D. Bjergaard², L. Bryngemark⁷, A. Buckley⁸, J. Butterworth⁹, M. Cacciari¹⁰, M. Campanelli⁹, T. Carli¹¹, M. Chala¹², B. Chapleau¹³, C. Chen¹⁴, J.P. Chou¹⁵, Th. Cornelissen¹⁶, D. Curtin¹⁷, M. Dasgupta¹⁸, A. Davison⁹, F. de Almeida Dias¹⁹, A. de Cosa²⁰, A. de Roeck¹¹, C. Debenedetti⁸, C. Doglioni²¹, S. D. Ellis²², F. Fassi²³, J. Ferrando²⁴, S. Fleischmann¹⁶, M. Freytsis²⁵, M.L. Gonzalez Silva²⁶, S. Gonzalez de la Hoz²³, F. Guescini²¹, Z. Han²⁷, A. Hook⁴, A. Hornig²², E. Izaguirre⁴, M. Jankowiak⁴, J. Juknevich²⁸, M. Kaci²³, D. Kar²⁴, G. Kasieczka²⁹, R. Kogler³⁰, A. Larkoski⁴, P. Loch³¹, D. Lopez Mateos²⁷, S. Marzani³², L. Masetti³³, V. Mateu³⁴, D. W. Miller³⁵, K. Mishra³⁶, P. Nef⁴, K. Nordstrom²⁴, E. Oliver Garcia²³, J. Penwell³⁷, J. Pilot³⁸, T. Plehn²⁹, S. Rappoccio³⁹, A. Rizzi⁴⁰, G. Rodrigo²³, A. Safonov⁴¹, G. P. Salam^{10,11}, J. Salt²³, S. Schaetzel²⁹, M. Schioppa⁴², A. Schmidt²⁹, J. Scholtz²², A. Schwartzman⁴, M. D. Schwartz²⁷, M. Segala⁴³, M. Son⁴⁴, G. Soyez⁴⁵, M. Spannowsky³², I. Stewart³⁴, D. Strom⁴⁶, M. Swiatlowski⁴, V. Sanchez Martinez²³, M. Takeuchi²⁹, J. Thaler³⁴, E. Thompson¹, N. V. Tran³⁶, C. Vermilion²⁵, M. Villaplana²³, M. Vos²³, J. Wacker⁴, and J. Walsh⁴⁷

¹ Columbia University, Nevis Laboratory, Irvington, NY 10533, USA

² Duke University, Durham, NC 27708, USA

³ Argonne National Laboratory, Lemont, IL 60439, USA

⁴ SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA

⁵ Deutsches Elektronen-Synchrotron, DESY, D-15738 Zeuthen, Germany

⁶ Cornell University, Ithaca, NY 14853, USA

⁷ Lund University, Lund, SE 22100, Sweden

⁸ University of Edinburgh, EH9 3JZ, UK

⁹ University College London, WC1E 6BT, UK

¹⁰ LPTHE, UPMC Univ. Paris 6 and CNRS UMR 7589, Paris, France

¹¹ CERN, CH-1211 Geneva 23, Switzerland

¹² CAFPE and U. of Granada, Granada, E-18071, Spain

¹³ McGill University, Montreal, Quebec H3A 2T8, Canada

¹⁴ Iowa State University, Ames, Iowa 50011, USA

¹⁵ Rutgers University, Piscataway, NJ 08854, USA

¹⁶ Bergische Universitaet Wuppertal, Wuppertal, D-42097, Germany

¹⁷ YITP, Stony Brook University, Stony Brook, NY 11794-3840, USA

¹⁸ University of Manchester, Manchester, M13 9PL, UK

¹⁹ UNESP - Universidade Estadual Paulista, Sao Paulo, 01140-070, Brazil

²⁰ INFN and University of Naples, IT80216, Italy

²¹ University of Geneva, CH-1211 Geneva 4, Switzerland

²² University of Washington, Seattle, WA 98195, USA

²³ Instituto de Física Corpuscular, IFIC/CSIC-UVEG, E-46071 Valencia, Spain

²⁴ University of Glasgow, Glasgow, G12 8QQ, UK

²⁵ Berkeley National Laboratory, University of California, Berkeley, CA 94720, USA

²⁶ Universidad de Buenos Aires, AR-1428, Argentina

²⁷ Harvard University, Cambridge, MA 02138, USA

²⁸ Weizmann Institute, 76100 Rehovot, Israel

²⁹ Universitaet Hamburg, DE-22761, Germany

³⁰ Universitaet Heidelberg, DE-69117, Germany

³¹ University of Arizona, Tucson, AZ 85719, USA

³² IPPP, University of Durham, Durham, DH1 3LE, UK

³³ Universitaet Mainz, DE 55099, Germany

³⁴ MIT, Cambridge, MA 02139, USA

³⁵ University of Chicago, IL 60637, USA

³⁶ Fermi National Accelerator Laboratory, Batavia, IL 60510, USA

³⁷ Indiana University, Bloomington, IN 47405, USA

³⁸ University of California, Davis, CA 95616, USA

³⁹ Johns Hopkins University, Baltimore, MD 21218, USA

⁴⁰ INFN and University of Pisa, Pisa, IT-56127, Italy

⁴¹ Texas A & M University, College Station, TX 77843, USA

⁴² INFN and University of Calabria, Rende, IT-87036, Italy

⁴³ Brown University, Richmond, RI 02912, USA

⁴⁴ Yale University, New Haven, CT 06511, USA

⁴⁵ CEA Saclay, Gif-sur-Yvette, FR-91191, France

⁴⁶ University of Illinois, Chicago, IL 60607, USA

⁴⁷ University of California, Berkeley, CA 94720, USA

Abstract Abstract for BOOST2013 report

Keywords boosted objects · jet substructure · beyond-the-Standard-Model physics searches · Large Hadron Collider

1 Introduction

Jet substructure has been around a while now, and it's time to study the correlations between the plethora of observables that have been developed and used. Previous BOOST reports [1, 2, 3] studied some of these things.

2 Monte Carlo Samples

Give details about how the samples we use have been generated.

3 Jet Algorithms and Grooming Approaches

Describe the jet algorithms and grooming approaches that we will use in the report. Give the nomenclature that we will use to refer to e.g. the groomed mass in the rest of the report.

4 Substructure Variables/Taggers

Describe the specific substructure variables and tagging approaches that we will be using in this report e.g. n-subjettiness, Q-jets, HTT, JH tagger. Give the nomenclature that we will use to refer to these variables/taggers in the rest of the report.

5 Boosted W -Tagging

In this section we study the performance of various jet algorithms in combination with jet substructure variables/taggers in terms of the identification of a boosted hadronically decaying W signal. For each jet algorithm we produce Receiver Operating Characteristic (ROC) curves that elucidate the performance of various variables that are capable of providing discrimination between a hadronic W signal and a QCD jet. These variables are then combined in a Boosted Decision Tree (BDT) and the performance of the resulting BDT discriminant explored through ROC curves to understand the degree to which variables are correlated and exploiting the same information. These studies are repeated in

different kinematic regimes, to explore both the performance and correlations as a function of the jet boost, and where substructure approaches may break down.

5.1 Methodology

These studies use the $X \rightarrow WW$ samples as signal and the XXX samples to model the QCD background.

Jets are reconstructed using the XXX jet algorithms described in the previous section. The following event selection is then applied to these samples....(presumably this will vary depending on which kinematic bin is used, as will the actual samples used - maybe summarize in a table).

Figure 1 shows background versus signal in some basic kinematic distributions. *Do we want to reweight signal kinematics to background or vice versa? Do we want to study quarks/gluons separately?*

Go on to explain how we produce the ROC curves.

5.2 Performance at Moderate Boosts

(this section is to cover the W -tagging performance for jet p_T 200-300 GeV and 500-600 GeV using $\sqrt{s} = 8$ TeV samples)

5.2.1 Single Variable Performance

Show plots of signal versus background for all single variables investigated.

Figure 2 the compares signal and background in the mass distributions for the different groomers, and Figure 3 in the different substructure variables.

Figure 4 shows the single variable ROC curves in the p_T 500 GeV bin for the anti- k_T $R=0.8$ algorithm, compared to the ROC curve for a BDT combination of all the variables. One can see that the best performant single variables for a reasonable signal efficiency are the groomed/filtered masses, which all have a similar level of performance with the exception of the soft drop mass with $\beta = -1$. *Would be good to split this into two plots, one using the masses and one for other variables, or somehow make the mass and other variable curves more distinct from one another by using same colour for all the mass curves.*

We want to look also at:

- *Dependence on R . So have the same single variable ROC for e.g. $R=1.2$, $R=0.4$. Then possibly have another plot which compares the best single variable (e.g. groomed mass) for different R .*




Fig. 1 Comparisons of the QCD background to the WW signal in the p_T 500 GeV bin using the anti- k_T $R=0.8$ algorithm: basic kinematic distributions.

- *Dependence on p_T .* Again want to repeat the plot for different kinematic bins, and then have a plot which compares the best performance in each kinematic bin to see the dependence of performance on kinematics.

5.2.2 Combined Performance

Figure 5 shows the BDT combinations of each mass variable with every other variable considered in the p_T 500 GeV bin using the anti- k_T $R=0.8$ algorithm. Can we drop the combinations of mass + mass from these plots to make them clearer? Also would be good to put the single variable mass curve on these plots, so you can see how much improvement the combination gives, and the “all variables” curve.

No combination with other variables can recover the poor performance of the ungroomed mass and the soft drop mass with $\beta = -1$. The other groomed/filtered masses are all most improved by combination with the $C_2^{\beta=1}$ energy correlation function. Show 2-D correlation plot of $C_2^{\beta=1}$ vs groomed mass - show that it is largely uncorrelated. Now show a plot which compares on one plot the best combined performance for each mass + X. e.g. mass + $C_2^{\beta=1}$, and compared also to the all variables curve. This plot is just for one R and one kinematic bin.

Repeat these studies for different R and different kinematic bins. Finally make plots which compare best combined performance for different R and kinematics.




Fig. 2 Comparisons of the QCD background to the WW signal in the p_T 500 GeV bin using the anti- k_T $R=0.8$ algorithm: leading jet mass distributions.




Fig. 3 Comparisons of the QCD background to the WW signal in the p_T 500 GeV bin using the anti- k_T R=0.8 algorithm: substructure variables.




Fig. 4 The ROC curve for all single variables considered for W tagging in the p_T 500 GeV bin using the anti- k_T $R=0.8$ algorithm.




Fig. 5 The BDT combinations of each mass variable with every other variable considered in the p_T 500 GeV bin using the anti- k_T $R=0.8$ algorithm.




Fig. 6 2-D plots showing the correlation between mMDT mass and various substructure variables in the p_T 500 GeV bin using the anti- k_T $R=0.8$ algorithm.

Do we want to look at other combinations of variables which don't involve mass? Practically I think we will always be making mass + X though.

5.3 Performance at High Boosts

(this section is to cover the W -tagging performance for jet p_T 1-1.1 TeV and > 1.5 TeV using $\sqrt{s} = 14$ TeV samples)

Maybe we don't need to divide into different medium/high boost sections.

6 Top Tagging

Top tagging studies go here.

7 Quark-Gluon Tagging

q/g tagging studies go here.

8 Summary & Conclusions

This report discussed the correlations between observables and looked forward to jet substructure at Run II

of the LHC at 14 TeV center-of-mass collisions energies.

Acknowledgements

We thank the Department of Physics at the University of Arizona and for hosting the conference at the Little America Hotel. We also thank Harvard University for hosting the event samples used in this report. We also thank Hallie Bolonkin for the BOOST2013 poster design and Jackson Boelts' ART465 class (fall 2012) at the University of Arizona School of Arts VisCom program. (NEED TO ASK PETER LOCH FOR MORE ACKNOWLEDGEMENTS)

References

1. A. Abdesselam, E. B. Kuutmann, U. Bitenc, G. Brooijmans, J. Butterworth, et al., *Boosted objects: A Probe of beyond the Standard Model physics*, *Eur.Phys.J.* **C71** (2011) 1661, [arXiv:1012.5412].
2. A. Altheimer, S. Arora, L. Asquith, G. Brooijmans, J. Butterworth, et al., *Jet Substructure at the Tevatron and LHC: New results, new tools, new benchmarks*, *J.Phys.* **G39** (2012) 063001, [arXiv:1201.0008].
3. A. Altheimer, A. Arce, L. Asquith, J. Backus Mayes, E. Bergeas Kuutmann, et al., *Boosted objects and jet substructure at the LHC*, arXiv:1311.2708.