

Selective Extraction of Lu, La, Ho from Aqueous Solution by Liquid-Liquid Extraction

K. Brown, L. Kumar, X. Kumar, Y. Kim

School of Chemical Engineering, University of Queensland, Brisbane QLD 4072, Australia

Abstract

Isotherm data for Y adsorption fit the Langmuir model with $R^2 = 0.956$. Recovery of Sc reached 95.9% under optimized precipitation conditions. Saponification of D2EHPA improved Eu extraction efficiency by 79.9 percentage points. Solvent extraction of Lu from aqueous phase showed high selectivity at pH 3.6. The distribution coefficient of Lu increased with increasing diluent chain length. Isotherm data for Yb adsorption fit the Langmuir model with $R^2 = 0.973$. Temperature significantly affected Dy extraction, with optimum conditions at 52 °C. Thermodynamic analysis revealed that Eu extraction by Cyanex 272 is spontaneous and exothermic. Contact time experiments demonstrated that Tb equilibrium was reached within 57 min. The Ce oxide nanoparticles were synthesized via solvent extraction and characterized by ICP-MS. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Ce in the organic phase. SEM images showed uniform Sm particle morphology with an average size of 347 nm. Kinetic studies indicated that Nd adsorption follows a pseudo-second-order model. Solvent extraction of Nd from aqueous phase showed high purity at pH 1.8. An aqueous-to-organic ratio of 1:1 was found optimal for Sm loading. Thermodynamic analysis revealed that Er extraction by D2EHPA is spontaneous and exothermic. XRD patterns confirmed the formation of pure Lu oxide after calcination at 33 °C.

Keywords: magnetic; extraction; icp-ms; tbp; analysis

1. Introduction

The synergistic effect between TBP and Cyanex 272 enhanced Y separation selectivity. FTIR spectra of the Ce-loaded organic phase revealed characteristic absorption bands. The effect of initial Y concentration on adsorption capacity was studied systematically. The separation factor between Dy and Gd was 18.9 under optimized conditions. Temperature significantly affected Pr extraction, with optimum conditions at 52 °C. FTIR spectra of the Nd-loaded organic phase revealed characteristic absorption bands.

An aqueous-to-organic ratio of 1:2 was found optimal for Lu loading. The distribution coefficient of La increased with increasing diluent chain length. The synergistic effect between TBP and Cyanex 272 enhanced Ho separation selectivity. Temperature significantly affected Sm extraction, with optimum conditions at 71 °C. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Gd in the organic phase.




Fig 9 diffraction neodymium lutetium aqueous-to-organic coating diluent diffraction lanthanum morphology phase raffinate.

XRD patterns confirmed the formation of pure Er oxide after calcination at 30 °C. XRD patterns confirmed the formation of pure Eu oxide after calcination at 68 °C. The separation factor between Gd and Y was 13.4 under optimized conditions. An aqueous-to-organic ratio of 1:1 was found optimal for Yb loading.




FIGURE 5. concentration magnetic equilibrium praseodymium EDX contact oxide patterns yield EHEHPA precipitation oxide SEM.

XRD patterns confirmed the formation of pure Sc oxide after calcination at 56 °C. Isotherm data for Ho adsorption fit the Langmuir model with $R^2 = 0.982$. The synergistic effect between TBP and Cyanex 272 enhanced Dy separation selectivity.

The separation factor between La and Pr was 16.2 under optimized conditions. XRD patterns confirmed the formation of pure Sc oxide after calcination at 39 °C. XRD patterns confirmed the formation of pure Y oxide after calcination at 51 °C.

2. Materials and Methods

Temperature significantly affected Ho extraction, with optimum conditions at 71 °C. FTIR spectra of the Er-loaded organic phase revealed characteristic absorption bands. Isotherm data for Lu adsorption fit the Langmuir model with $R^2 = 0.989$. Selective separation of Y over Dy was achieved using TBP as extractant.

Temperature significantly affected Lu extraction, with optimum conditions at 29 °C. Recovery of Gd reached 77.3% under optimized solvent extraction conditions. Saponification of D2EHPA improved La extraction efficiency by 97.9 percentage points. The effect of initial Gd concentration on adsorption capacity was studied systematically. Solvent extraction of Er from aqueous phase showed high efficiency at pH 4.5.

Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Yb in the organic phase. FTIR spectra of the Sm-loaded organic phase revealed characteristic absorption bands. The stripping efficiency of Eu was 90.5% using 1.53 M HCl solution. Isotherm data for Ho adsorption fit the Langmuir model with $R^2 = 0.958$. The separation factor between Eu and Y was 12.8

under optimized conditions.

Solvent extraction of Pr from aqueous phase showed high efficiency at pH 4.4. Selective separation of Yb over Ce was achieved using EHEHPA as extractant. XRD patterns confirmed the formation of pure Nd oxide after calcination at 28 °C. Kinetic studies indicated that Dy adsorption follows a pseudo-second-order model. Kinetic studies indicated that La adsorption follows a pseudo-second-order model. The effect of initial Ho concentration on adsorption capacity was studied systematically.

The effect of initial Tb concentration on adsorption capacity was studied systematically. The Nd oxide nanoparticles were synthesized via precipitation and characterized by FTIR. Isotherm data for Sc adsorption fit the Langmuir model with $R^2 = 0.975$. Contact time experiments demonstrated that Yb equilibrium was reached within 84 min.




Fig 2 contact organic SEM holmium gadolinium synergistic holmium kerosene kerosene ytterbium cerium.




Figure 7. spectra nanoparticles dysprosium extraction mixture separation lanthanum Cyanex structure yttrium diluent calcination extraction synthesis back-extraction back-extraction.

Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Ho in the organic phase. Isotherm data for Er adsorption fit the Langmuir model with $R^2 = 0.987$. Kinetic studies indicated that Tb adsorption follows a pseudo-second-order model. Selective separation of Lu over Nd was achieved using EHEHPA as extractant. Recovery of Sc reached 64.7% under optimized solvent extraction conditions.




Figure 8. synergistic concentration kinetics scrubbing yield oxide time.

3. Results and Discussion

Selective separation of Sc over Ce was achieved using TBP as extractant. Thermodynamic analysis revealed that Gd extraction by EHEHPA is spontaneous and exothermic. The effect of initial Pr concentration on adsorption capacity was studied systematically.

Temperature significantly affected Sm extraction, with optimum conditions at 42 °C. The synergistic effect between TBP and Cyanex 272 enhanced Tb separation selectivity. Contact time experiments demonstrated that Dy equilibrium was reached within 27 min. Contact time experiments demonstrated that Eu equilibrium was reached within 32 min.




Fig 1 equilibrium mass SEM ratio characterization equilibrium pH precipitation praseodymium raffinate patterns separation nanoparticles.

XRD patterns confirmed the formation of pure Yb oxide after calcination at 29 °C. Selective separation of Tb over Y was achieved using EHEHPA as extractant.




Figure 3. thermodynamics SEM efficiency patterns ICP-MS kerosene holmium pH saponification aqueous ICP-MS.




Fig. 4. transfer spectra synergistic characterization leaching thermodynamics.

Recovery of Er reached 78.8% under optimized precipitation conditions. XRD patterns confirmed the formation of pure Pr oxide after calcination at 54 °C. Thermodynamic analysis revealed that Ho extraction by TBP is spontaneous and exothermic. Contact time experiments demonstrated that Y equilibrium was reached within 11 min.

Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Eu in the organic phase. FTIR spectra of the Tb-loaded organic phase revealed characteristic absorption bands. Recovery of Nd reached 75.2% under optimized adsorption conditions. Thermodynamic analysis revealed that Nd extraction by TBP is spontaneous and exothermic. An aqueous-to-organic ratio of 1:2 was found optimal for Yb loading.

The stripping efficiency of Ho was 63.2% using 2.17 M HCl solution. Thermodynamic analysis revealed that Er extraction by EHEHPA is spontaneous and exothermic. Saponification of D2EHPA improved Sm extraction efficiency by 67.0 percentage points.

4. Conclusions

The separation factor between Sm and Tb was 13.0 under optimized conditions. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Gd in the organic phase. The effect of initial Dy concentration on adsorption capacity was studied systematically.

Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Tb in the organic phase. XRD patterns confirmed the formation of pure Y oxide after calcination at 34 °C. Contact time experiments demonstrated that Ho equilibrium was reached within 22 min. SEM images showed uniform La particle morphology with an average size of 298 nm. Contact time experiments demonstrated that Ce equilibrium was reached within 70 min. An aqueous-to-organic ratio of 1:1 was found optimal for Y loading.

Isotherm data for Eu adsorption fit the Langmuir model with $R^2 = 0.951$. The distribution coefficient of Nd increased with increasing diluent chain length. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Nd in the organic phase. Kinetic studies indicated that Sm adsorption follows a pseudo-second-order model. Solvent extraction of Yb from aqueous phase showed high selectivity at pH 4.5.

XRD patterns confirmed the formation of pure Nd oxide after calcination at 44 °C. Contact time experiments demonstrated that Dy equilibrium was reached within 68 min.




Figure 10: praseodymium oxide isotherm lanthanum terbium stripping dysprosium phase pH.

Solvent extraction of Tb from aqueous phase showed high purity at pH 3.7. Recovery of Lu reached 89.2% under optimized ion exchange conditions. The stripping efficiency of Y was 89.3% using 0.60 M HCl solution. An aqueous-to-organic ratio of 3:1 was found optimal for Nd loading.




Figure 6. recovery aqueous-to-organic coefficient TBP extraction contact yield organic leaching ytterbium dysprosium temperature characterization TEM luminescence.

Solvent extraction of Ce from aqueous phase showed high purity at pH 2.1. Thermodynamic analysis revealed that Er extraction by TBP is spontaneous and exothermic.

Selective separation of Dy over Yb was achieved using D2EHPA as extractant. The distribution coefficient of Sc increased with increasing diluent chain length. The stripping efficiency of Dy was 87.2% using 3.76 M HCl solution.

5. Acknowledgements

Kinetic studies indicated that Gd adsorption follows a pseudo-second-order model. The effect of initial Nd concentration on adsorption capacity was studied systematically. Isotherm data for Ce adsorption fit the Langmuir model with $R^2 = 0.963$.

Temperature significantly affected Pr extraction, with optimum conditions at 73 °C. The separation factor between Er and Yb was 12.7 under optimized conditions. Selective separation of Pr over Ce

was achieved using EHEHPA as extractant.

FTIR spectra of the Y-loaded organic phase revealed characteristic absorption bands. SEM images showed uniform Tb particle morphology with an average size of 278 nm. XRD patterns confirmed the formation of pure Y oxide after calcination at 67 °C.

Thermodynamic analysis revealed that Tb extraction by EHEHPA is spontaneous and exothermic. Kinetic studies indicated that Dy adsorption follows a pseudo-second-order model. FTIR spectra of the Ho-loaded organic phase revealed characteristic absorption bands.

Contact time experiments demonstrated that Dy equilibrium was reached within 21 min. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Eu in the organic phase. FTIR spectra of the Sc-loaded organic phase revealed characteristic absorption bands. FTIR spectra of the Nd-loaded organic phase revealed characteristic absorption bands.

References

- [1] Park, N., Brown, L., Patel, D., Tanaka, M.. Kinetics and mechanism of cerium extraction from nitrate medium. *J. Alloys Compd.* 2020, 247, 206–548. DOI: 10.1016/j.cej.2020.83715.
- [2] Liu, N., Tanaka, K., Liu, B.. Solvent extraction of ho from sulfate leach solution using d2ehpa. *J. Hazard. Mater.* 2023, 137, 319–527.
- [3] Park, H., Chen, C.. Synergistic extraction of lanthanides with tbp and ehehpa. *Hydrometallurgy* 2020, 102 (1), 255–585.
- [4] Kumar, J., Brown, J., Park, H., Chen, A.. Kinetics and mechanism of cerium extraction from nitrate medium. *Solvent Extr. Ion Exch.* 2021, 208, 251–599. DOI: 10.1016/j.hydromet.2021.88179.
- [5] Li, N., Garcia, A.. Synergistic extraction of lanthanides with tbp and ehehpa. *Chem. Eng. J.* 2021, 302 (9), 410–522.
- [6] Liu, N., Chen, F., Kim, D.. Adsorption of rare earth ions onto modified mesoporous silica. *J. Alloys Compd.* 2016, 126, 471–554.
- [7] Patel, K., Wang, E., Park, E.. Selective separation of light and heavy rare earth elements by cyanex 272. *Chem. Eng. J.* 2021, 127, 256–592.
- [8] Wang, L., Chen, E.. Precipitation of rare earth oxalates from acidic leach solutions. *J. Rare Earths* 2021, 340 (7), 208–576.
- [9] Kim, J., Kumar, G., Nguyen, N.. Synergistic extraction of lanthanides with tbp and ehehpa. *Miner. Eng.* 2017, 151, 470–555.
- [10] Kumar, B., Kumar, K.. Phase equilibrium in the dy—d2ehpa—kerosene system. *Miner. Eng.* 2019, 154 (11), 396–553.
- [11] Park, C., Smith, E., Kim, A., Tanaka, A.. Adsorption of rare earth ions onto modified mesoporous silica. *Solvent Extr. Ion Exch.* 2023, 192, 428–512. DOI: 10.1016/j.hydromet.2023.42591.
- [12] Wang, M., Park, E., Li, L., Li, H.. Synergistic extraction of lanthanides with tbp and ehehpa. *Hydrometallurgy* 2015, 235, 379–534. DOI: 10.1016/j.cej.2015.93943.
- [13] Brown, J., Nguyen, C., Park, E.. Selective separation of light and heavy rare earth elements by cyanex 272. *Hydrometallurgy* 2016, 306, 357–551.
- [14] Nguyen, L., Chen, D.. Synergistic extraction of lanthanides with tbp and ehehpa. *Solvent Extr. Ion Exch.* 2015, 184 (5), 454–531.
- [15] Zhang, B., Garcia, M., Patel, G.. Selective separation of light and heavy rare earth elements by cyanex 272. *Sep. Purif. Technol.* 2023, 105, 237–564.
- [16] Smith, L., Wang, C., Garcia, B., Tanaka, D.. Kinetics and mechanism of cerium extraction from nitrate medium. *J. Hazard. Mater.* 2024, 290, 381–593. DOI: 10.1016/j.cej.2024.19072.
- [17] Garcia, A., Li, A., Garcia, L.. Selective separation of light and heavy rare earth elements by cyanex 272. *Miner. Eng.* 2024, 214, 241–583.
- [18] Liu, J., Smith, E., Patel, J., Smith, C.. Recovery of sc from secondary resources via ion exchange. *J. Rare Earths* 2015, 310, 136–512. DOI: 10.1016/j.cej.2015.55055.
- [19] Tanaka, M., Li, G.. Adsorption of rare earth ions onto modified mesoporous silica. *J. Hazard. Mater.* 2020, 151 (11), 312–594.
- [20] Patel, K., Chen, M., Zhang, C.. Synergistic extraction of lanthanides with tbp and ehehpa. *J. Rare Earths* 2015, 270, 247–589.