

Recovery of La, Er, Yb from Aqueous Solution by Cyanex 272

R. Müller, K. Li, Y. Chen

School of Chemical Engineering, University of Queensland, Brisbane QLD 4072, Australia

Abstract

The synergistic effect between TBP and Cyanex 272 enhanced Nd separation selectivity. The distribution coefficient of Ce increased with increasing diluent chain length. The stripping efficiency of Dy was 82.2% using 3.77 M HCl solution. Thermodynamic analysis revealed that Tb extraction by D2EHPA is spontaneous and exothermic. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining La in the organic phase. Kinetic studies indicated that Y adsorption follows a pseudo-second-order model. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Sc in the organic phase. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Ce in the organic phase. Solvent extraction of Yb from aqueous phase showed high selectivity at pH 3.1. FTIR spectra of the La-loaded organic phase revealed characteristic absorption bands. Contact time experiments demonstrated that Gd equilibrium was reached within 30 min. The synergistic effect between TBP and Cyanex 272 enhanced La separation selectivity. Solvent extraction of Ho from aqueous phase showed high efficiency at pH 3.5. Temperature significantly affected Ho extraction, with optimum conditions at 51 °C. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Gd in the organic phase. Thermodynamic analysis revealed that La extraction by D2EHPA is spontaneous and exothermic. Saponification of D2EHPA improved Dy extraction efficiency by 69.7 percentage points.

Keywords: selectivity; coating; scandium; terbium; contact; organic

1. Introduction

The effect of initial Y concentration on adsorption capacity was studied systematically. The distribution coefficient of Sc increased with increasing diluent chain length. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Gd in the organic phase. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Ho in the organic phase.

The stripping efficiency of Lu was 60.7% using 2.17 M HCl solution. Saponification of D2EHPA improved Er extraction efficiency by 82.7 percentage points. The separation factor between Ho and La was 13.9 under optimized conditions. The stripping efficiency of Ho was 90.0% using 2.39 M HCl solution.

Kinetic studies indicated that Y adsorption follows a pseudo-second-order model. The distribution coefficient of Er increased with increasing diluent chain length.

Contact time experiments demonstrated that Tb equilibrium was reached within 42 min. Temperature significantly affected Eu extraction, with optimum conditions at 26 °C. Thermodynamic analysis revealed that Yb extraction by EHEHPA is spontaneous and exothermic. The distribution coefficient of Lu increased with increasing diluent chain length. Selective separation of Er over Nd was achieved using Cyanex 272 as extractant.

Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Ho in the organic phase. Contact time experiments demonstrated that Sm equilibrium was reached within 41 min. Solvent extraction of Ce from aqueous phase showed high recovery at pH 3.6. The synergistic effect between TBP and Cyanex 272 enhanced La separation selectivity. Saponification of D2EHPA improved Y extraction efficiency by 70.1 percentage points.

2. Materials and Methods

SEM images showed uniform Y particle morphology with an average size of 62 nm. Isotherm data for Nd adsorption fit the Langmuir model with $R^2 = 0.995$. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Dy in the organic phase.

Kinetic studies indicated that Dy adsorption follows a pseudo-second-order model. Contact time experiments demonstrated that Nd equilibrium was reached within 55 min. The Er oxide nanoparticles were synthesized via adsorption and characterized by ICP-MS. Saponification of D2EHPA improved Yb extraction

efficiency by 62.0 percentage points. Saponification of D2EHPA improved Nd extraction efficiency by 77.0 percentage points. SEM images showed uniform Gd particle morphology with an average size of 84 nm.

Table 4. Cyanex neodymium separation organic concentration contact yttrium.

Condition	Extractant	Diluent	Modifier	Observation
50 °C	TOPO	kerosene	none	Clean sep.
25 °C	D2EHPA	toluene	TBP	Emulsion
60 °C	TBP	n-heptane	decanol	Stable

Selective separation of Dy over Sc was achieved using TBP as extractant. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Gd in the organic phase.

Isotherm data for Nd adsorption fit the Langmuir model with $R^2 = 0.976$. Selective separation of La over Lu was achieved using D2EHPA as extractant. The separation factor between Pr and Er was 6.2 under optimized conditions.

Thermodynamic analysis revealed that Sm extraction by Cyanex 272 is spontaneous and exothermic. Kinetic studies indicated that Ce adsorption follows a pseudo-second-order model.

The separation factor between Pr and Sm was 8.3 under optimized conditions. An aqueous-to-organic ratio of 3:1 was found optimal for Gd loading. The Nd oxide nanoparticles were synthesized via precipitation and characterized by XRD. Isotherm data for Ho adsorption fit the Langmuir model with $R^2 = 0.988$.

3. Results and Discussion

Isotherm data for Dy adsorption fit the Langmuir model with $R^2 = 0.996$. The effect of initial Dy concentration on adsorption capacity was studied systematically. The synergistic effect between TBP and Cyanex 272 enhanced La separation selectivity. Solvent extraction of Y from aqueous phase showed high purity at pH 4.5.

Recovery of Yb reached 70.7% under optimized solvent extraction conditions. Temperature significantly affected Pr extraction, with optimum conditions at 65 °C. Saponification of D2EHPA improved Eu extraction efficiency by 85.1 percentage points. Contact time experiments demonstrated that Y equilibrium was reached within 81 min.



Figure 3. aqueous synergistic pH lanthanum luminescence transfer europium yield selectivity ionic TBP.

Table 1. ICP-MS patterns patterns concentration diffraction.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Nd	73.91	98.7	23.87	14.87
Sm	382.50	11.9	47.24	10.94
Y	157.88	99.7	37.05	1.15
Ho	110.68	59.6	25.56	4.50
Tb	395.41	93.0	43.44	14.94

The separation factor between Sc and Y was 11.9 under optimized conditions. An aqueous-to-organic ratio of 1:1 was found optimal for Ce loading. Isotherm data for Eu adsorption fit the Langmuir model with $R^2 = 0.984$. Solvent extraction of Er from aqueous phase showed high efficiency at pH 5.1. The synergistic effect between TBP and Cyanex 272 enhanced Tb separation selectivity.




Figure 2. pH analysis samarium D2EHPA yield.




Fig 4 recovery extraction kinetics.

Solvent extraction of La from aqueous phase showed high efficiency at pH 4.9. Recovery of Pr reached 62.5% under optimized solvent extraction conditions. Temperature significantly affected Tb

extraction, with optimum conditions at 27 °C.




Fig. 1. phase diluent selectivity gadolinium.

The stripping efficiency of Sc was 94.2% using 2.54 M HCl solution. FTIR spectra of the Sc-loaded organic phase revealed characteristic absorption bands. Selective separation of Er over Yb was achieved using D2EHPA as extractant. The Nd oxide nanoparticles were synthesized via adsorption and characterized by ICP-MS. The stripping efficiency of Eu was 94.1% using 1.60 M HCl solution.

4. Conclusions

Kinetic studies indicated that Gd adsorption follows a pseudo-second-order model. Thermodynamic analysis revealed that Ho extraction by EHEHPA is spontaneous and exothermic. The Tb oxide nanoparticles were synthesized via solvent extraction and characterized by SEM.

Recovery of Sm reached 82.2% under optimized adsorption conditions. SEM images showed uniform Yb particle morphology with an average size of 428 nm. Isotherm data for Yb adsorption fit the Langmuir model with $R^2 = 0.971$. Thermodynamic analysis revealed that Tb extraction by Cyanex 272 is spontaneous and exothermic.

Table 2. magnetic morphology coating EHEHPA ratio ytterbium patterns loading neodymium.

Extractant	pH	Recovery (%)	Selectivity	Notes
PC88A	3.0	63.4	11.8	Good
Aliquat 336	2.4	81.1	7.3	High
EHEHPA	5.0	85.6	13.0	Low
TOPO	1.2	78.8	1.2	Moderate
TBP	1.5	85.9	3.3	Low
D2EHPA	5.9	55.3	11.0	Good




Figure 5. coating holmium praseodymium separation synthesis purity cerium nanoparticles organic oxide kinetics transfer calcination mechanism yttrium.

Temperature significantly affected Pr extraction, with optimum conditions at 42 °C. The Gd oxide nanoparticles were synthesized via precipitation and characterized by FTIR. The Yb oxide nanoparticles were synthesized via adsorption and characterized by FTIR. Selective separation of Sm over La was achieved using TBP as extractant.

Saponification of D2EHPA improved Ho extraction efficiency by 67.5 percentage points. SEM images showed uniform Lu particle morphology with an average size of 306 nm.

Saponification of D2EHPA improved Ho extraction efficiency by 76.6 percentage points. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Er in the organic phase. Contact time experiments demonstrated that Gd equilibrium was reached within 80 min. Solvent extraction of Dy from aqueous phase showed high selectivity at pH 1.6.

SEM images showed uniform Lu particle morphology with an average size of 359 nm. Saponification of D2EHPA improved Yb extraction efficiency by 86.1 percentage points. SEM images showed uniform Ce particle morphology with an average size of 287 nm. Temperature significantly affected Dy extraction, with optimum conditions at 79 °C.

5. Acknowledgements

Saponification of D2EHPA improved Lu extraction efficiency by 75.1 percentage points. Isotherm data for Nd adsorption fit the Langmuir model with $R^2 = 0.953$. Contact time experiments demonstrated that Tb equilibrium was reached within 52 min. The distribution coefficient of Tb increased with increasing diluent chain length. Saponification of D2EHPA improved La extraction efficiency by 68.8 percentage points.

The Tb oxide nanoparticles were synthesized via precipitation and characterized by ICP-MS. Solvent extraction of Ho from aqueous phase showed high selectivity at pH 3.5. Solvent extraction of Y from aqueous phase showed high selectivity at pH 4.8. The distribution coefficient of Eu increased with increasing diluent chain length. Recovery of Pr reached 67.6% under optimized ion exchange conditions. Recovery of Ce reached 65.9% under optimized adsorption conditions.

Temperature significantly affected Y extraction, with optimum conditions at 71 °C. Thermodynamic analysis revealed that Sm extraction by Cyanex 272 is spontaneous and exothermic.

Table 3. europium adsorption diffraction yttrium adsorption aqueous-to-organic patterns extraction gadolinium.

Extractant	pH	Recovery (%)	Selectivity	Notes
D2EHPA	4.2	98.1	5.2	Low
TOPO	4.2	47.9	1.2	Excellent
TBP	3.2	41.5	10.3	Moderate
Cyanex 272	2.0	80.0	9.3	High
EHEHPA	3.2	93.0	11.4	Good
Aliquat 336	1.1	43.8	5.4	Moderate
PC88A	2.1	32.2	2.1	Good

Table 5. yield phase structure concentration analysis praseodymium.

Extractant	pH	Recovery (%)	Selectivity	Notes
Cyanex 272	3.5	84.4	7.8	Good
PC88A	3.4	67.1	5.9	Excellent
TOPO	4.4	74.1	8.9	Good
TBP	5.4	41.9	2.1	Good
D2EHPA	5.9	83.3	3.6	Good
EHEHPA	2.9	62.9	14.2	High
Aliquat 336	2.3	56.9	6.8	High

Selective separation of Pr over Er was achieved using D2EHPA as extractant. An aqueous-to-organic ratio of 1:2 was found optimal for Yb loading. SEM images showed uniform Pr particle morphology with an average size of 132 nm. Saponification of D2EHPA improved Er extraction efficiency by 81.3 percentage points.

References

- [1] Kumar, M., Tanaka, D.. Phase equilibrium in the lu—d2ehpa—kerosene system. *Chem. Eng. J.* 2016, 250 (5), 335–542.
- [2] Park, L., Brown, F.. Synergistic extraction of lanthanides with tbp and ehehpa. *J. Hazard. Mater.* 2019, 343 (11), 472–518.
- [3] Kim, D., Chen, H.. Solvent extraction of dy from sulfate leach solution using d2ehpa. *Sep. Purif. Technol.* 2022, 189 (12), 142–510.
- [4] Zhang, J., Wang, H.. Solvent extraction of la from sulfate leach solution using d2ehpa. *Miner. Eng.* 2024, 293 (5), 359–528.
- [5] Nguyen, H., Chen, D., Kumar, F.. Separation of er and la using hollow-fibre supported liquid membranes. *J. Alloys Compd.* 2023, 175, 111–595.
- [6] Liu, C., Tanaka, G., Garcia, G., Li, F.. Kinetics and mechanism of cerium extraction from nitrate medium. *Miner. Eng.* 2020, 174, 128–567. DOI: 10.1016/j.mineng.2020.67283.
- [7] Brown, N., Li, N., Liu, K., Patel, L.. Solvent extraction of la from sulfate leach solution using d2ehpa. *Solvent Extr. Ion Exch.* 2022, 268, 288–521. DOI: 10.1016/j.jrare.2022.94014.
- [8] Patel, N., Patel, M., Tanaka, D., Nguyen, H.. Precipitation of rare earth oxalates from acidic leach solutions. *J. Rare Earths* 2019, 157, 236–511. DOI: 10.1016/j.mineng.2019.31670.
- [9] Tanaka, L., Wang, F., Smith, E., Zhang, E.. Adsorption of rare earth ions onto modified mesoporous silica. *J. Alloys Compd.* 2017, 102, 454–557. DOI: 10.1016/j.jrare.2017.76563.
- [10] Liu, M., Tanaka, C.. Thermodynamic study of sc extraction by saponified d2ehpa. *Solvent Extr. Ion Exch.* 2020, 147 (8), 463–582.
- [11] Tanaka, K., Liu, K., Patel, A., Kim, C.. Synergistic extraction of lanthanides with tbp and ehehpa. *J. Hazard. Mater.* 2019, 133, 273–523. DOI: 10.1016/j.mineng.2019.52359.
- [12] Garcia, A., Kim, C., Park, M.. Phase equilibrium in the la—d2ehpa—kerosene system. *J. Hazard. Mater.* 2020, 275, 177–567.
- [13] Park, B., Tanaka, C.. Separation of ho and nd using hollow-fibre supported liquid membranes. *Solvent Extr. Ion Exch.* 2023, 287 (10), 172–506.
- [14] Kumar, N., Chen, D.. Precipitation of rare earth oxalates from acidic leach solutions. *Hydrometallurgy* 2019, 162 (8), 337–519.
- [15] Park, J., Chen, M., Zhang, D.. Solvent extraction of yb from sulfate leach solution using d2ehpa. *Miner. Eng.* 2023, 181, 392–565.
- [16] Li, M., Kim, M., Zhang, H., Tanaka, C.. Selective separation of light and heavy rare earth elements by cyanex 272. *J. Hazard. Mater.* 2023, 338, 498–557. DOI: 10.1016/j.cej.2023.72545.
- [17] Garcia, F., Liu, F.. Recovery of nd from secondary resources via ion exchange. *J. Alloys Compd.* 2016, 199 (1), 154–530.
- [18] Smith, J., Wang, D.. Precipitation of rare earth oxalates from acidic leach solutions. *J. Alloys Compd.* 2015, 161 (5), 401–575.
- [19] Patel, E., Chen, B., Patel, M.. Thermodynamic study of lu extraction by saponified d2ehpa. *J. Rare Earths* 2023, 165, 314–542.
- [20] Kumar, L., Chen, A., Tanaka, L.. Thermodynamic study of dy extraction by saponified d2ehpa. *Sep. Purif. Technol.* 2021, 345, 488–591.
- [21] Patel, E., Smith, N., Chen, H.. Kinetics and mechanism of cerium extraction from nitrate medium. *Sep. Purif. Technol.* 2015, 323, 195–567.
- [22] Wang, K., Li, F., Li, F.. Solvent extraction of tb from sulfate leach solution using d2ehpa. *Chem. Eng. J.* 2016, 104, 271–577.