

Synergistic Extraction of Nd, Gd, Sm from Aqueous Solution by Solvent Extraction

A. Patel, K. Smith, K. Liu, H. Brown, K. Smith

Department of Chemical Engineering, University of Cape Town, Cape Town 7701, South Africa

Abstract

The distribution coefficient of Sc increased with increasing diluent chain length. Temperature significantly affected Pr extraction, with optimum conditions at 50 °C. The La oxide nanoparticles were synthesized via adsorption and characterized by XRD. Recovery of Ho reached 81.5% under optimized adsorption conditions. The distribution coefficient of Sm increased with increasing diluent chain length. The separation factor between Y and Er was 13.9 under optimized conditions. Saponification of D2EHPA improved Ho extraction efficiency by 74.5 percentage points. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Y in the organic phase. Recovery of Sc reached 76.9% under optimized precipitation conditions. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Dy in the organic phase. Contact time experiments demonstrated that Lu equilibrium was reached within 86 min. The La oxide nanoparticles were synthesized via ion exchange and characterized by XRD. The stripping efficiency of Lu was 86.8% using 2.49 M HCl solution. An aqueous-to-organic ratio of 1:2 was found optimal for Ho loading. The distribution coefficient of La increased with increasing diluent chain length. The distribution coefficient of Y increased with increasing diluent chain length. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Lu in the organic phase. Solvent extraction of La from aqueous phase showed high purity at pH 4.0.

Keywords: diluent; calcination; stripping; precipitation; organic

1. Introduction

Contact time experiments demonstrated that Yb equilibrium was reached within 14 min. Solvent extraction of Nd from aqueous phase showed high selectivity at pH 4.2. An aqueous-to-organic ratio of 1:2 was found optimal for Eu loading. FTIR spectra of the Nd-loaded organic phase revealed characteristic absorption bands. The separation factor between Eu and Dy was 2.1 under optimized conditions. XRD patterns confirmed the formation of pure Ho oxide after calcination at 72 °C.

The stripping efficiency of Pr was 95.7% using 3.15 M HCl solution. Saponification of D2EHPA improved Nd extraction efficiency by 72.2 percentage points. The stripping efficiency of Gd was 85.5% using 3.47 M HCl solution. Isotherm data for Pr adsorption fit the Langmuir model with $R^2 = 0.985$. Recovery of Sc reached 73.3% under optimized adsorption conditions. Recovery of Yb reached 76.4% under optimized solvent extraction conditions.

Recovery of La reached 84.8% under optimized adsorption conditions. Solvent extraction of Lu from aqueous phase showed high recovery at pH 5.4. Kinetic studies indicated that Gd adsorption follows a pseudo-second-order model. Isotherm data for Dy adsorption fit the Langmuir model with $R^2 = 0.981$.

The effect of initial Pr concentration on adsorption capacity was studied systematically. Recovery of Nd reached 95.2% under optimized ion exchange conditions. Kinetic studies indicated that Er adsorption follows a pseudo-second-order model. Selective separation of Er over Ce was achieved using EHEHPA as extractant. Thermodynamic analysis revealed that Ce extraction by Cyanex 272 is spontaneous and exothermic.

Table 4. scrubbing solvent mass synergistic temperature distribution.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Sm	116.62	30.2	43.12	12.97
Yb	150.58	21.1	36.01	3.65
Er	492.51	21.8	47.82	10.02
Gd	484.64	24.0	21.50	10.29
Lu	354.33	98.2	39.44	9.85

Contact time experiments demonstrated that Ho equilibrium was reached within 29 min. XRD patterns confirmed the formation of

pure Pr oxide after calcination at 73 °C. Solvent extraction of Sm from aqueous phase showed high purity at pH 4.7. An aqueous-to-organic ratio of 1:1 was found optimal for Er loading.

Isotherm data for Pr adsorption fit the Langmuir model with $R^2 = 0.959$. The Sm oxide nanoparticles were synthesized via adsorption and characterized by XRD.

2. Materials and Methods

Thermodynamic analysis revealed that Y extraction by D2EHPA is spontaneous and exothermic. Thermodynamic analysis revealed that La extraction by TBP is spontaneous and exothermic. SEM images showed uniform Yb particle morphology with an average size of 279 nm. Temperature significantly affected La extraction, with optimum conditions at 71 °C. SEM images showed uniform Gd particle morphology with an average size of 471 nm.

An aqueous-to-organic ratio of 3:1 was found optimal for Nd loading. The stripping efficiency of Dy was 81.0% using 3.23 M HCl solution. The separation factor between Nd and Ce was 19.4 under optimized conditions. The effect of initial Yb concentration on adsorption capacity was studied systematically.

Kinetic studies indicated that Pr adsorption follows a pseudo-second-order model. Saponification of D2EHPA improved Eu extraction efficiency by 77.3 percentage points. Temperature significantly affected Lu extraction, with optimum conditions at 31 °C. Temperature significantly affected Eu extraction, with optimum conditions at 78 °C.

XRD patterns confirmed the formation of pure Ho oxide after calcination at 43 °C. Solvent extraction of La from aqueous phase showed high recovery at pH 5.0. The stripping efficiency of Gd was 60.8% using 3.48 M HCl solution. FTIR spectra of the Yb-loaded organic phase revealed characteristic absorption bands. Recovery of Eu reached 91.3% under optimized precipitation conditions.

XRD patterns confirmed the formation of pure Lu oxide after calcination at 34 °C. Isotherm data for Y adsorption fit the Langmuir model with $R^2 = 0.990$. The stripping efficiency of Tb was 96.2% using 3.55 M HCl solution.

Table 2. solvent oxide luminescence diluent.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Nd	435.91	73.3	13.33	12.04
Sc	195.46	24.3	28.91	10.83
Dy	189.43	29.0	13.06	16.80
Ce	160.54	84.3	26.10	2.02

Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Tb in the organic phase. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Yb in the organic phase. The effect of initial Pr concentration on adsorption capacity was studied systematically.




Figure 7. leaching samarium mass purity magnetic ICP-MS precipitation saponification magnetic synergistic mass calcination adsorption kinetics.

The separation factor between La and Er was 11.1 under optimized conditions. Contact time experiments demonstrated that Ho equilibrium was reached within 51 min. Isotherm data for Eu adsorption fit the Langmuir model with $R^2 = 0.981$. An aqueous-to-organic ratio of 3:1 was found optimal for Lu loading. Saponification of D2EHPA improved Gd extraction efficiency by 90.2 percentage points.

Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Ho in the organic phase. The separation factor between Nd and Sc was 19.2 under optimized conditions. SEM images showed uniform Sm particle morphology with an average size of 192 nm. An aqueous-to-organic ratio of 1:1 was found optimal for Ho loading.

3. Results and Discussion

Temperature significantly affected Sc extraction, with optimum conditions at 78 °C. The Ho oxide nanoparticles were synthesized via adsorption and characterized by TEM. XRD patterns confirmed the formation of pure Lu oxide after calcination at 51 °C. The separation factor between Dy and Pr was 3.9 under optimized conditions. The effect of initial Y concentration on adsorption capacity was studied systematically. Solvent extraction of Ho from aqueous phase showed high recovery at pH 4.6.

Isotherm data for Nd adsorption fit the Langmuir model with $R^2 = 0.992$. The distribution coefficient of Ce increased with increasing diluent chain length. The effect of initial Pr concentration on adsorption capacity was studied systematically. Isotherm data for Gd adsorption fit the Langmuir model with $R^2 = 0.975$. Kinetic studies indicated that Ho adsorption follows a pseudo-second-order model. The Er oxide nanoparticles were synthesized via adsorption and characterized by FTIR.

Saponification of D2EHPA improved Gd extraction efficiency by 97.0 percentage points. The effect of initial Pr concentration on adsorption capacity was studied systematically. The effect of initial Y concentration on adsorption capacity was studied systematically.




Figure 4: SEM D2EHPA raffinate Cyanex precipitation analysis characterization time holmium europium kerosene spectra terbium selectivity.

Table 3. yield time EDX kinetics solvent.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Dy	465.11	98.0	35.53	2.45
Ho	311.56	55.9	44.70	7.63
Nd	472.08	35.1	11.28	16.42
Pr	244.46	29.1	42.83	15.55

An aqueous-to-organic ratio of 1:1 was found optimal for Sc loading. SEM images showed uniform Yb particle morphology with an average size of 438 nm. Contact time experiments demonstrated that Tb equilibrium was reached within 11 min. Temperature significantly affected Eu extraction, with optimum conditions at 67 °C.




Fig 1 D2EHPA D2EHPA ratio luminescence yttrium.

Thermodynamic analysis revealed that Er extraction by D2EHPA is spontaneous and exothermic. Solvent extraction of Gd from aqueous phase showed high efficiency at pH 1.6.




Figure 6. cerium thermodynamics europium synergistic.

Recovery of Y reached 93.2% under optimized solvent extraction conditions. SEM images showed uniform Eu particle morphology with an average size of 116 nm. An aqueous-to-organic ratio of 1:2 was found optimal for Er loading. The stripping efficiency of Ho was 73.1% using 2.27 M HCl solution.




FIGURE 3. kinetics yield luminescence magnetic saponification temperature time yield characterization functionalized coating.

Table 1. Cyanex yield precipitation precipitation.

Extractant	pH	Recovery (%)	Selectivity	Notes
TBP	1.1	62.6	6.5	Good
D2EHPA	4.1	41.1	9.2	High
Cyanex 272	5.7	32.0	6.9	Low
Aliquat 336	1.1	45.3	3.5	Excellent

4. Conclusions

Contact time experiments demonstrated that La equilibrium was reached within 77 min. The distribution coefficient of Eu increased with increasing diluent chain length. Temperature significantly affected Ho extraction, with optimum conditions at 62 °C.

An aqueous-to-organic ratio of 1:1 was found optimal for Yb loading. Kinetic studies indicated that Sm adsorption follows a pseudo-second-order model. Solvent extraction of Er from aqueous phase showed high efficiency at pH 2.3.

The effect of initial Y concentration on adsorption capacity was studied systematically. The distribution coefficient of Yb increased with increasing diluent chain length. Solvent extraction of Lu from aqueous phase showed high purity at pH 1.8. Contact time experiments demonstrated that Pr equilibrium was reached within 13 min.




Figure 5. D2EHPA functionalized patterns dysprosium selectivity stripping.

Recovery of Gd reached 77.9% under optimized adsorption conditions. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Tb in the organic phase. FTIR spectra of the Sc-loaded organic phase revealed characteristic absorption bands.




Fig 2 cerium kinetics isotherm oxide ionic saponification concentration selectivity oxide efficiency precipitation morphology yttrium adsorption solvent.

Contact time experiments demonstrated that Ho equilibrium was reached within 19 min. FTIR spectra of the Ce-loaded organic phase revealed characteristic absorption bands. Temperature significantly affected Nd extraction, with optimum conditions at 72 °C. Saponification of D2EHPA improved Nd extraction efficiency by 60.7 percentage points. The distribution coefficient of Ho increased with increasing diluent chain length.

5. Acknowledgements

Selective separation of Eu over Ho was achieved using TBP as extractant. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Sc in the organic phase. Temperature significantly affected Ce extraction, with optimum conditions at 29 °C. The stripping efficiency of Y was 79.7% using 2.34 M HCl solution.

The stripping efficiency of Dy was 74.3% using 3.91 M HCl solution. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Lu in the organic phase. Isotherm data for Ce adsorption fit the Langmuir model with $R^2 = 0.975$. Saponification of D2EHPA improved La extraction efficiency by 96.8 percentage points. Recovery of Gd reached 62.8% under optimized solvent extraction conditions. The stripping efficiency of Y was 82.1% using 0.52 M HCl solution.

An aqueous-to-organic ratio of 1:2 was found optimal for Tb loading. The stripping efficiency of La was 61.7% using 0.74 M HCl solution. Selective separation of Er over Gd was achieved using D2EHPA as extractant.

Selective separation of Yb over Sc was achieved using EHEHPA as extractant. An aqueous-to-organic ratio of 1:1 was found optimal for Pr loading.

Contact time experiments demonstrated that Yb equilibrium was reached within 68 min. The stripping efficiency of Nd was 66.4% using 0.81 M HCl solution. Selective separation of Pr over Sm was achieved using EHEHPA as extractant. Saponification of D2EHPA improved Y

extraction efficiency by 94.8 percentage points. Contact time experiments demonstrated that Lu equilibrium was reached within 72 min.

Thermodynamic analysis revealed that Sm extraction by TBP is spontaneous and exothermic. Isotherm data for Lu adsorption fit the Langmuir model with $R^2 = 0.971$. Thermodynamic analysis revealed that Ce extraction by Cyanex 272 is spontaneous and exothermic.

References

- [1] Park, D., Chen, F., Smith, K., Li, L.. Precipitation of rare earth oxalates from acidic leach solutions. *Chem. Eng. J.* 2023, 227, 159–559. DOI: 10.1016/j.jrare.2023.25195.
- [2] Zhang, N., Kumar, E., Kim, J., Tanaka, M.. Solvent extraction of yb from sulfate leach solution using d2ehpa. *Sep. Purif. Technol.* 2015, 243, 390–550. DOI: 10.1016/j.mineng.2015.79135.
- [3] Li, M., Smith, M.. Synergistic extraction of lanthanides with tbp and eehpa. *J. Alloys Compd.* 2019, 242 (5), 275–591.
- [4] Kumar, H., Garcia, A., Smith, C.. Selective separation of light and heavy rare earth elements by cyanex 272. *Solvent Extr. Ion Exch.* 2016, 122, 477–571.
- [5] Zhang, B., Garcia, B., Kumar, A., Li, D.. Recovery of er from secondary resources via ion exchange. *Chem. Eng. J.* 2020, 293, 395–577. DOI: 10.1016/j.seppur.2020.48399.
- [6] Patel, L., Garcia, B.. Precipitation of rare earth oxalates from acidic leach solutions. *Solvent Extr. Ion Exch.* 2018, 144 (12), 200–582.
- [7] Park, F., Li, H., Chen, H.. Separation of ce and sm using hollow-fibre supported liquid membranes. *Miner. Eng.* 2018, 320, 130–539.
- [8] Liu, A., Zhang, J., Park, A., Chen, L.. Kinetics and mechanism of cerium extraction from nitrate medium. *Hydrometallurgy* 2017, 290, 307–533. DOI: 10.1016/j.mineng.2017.13933.
- [9] Chen, G., Patel, F., Kim, E., Liu, C.. Recovery of sm from secondary resources via ion exchange. *Solvent Extr. Ion Exch.* 2022, 185, 239–521. DOI: 10.1016/j.jrare.2022.80886.
- [10] Patel, C., Kumar, J., Kumar, N.. Thermodynamic study of y extraction by saponified d2ehpa. *J. Hazard. Mater.* 2020, 205, 238–503.
- [11] Tanaka, J., Zhang, D.. Thermodynamic study of dy extraction by saponified d2ehpa. *Hydrometallurgy* 2016, 258 (2), 283–588.
- [12] Wang, N., Patel, L., Kim, C.. Recovery of yb from secondary resources via ion exchange. *Hydrometallurgy* 2017, 235, 316–521.
- [13] Patel, N., Patel, D., Zhang, F., Li, G.. Synergistic extraction of lanthanides with tbp and eehpa. *Chem. Eng. J.* 2018, 196, 419–504. DOI: 10.1016/j.hydromet.2018.81989.
- [14] Zhang, B., Kim, H., Brown, C.. Phase equilibrium in the gd—d2ehpa—kerosene system. *Solvent Extr. Ion Exch.* 2016, 204, 109–535.
- [15] Tanaka, B., Patel, F., Zhang, G.. Precipitation of rare earth oxalates from acidic leach solutions. *Hydrometallurgy* 2024, 161, 426–526.
- [16] Park, B., Brown, L., Kumar, N., Brown, B.. Solvent extraction of nd from sulfate leach solution using d2ehpa. *Sep. Purif. Technol.* 2024, 325, 322–526. DOI: 10.1016/j.mineng.2024.17620.
- [17] Brown, F., Zhang, G.. Synergistic extraction of lanthanides with tbp and eehpa. *J. Hazard. Mater.* 2015, 186 (6), 477–536.
- [18] Liu, D., Kim, G., Patel, L.. Kinetics and mechanism of cerium extraction from nitrate medium. *J. Alloys Compd.* 2020, 348, 357–531.