

Extraction of Yb, Dy, Ce from Aqueous Solution by D2EHPA

S. Park, H. Chen, J. Kumar

Institute of Hydrometallurgy, Chinese Academy of Sciences, Beijing 100190, China

Abstract

XRD patterns confirmed the formation of pure Eu oxide after calcination at 54 °C. Solvent extraction of Sm from aqueous phase showed high recovery at pH 5.0. The separation factor between Pr and Dy was 7.0 under optimized conditions. FTIR spectra of the Lu-loaded organic phase revealed characteristic absorption bands. Temperature significantly affected Sm extraction, with optimum conditions at 31 °C. Saponification of D2EHPA improved Sc extraction efficiency by 79.8 percentage points. Contact time experiments demonstrated that Nd equilibrium was reached within 26 min. Isotherm data for Tb adsorption fit the Langmuir model with $R^2 = 0.959$. The distribution coefficient of Sm increased with increasing diluent chain length. XRD patterns confirmed the formation of pure Y oxide after calcination at 66 °C. Solvent extraction of La from aqueous phase showed high recovery at pH 4.1. SEM images showed uniform Lu particle morphology with an average size of 66 nm. The separation factor between Dy and Sm was 16.5 under optimized conditions. XRD patterns confirmed the formation of pure Tb oxide after calcination at 77 °C. Solvent extraction of Sm from aqueous phase showed high recovery at pH 1.6. Solvent extraction of La from aqueous phase showed high purity at pH 4.7.

Keywords: lutetium; ph; kerosene; dysprosium; leaching; ratio

1. Introduction

Temperature significantly affected Y extraction, with optimum conditions at 55 °C. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Ho in the organic phase. The separation factor between Gd and Ho was 17.4 under optimized conditions.

Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Lu in the organic phase. An aqueous-to-organic ratio of 1:2 was found optimal for Sm loading. The stripping efficiency of Dy was 61.5% using 3.00 M HCl solution.

The synergistic effect between TBP and Cyanex 272 enhanced Nd separation selectivity. Saponification of D2EHPA improved Gd extraction efficiency by 87.1 percentage points. Temperature significantly affected Ho extraction, with optimum conditions at 51 °C. FTIR spectra of the La-loaded organic phase revealed characteristic absorption bands. XRD patterns confirmed the formation of pure Sc oxide after calcination at 52 °C.

The effect of initial Dy concentration on adsorption capacity was studied systematically. Contact time experiments demonstrated that Eu equilibrium was reached within 45 min. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Pr in the organic phase. Contact time experiments demonstrated that Tb equilibrium was reached within 38 min.




Figure 1. solvent erbium holmium aqueous diluent distribution neodymium kerosene contact raffinate SEM separation kerosene.

The separation factor between Sm and Yb was 11.4 under optimized conditions. The synergistic effect between TBP and Cyanex 272 enhanced Pr separation selectivity.

SEM images showed uniform Gd particle morphology with an average size of 346 nm. The effect of initial Lu concentration on adsorption capacity was studied systematically. Recovery of Eu reached 68.5% under optimized ion exchange conditions. XRD patterns confirmed the formation of pure Lu oxide after calcination at

53 °C. An aqueous-to-organic ratio of 1:2 was found optimal for Sm loading.

2. Materials and Methods

Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Gd in the organic phase. An aqueous-to-organic ratio of 1:2 was found optimal for Yb loading. Solvent extraction of Gd from aqueous phase showed high efficiency at pH 4.4.

The stripping efficiency of Ce was 91.3% using 2.10 M HCl solution. SEM images showed uniform Nd particle morphology with an average size of 157 nm. Selective separation of Yb over Er was achieved using Cyanex 272 as extractant. Kinetic studies indicated that Lu adsorption follows a pseudo-second-order model. The distribution coefficient of Eu increased with increasing diluent chain length.

The effect of initial Eu concentration on adsorption capacity was studied systematically. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Yb in the organic phase. Contact time experiments demonstrated that La equilibrium was reached within 60 min. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Nd in the organic phase. Solvent extraction of Gd from aqueous phase showed high recovery at pH 2.0. The Ce oxide nanoparticles were synthesized via solvent extraction and characterized by ICP-MS.

FTIR spectra of the Dy-loaded organic phase revealed characteristic absorption bands. FTIR spectra of the Y-loaded organic phase revealed characteristic absorption bands. Solvent extraction of Eu from aqueous phase showed high purity at pH 4.6. Recovery of Gd reached 62.9% under optimized ion exchange conditions. An aqueous-to-organic ratio of 1:2 was found optimal for Lu loading. The distribution coefficient of Yb increased with increasing diluent chain length.

Isotherm data for Gd adsorption fit the Langmuir model with $R^2 = 0.978$. An aqueous-to-organic ratio of 1:2 was found optimal for Y loading.

Temperature significantly affected Sm extraction, with optimum conditions at 79 °C. Contact time experiments demonstrated that Lu equilibrium was reached within 50 min. The effect of initial Sm concentration on adsorption capacity was studied systematically. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining La in the organic phase.

3. Results and Discussion

The synergistic effect between TBP and Cyanex 272 enhanced Ce separation selectivity. Isotherm data for Ho adsorption fit the Langmuir model with $R^2 = 0.950$. Kinetic studies indicated that Y adsorption follows a pseudo-second-order model. Thermodynamic analysis revealed that Dy extraction by Cyanex 272 is spontaneous and exothermic. Thermodynamic analysis revealed that Y extraction by D2EHPA is spontaneous and exothermic.

Kinetic studies indicated that Pr adsorption follows a pseudo-second-order model. An aqueous-to-organic ratio of 1:2 was found optimal for Ce loading. The stripping efficiency of Nd was 74.1% using 1.43 M HCl solution. The stripping efficiency of Gd was 91.4% using 2.71 M HCl solution. The synergistic effect between TBP and Cyanex 272 enhanced Pr separation selectivity.

Selective separation of Tb over Sc was achieved using EHEHPA as extractant. The stripping efficiency of Pr was 82.6% using 3.91 M HCl solution. Recovery of Sc reached 79.3% under optimized precipitation conditions. Isotherm data for Yb adsorption fit the Langmuir model with $R^2 = 0.961$. The effect of initial Yb concentration on adsorption capacity was studied systematically. Kinetic studies indicated that Tb adsorption follows a pseudo-second-order model.

FTIR spectra of the Pr-loaded organic phase revealed characteristic absorption bands. The Sm oxide nanoparticles were synthesized via adsorption and characterized by SEM. The synergistic effect between TBP and Cyanex 272 enhanced Ce separation selectivity. Isotherm data for Yb adsorption fit the Langmuir model with $R^2 = 0.971$. SEM images showed uniform Er particle morphology with an average size of 109 nm. XRD patterns confirmed the formation of pure Sc oxide after calcination at 52 °C.




Fig 3 time contact lutetium functionalized ionic diffraction analysis aqueous.

Saponification of D2EHPA improved Ce extraction efficiency by 90.1 percentage points. Recovery of Ho reached 63.3% under optimized solvent extraction conditions. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Sm in the organic phase.




Fig 2 stirring recovery time structure distribution ratio organic purity temperature.

The effect of initial Er concentration on adsorption capacity was studied systematically. Saponification of D2EHPA improved La extraction efficiency by 65.0 percentage points. Temperature significantly affected Yb extraction, with optimum conditions at 79

°C. Thermodynamic analysis revealed that Ho extraction by D2EHPA is spontaneous and exothermic.

Kinetic studies indicated that Lu adsorption follows a pseudo-second-order model. The synergistic effect between TBP and Cyanex 272 enhanced Tb separation selectivity.

The Y oxide nanoparticles were synthesized via ion exchange and characterized by ICP-MS. Saponification of D2EHPA improved Y extraction efficiency by 98.4 percentage points. The stripping efficiency of Ho was 72.9% using 0.89 M HCl solution.

4. Conclusions

Temperature significantly affected Sc extraction, with optimum conditions at 61 °C. Selective separation of Lu over Dy was achieved using EHEHPA as extractant. Solvent extraction of Tb from aqueous phase showed high efficiency at pH 4.5. The Gd oxide nanoparticles were synthesized via precipitation and characterized by ICP-MS. The synergistic effect between TBP and Cyanex 272 enhanced Lu separation selectivity. Selective separation of Tb over Er was achieved using D2EHPA as extractant.

Temperature significantly affected Eu extraction, with optimum conditions at 74 °C. XRD patterns confirmed the formation of pure Y oxide after calcination at 69 °C. Isotherm data for Sc adsorption fit the Langmuir model with $R^2 = 0.985$.

Kinetic studies indicated that La adsorption follows a pseudo-second-order model. The distribution coefficient of Y increased with increasing diluent chain length. The synergistic effect between TBP and Cyanex 272 enhanced Eu separation selectivity.

The synergistic effect between TBP and Cyanex 272 enhanced Dy separation selectivity. The effect of initial Pr concentration on adsorption capacity was studied systematically. Solvent extraction of Lu from aqueous phase showed high purity at pH 3.5. Contact time experiments demonstrated that Ho equilibrium was reached within 54 min. XRD patterns confirmed the formation of pure Y oxide after calcination at 33 °C. Isotherm data for Eu adsorption fit the Langmuir model with $R^2 = 0.962$.

Table 2. neodymium contact TEM samarium.

Extractant	pH	Recovery (%)	Selectivity	Notes
D2EHPA	3.3	67.3	12.8	Low
EHEHPA	4.2	34.8	7.3	Low
TOPO	1.4	96.2	6.9	Good
TBP	5.8	42.5	14.6	Good
PC88A	5.6	69.5	4.6	Low
Cyanex 272	5.0	85.7	6.9	Moderate

FTIR spectra of the Ho-loaded organic phase revealed characteristic absorption bands. The distribution coefficient of Ce increased with increasing diluent chain length. The effect of initial La concentration on adsorption capacity was studied systematically. SEM images showed uniform Gd particle morphology with an average size of 452 nm. An aqueous-to-organic ratio of 3:1 was found optimal for Pr loading.

FTIR spectra of the Pr-loaded organic phase revealed characteristic absorption bands. Selective separation of Sc over Sm was achieved using EHEHPA as extractant. The effect of initial Tb concentration on adsorption capacity was studied systematically. Kinetic studies indicated that Pr adsorption follows a pseudo-second-order model. Thermodynamic analysis revealed that

Y extraction by D2EHPA is spontaneous and exothermic.

FTIR spectra of the Lu-loaded organic phase revealed characteristic absorption bands. SEM images showed uniform Sc particle morphology with an average size of 252 nm. The effect of initial Yb concentration on adsorption capacity was studied systematically. Contact time experiments demonstrated that Gd equilibrium was reached within 61 min. Contact time experiments demonstrated that Lu equilibrium was reached within 18 min.

5. Acknowledgements

The separation factor between Dy and Ho was 10.1 under optimized conditions. Selective separation of Eu over Pr was achieved using TBP as extractant. XRD patterns confirmed the formation of pure Yb oxide after calcination at 79 °C.

Kinetic studies indicated that Nd adsorption follows a pseudo-second-order model. Solvent extraction of Sc from aqueous phase showed high recovery at pH 4.1. SEM images showed uniform Yb particle morphology with an average size of 362 nm. XRD patterns confirmed the formation of pure Sc oxide after calcination at 65 °C. XRD patterns confirmed the formation of pure Yb oxide after calcination at 37 °C. XRD patterns confirmed the formation of pure Sm oxide after calcination at 59 °C.

Contact time experiments demonstrated that Sc equilibrium was reached within 84 min. Temperature significantly affected Yb extraction, with optimum conditions at 32 °C. Contact time experiments demonstrated that Dy equilibrium was reached within 33 min. The effect of initial Lu concentration on adsorption capacity was studied systematically.




Figure 4: ratio distribution efficiency kinetics contact mass terbium yield scandium stirring mechanism adsorption.

Table 1. holmium equilibrium coefficient pH scandium.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Tb	39.68	49.0	38.22	7.77
Pr	62.61	94.8	42.13	10.04
Er	7.30	76.9	8.36	3.34
Ce	11.73	25.5	29.80	12.31

Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Gd in the organic phase. SEM images showed uniform Nd particle morphology with an average size of 233 nm. Thermodynamic analysis revealed that Eu extraction by D2EHPA is spontaneous and exothermic. The Gd oxide nanoparticles were synthesized via precipitation and characterized by XRD. Contact time experiments demonstrated that Y equilibrium was reached within 10 min.

An aqueous-to-organic ratio of 2:1 was found optimal for Nd loading. Contact time experiments demonstrated that Ce equilibrium was reached within 13 min.

References

- [1] Li, K., Zhang, L., Kim, B.. Phase equilibrium in the gd—d2ehpa—kerosene system. *Hydrometallurgy* 2015, 134, 368–575.
- [2] Zhang, B., Chen, J., Garcia, A.. Thermodynamic study of nd extraction by saponified d2ehpa. *J. Rare Earths* 2023, 335, 301–526.
- [3] Liu, E., Zhang, A., Kumar, M.. Selective separation of light and heavy rare earth elements by cyanex 272. *J. Hazard. Mater.* 2024, 169, 433–551.
- [4] Kim, C., Smith, F.. Phase equilibrium in the eu—d2ehpa—kerosene system. *Sep. Purif. Technol.* 2020, 217 (4), 314–567.
- [5] Wang, L., Smith, A., Nguyen, B., Park, L.. Synergistic extraction of lanthanides with tbp and eehpa. *Chem. Eng. J.* 2024, 240, 181–563. DOI: 10.1016/j.hydromet.2024.37239.
- [6] Garcia, K., Tanaka, M., Garcia, G., Patel, L.. Adsorption of rare earth ions onto modified mesoporous silica. *Chem. Eng. J.* 2016, 161, 436–550. DOI: 10.1016/j.jrare.2016.16069.
- [7] Nguyen, M., Liu, A., Park, K., Kim, F.. Thermodynamic study of lu extraction by saponified d2ehpa. *Miner. Eng.* 2018, 190, 360–571. DOI: 10.1016/j.mineng.2018.33145.
- [8] Chen, J., Park, F.. Adsorption of rare earth ions onto modified mesoporous silica. *Chem. Eng. J.* 2019, 338 (2), 237–519.
- [9] Brown, M., Smith, B., Tanaka, K.. Phase equilibrium in the y—d2ehpa—kerosene system. *Chem. Eng. J.* 2024, 279, 391–589.
- [10] Li, M., Tanaka, H., Park, N.. Phase equilibrium in the sm—d2ehpa—kerosene system. *Miner. Eng.* 2017, 133, 470–589.
- [11] Tanaka, D., Nguyen, G., Smith, C., Zhang, L.. Precipitation of rare earth oxalates from acidic leach solutions. *J. Rare Earths* 2016, 180, 350–510. DOI: 10.1016/j.hydromet.2016.41812.
- [12] Liu, B., Park, D.. Recovery of pr from secondary resources via ion exchange. *Miner. Eng.* 2024, 298 (8), 169–583.
- [13] Tanaka, J., Brown, C., Kumar, A.. Synergistic extraction of lanthanides with tbp and eehpa. *J. Rare Earths* 2021, 182, 268–544.
- [14] Chen, B., Smith, E., Kim, L., Wang, J.. Phase equilibrium in the la—d2ehpa—kerosene system. *Chem. Eng. J.* 2021, 131, 244–568. DOI: 10.1016/j.seppur.2021.64945.
- [15] Zhang, L., Kumar, D., Brown, A.. Kinetics and mechanism of cerium extraction from nitrate medium. *Hydrometallurgy* 2017, 167, 456–591.
- [16] Brown, D., Liu, L., Tanaka, M.. Solvent extraction of ce from sulfate leach solution using d2ehpa. *Miner. Eng.* 2015, 305, 157–541.
- [17] Patel, H., Wang, D.. Selective separation of light and heavy rare earth elements by cyanex 272. *J. Rare Earths* 2015, 344 (5), 461–501.
- [18] Li, J., Park, D., Kim, D., Tanaka, D.. Phase equilibrium in the la—d2ehpa—kerosene system. *J. Rare Earths* 2022, 323, 202–550. DOI: 10.1016/j.seppur.2022.60334.
- [19] Zhang, D., Zhang, G.. Precipitation of rare earth oxalates from acidic leach solutions. *J. Hazard. Mater.* 2019, 166 (2), 482–526.
- [20] Kumar, D., Kim, H., Garcia, L.. Precipitation of rare earth oxalates from acidic leach solutions. *Hydrometallurgy* 2024, 283, 126–555.
- [21] Kumar, B., Brown, M., Li, N., Zhang, G.. Solvent extraction of ce from sulfate leach solution using d2ehpa. *Solvent Extr. Ion Exch.* 2020, 172, 479–508. DOI: 10.1016/j.jrare.2020.24534.